

L_2 DISCREPANCY OF LINEARLY DIGIT SCRAMBLED ZAREMBA POINT SETS

HENRI FAURE, FRIEDRICH PILLICHSHAMMER AND GOTTLIEB PIRSIC

ABSTRACT. We give an exact formula for the L_2 discrepancy of a class of generalized two-dimensional Hammersley point sets in base b , namely generalized Zaremba point sets. For the construction of such point sets one needs sequences of permutations of the form $\pi_l(k) = \alpha k + l \pmod{b}$ for $k, l \in \{0, \dots, b-1\}$. As a corollary we obtain a condition on these sequences which yields the best possible order of L_2 discrepancy of generalized Zaremba point sets in the sense of Roth's lower bound, with very small leading constants.

Communicated by István Berkes

1. Introduction

For a point set $\mathcal{P} = \{\mathbf{x}_1, \dots, \mathbf{x}_N\}$ of $N \geq 1$ points in the unit-square $[0, 1]^2$ the L_2 discrepancy is defined by

$$L_2(\mathcal{P}) := \left(\int_0^1 \int_0^1 |E(x; y; \mathcal{P})|^2 dx dy \right)^{1/2},$$

where the so-called *discrepancy function* is given by $E(x; y; \mathcal{P}) = A([0, x) \times [0, y); N; \mathcal{P}) - Nxy$, where $A([0, x) \times [0, y); N; \mathcal{P})$ denotes the number of indices $1 \leq M \leq N$ for which $\mathbf{x}_M \in [0, x) \times [0, y)$. The L_2 discrepancy is a quantitative measure for the irregularity of distribution of \mathcal{P} , i.e., the deviation from ideal uniform distribution. See [5, 6, 16, 18, 19, 20] for more information on L_2 discrepancy and its relation to numerical integration.

The asymptotic behavior of the minimal possible L_2 discrepancy of an N -element point set as N tends to infinity is well-known. It was first shown by

2010 Mathematics Subject Classification: 11K06, 11K38.

Keywords: L_2 discrepancy, generalized Hammersley point set, linear digit scrambling.

F. P. and G. P. are supported by the Austrian Science Foundation (FWF), Project S9609, that is part of the Austrian National Research Network "Analytic Combinatorics and Probabilistic Number Theory".

Roth [23] that there is a constant $c > 0$ with the following property: for any $N \in \mathbb{N}$ and for any N -element point set \mathcal{P} in $[0, 1]^2$ we have

$$L_2(\mathcal{P}) \geq c\sqrt{\log N}. \quad (1)$$

On the other hand it is known that there is a constant $C > 0$ such that for all $N \in \mathbb{N}$, $N \geq 2$, there exists an N -element point set \mathcal{P} in $[0, 1]^2$ with

$$L_2(\mathcal{P}) \leq C\sqrt{\log N} + O(1). \quad (2)$$

Today a lot of explicit constructions of point sets are known which achieve an upper bound of the form (2); see, for example, [1, 4, 9, 10, 11, 12, 13, 15, 17, 22, 25] and [2, 3, 5] for corresponding results in arbitrary dimension $s \geq 2$.

If one is interested in the constants c and C in (1) and (2) for large N it is advisable to study the constants \underline{c} and \bar{c} defined as

$$\underline{c} := \liminf_{N \rightarrow \infty} \inf_{\substack{\mathcal{P} \subset [0, 1]^s \\ \#\mathcal{P} = N}} \frac{L_2(\mathcal{P})}{\sqrt{\log N}} \quad \text{and} \quad \bar{c} := \limsup_{N \rightarrow \infty} \inf_{\substack{\mathcal{P} \subset [0, 1]^s \\ \#\mathcal{P} = N}} \frac{L_2(\mathcal{P})}{\sqrt{\log N}}.$$

In [14] it is shown that (1) holds with $c = 7/(216\sqrt{\log 2}) = 0.038925\dots$. Furthermore, a construction presented in [12] shows that (2) holds with $C = 0.17907\dots$ for infinitely many N . Therefore the best estimates for \underline{c} and \bar{c} known so far are

$$0.038925\dots \leq \underline{c} \leq \bar{c} \leq 0.17907\dots \quad (3)$$

The value $C = 0.17907\dots$ can be achieved with so-called generalized Hammersley point sets whose definition will be presented now.

Throughout the paper the base $b \geq 2$ is an integer and \mathfrak{S}_b is the set of all permutations of $\{0, \dots, b-1\}$.

DEFINITION 1 (generalized Hammersley point set). Let $b \geq 2$ and $n \geq 1$ be integers and let $\Sigma = (\sigma_0, \dots, \sigma_{n-1}) \in \mathfrak{S}_b^n$. For an integer $1 \leq N \leq b^n$, write $N-1 = \sum_{r=0}^{n-1} a_r(N)b^r$ in the b -adic system and define $S_b^\Sigma(N) := \sum_{r=0}^{n-1} \sigma_r(a_r(N))b^{-r-1}$. Then the *generalized two-dimensional Hammersley point set in base b* consisting of b^n points associated to Σ is defined by

$$\mathcal{H}_{b,n}^\Sigma := \left\{ \left(S_b^\Sigma(N), \frac{N-1}{b^n} \right) : 1 \leq N \leq b^n \right\}.$$

If we choose in the above definition $\sigma_i = \text{id}$ — the identity in \mathfrak{S}_b — for all $i \in \{0, \dots, n-1\}$, then we obtain the classical two-dimensional Hammersley point set in base b .

A lot of sequences of permutations $\Sigma = (\sigma_0, \dots, \sigma_{n-1}) \in \mathfrak{S}_b^n$ are known which achieve an order $O(\sqrt{\log N})$ for the L_2 discrepancy of $\mathcal{H}_{b,n}^\Sigma$, see [9, 10, 11, 12, 15, 25] and the references therein.

Here we deal with sequences of permutations of the following form: for $\sigma \in \mathfrak{S}_b$ and $l \in \{0, \dots, b-1\}$ let $\sigma_l(k) := \sigma(k) + l \pmod{b}$ for all $k \in \{0, \dots, b-1\}$. We call σ_l the *shifted permutation σ with shift l* .

DEFINITION 2 (generalized Zaremba point set). Let $\sigma \in \mathfrak{S}_b$. A generalized Hammersley point set $\mathcal{H}_{b,n}^\Sigma$ where $\Sigma \in \{\sigma_l : 0 \leq l < b\}^n$ is called a *generalized Zaremba point set*.

This terminology goes back to White [25] who considered sequences Σ of the form

$$(\text{id}_0, \text{id}_1, \dots, \text{id}_{b-1}, \text{id}_0, \text{id}_1, \dots, \text{id}_{b-1}, \dots) \quad (4)$$

of length n and who gave an exact formula for the L_2 discrepancy of the corresponding generalized Hammersley point set, which he named *Zaremba point set*. This result is generalized in [11] to arbitrary sequences $\Sigma \in \{\text{id}_l : 0 \leq l < b\}^n$. The main result in [11] states that

$$(L_2(\mathcal{H}_{b,n}^\Sigma))^2 = n \frac{(b^2 - 1)(3b^2 + 13)}{720b^2} + O(1) \quad (5)$$

whenever the permutations id_l for $0 \leq l < b$ appear with the same frequency in the sequence Σ . It is interesting that the specific order of the id_l 's is of no importance contrary to (4). This observation was already made in [15].

It is the aim of this paper to generalize the results from [11] to sequences of permutations Σ belonging to $\{\pi_l : 0 \leq l < b\}^n$, where π is a linear permutation in \mathfrak{S}_b , i.e., of the form $\pi(k) = \alpha k \pmod{b}$ for some $\alpha \in \{1, \dots, b-1\}$ with $\gcd(\alpha, b) = 1$. This generalization allows a drastic improvement of the leading factor $\frac{(b^2-1)(3b^2+13)}{720b^2}$ in Formula (5).

Doing so, we continue to explore the L_2 discrepancy of various classes of Hammersley point sets with the help of an exact formula for the discrepancy function first used in [9] and then in [10, 11, 12] (see Lemma 3 below). Apart from number theory, permutations π_l are of interest since they are widely used in quasi-Monte Carlo methods under the name *linear digit scramblings*, to improve Halton sequences and $(0, s)$ -sequences (so-called Faure sequences); see for instance [8, 18].

The paper is organized as follows. In Section 2 we present the main results of the paper and in Section 3 we provide some auxiliary results which are necessary for the proofs in Section 4. We close this introductory section with some notations which are used throughout the paper.

Basic Notations. The analysis of the L_2 discrepancy is based on special functions which have been first introduced by Faure [7] and which are defined as follows:

For $\sigma \in \mathfrak{S}_b$ let $\mathcal{Z}_b^\sigma = (\sigma(0)/b, \sigma(1)/b, \dots, \sigma(b-1)/b)$. For $h \in \{0, 1, \dots, b-1\}$ and $x \in [(k-1)/b, k/b)$, where $k \in \{1, \dots, b\}$, we define

$$\varphi_{b,h}^\sigma(x) = \begin{cases} A([0, h/b]; k; \mathcal{Z}_b^\sigma) - hx & \text{if } 0 \leq h \leq \sigma(k-1), \\ (b-h)x - A([h/b, 1]; k; \mathcal{Z}_b^\sigma) & \text{if } \sigma(k-1) < h < b, \end{cases}$$

where here for a sequence $X = (x_M)_{M \geq 1}$ we denote by $A(I; k; X)$ the number of indices $1 \leq M \leq k$ such that $x_M \in I$. Further, the function $\varphi_{b,h}^\sigma$ is extended to the reals by periodicity. Note that $\varphi_{b,0}^\sigma = 0$ for any σ and that $\varphi_{b,h}^\sigma(0) = 0$ for any $\sigma \in \mathfrak{S}_b$ and any $0 \leq h < b$.

For $r \in \mathbb{N}$ define $\varphi_b^{\sigma,(r)} := \sum_{h=0}^{b-1} (\varphi_{b,h}^\sigma)^r$ and we simply write $\varphi_b^\sigma := \varphi_b^{\sigma,(1)}$. Note that φ_b^σ is continuous (see [1, Propriété 3.3] and [7, Propriété 3.2.2]), piecewise linear on the intervals $[k/b, (k+1)/b)$ and $\varphi_b^\sigma(0) = \varphi_b^\sigma(1)$.

Further, for our purpose, we will need the integrals $\Phi_b^\sigma := (1/b) \int_0^1 \varphi_b^\sigma(x) dx$ and $\Phi_b^{\sigma,(2)} := (1/b) \int_0^1 \varphi_b^{\sigma,(2)}(x) dx$.

2. The results

First we state a generalization of [11, Theorem 1] and of [15, Theorem 1].

PROPOSITION 1. *Let $\pi \in \mathfrak{S}_b$ be linear and let $\Sigma = (\sigma_0, \dots, \sigma_{n-1}) \in \{\pi_l : 0 \leq l < b\}^n$. For $0 \leq l < b$ define $\lambda_l := \#\{0 \leq i < n : \sigma_i = \pi_l\}$. Then we have*

$$\begin{aligned} (L_2(\mathcal{H}_{b,n}^\Sigma))^2 &= \left(\sum_{l=0}^{b-1} \lambda_l \Phi_b^{\pi_l} \right)^2 + n \Phi_b^{\pi,(2)} + \sum_{l=0}^{b-1} \lambda_l \Phi_b^{\pi_l} \\ &\quad + \sum_{l=0}^{b-1} \frac{\lambda_l}{b} \left[\varphi_b^{\pi,(2)} \left(\frac{\pi^{-1}(l)}{b} \right) - 2F_b^\pi(l) - b(\Phi_b^{\pi_l})^2 \right] + O(1), \end{aligned}$$

where $F_b^\pi(l) := (1/b) \sum_{h,j=0}^{b-1} \varphi_{b,h}^\pi(\pi^{-1}(l)/b) \varphi_{b,h}^\pi(j/b)$.

The proof of this result will be presented in Section 4. Although the idea for the proof is the same as for $\pi = \text{id}$ in [11], the proof is much more sophisticated and a lot of technical difficulties must be overcome before to reach the proposed formula. A short outline is given below, just before Lemma 4.

The following result provides a choice of Σ which yields the best possible order of L_2 discrepancy with respect to Roth's lower bound (1). This result generalizes [11, Corollary 1] (see formula (5)).

L_2 DISCREPANCY OF LINEARLY DIGIT SCRAMBLED ZAREMBA POINT SETS




FIGURE 1. Comparison of $\sqrt{C_{\text{opt}}(b)/\log b}$ and $\sqrt{C_1(b)/\log b}$ for $b = 2, \dots, 50$.

THEOREM 1. *Let $\pi \in \mathfrak{S}_b$ be linear and let $\Sigma \in \{\pi_l : 0 \leq l < b\}^n$ be such that $\lambda_l = \lfloor n/b \rfloor + \theta_l$ with $\theta_l \in \{0, 1\}$ for all $0 \leq l < b$. Then we have*

$$(L_2(\mathcal{H}_{b,n}^\Sigma))^2 = n(\Phi_b^{\pi,(2)} - (\Phi_b^\pi)^2) + O(1).$$

Thus we obtain the optimal order $O(\sqrt{n})$ for the L_2 discrepancy of $\mathcal{H}_{b,n}^\Sigma$ whenever the permutations π_l for $0 \leq l < b$ appear with the same frequency in the sequence Σ , independently of the specific order of the π_l 's.

The result of Theorem 1 yields a drastic improvement of [11, Corollary 1] (see formula (5)) when we use the “optimal” linear permutation $\pi(k) = \alpha k \pmod{b}$ instead of $\pi(k) = \text{id}$ (i.e. $\alpha = 1$). For $\alpha = 1$ we have $\Phi_b^{\pi,(2)} - (\Phi_b^\pi)^2 = \frac{(b^2-1)(3b^2+13)}{720b^2} =: C_1(b)$. Let now $C_{\text{opt}}(b) := \min\{\Phi_b^{\pi,(2)} - (\Phi_b^\pi)^2 : 1 \leq \alpha < b \text{ and } \gcd(\alpha, b) = 1\}$. Using results from [12] the value $\Phi_b^{\pi,(2)} - (\Phi_b^\pi)^2$ for a given permutation π can easily be calculated, for example with MATHEMATICA. A comparison of the two quantities $C_{\text{opt}}(b)$ and $C_1(b)$ for $b = 2, \dots, 50$ can be found in Figure 1. To compare the result also with the constants \underline{c} and \bar{c} from (3) we plot $\sqrt{C_{\text{opt}}(b)/\log b}$ and $\sqrt{C_1(b)/\log b}$, respectively. Notice that the constants $C_{\text{opt}}(b)$ are very close to the analogous constants in [12], even though a bit larger.

In Theorem 1 we require b permutations to obtain the optimal order of L_2 discrepancy. The following result shows that the optimal order can be obtained with only one permutation.

THEOREM 2. *If $b \in \mathbb{N}$, $b \geq 2$, $\alpha \in \{1, \dots, b-1\}$ with $\gcd(\alpha, b) = 1$ and $l \in \{0, \dots, b-1\}$ are chosen such that $\Phi_b^{\pi_l} = 0$, then with $\Sigma = (\pi_l, \dots, \pi_l)$ we have $L_2(\mathcal{H}_{b,n}^\Sigma) = O(\sqrt{n})$.*

For $\alpha = 1$ there exist infinitely many $b \geq 2$ and corresponding l such that $\Phi_b^{\pi_l} = \Phi_b^{\text{id}_l} = 0$; see [10, Corollary 1] for a necessary and sufficient condition. Many further examples of (b, α, l) for which $\Phi_b^{\pi_l} = 0$ can be found numerically, for example using MATHEMATICA. Until now we were not able to give a characterization of those (b, α, l) which yield $\Phi_b^{\pi_l} = 0$. Finding such a characterization remains open for the moment.

3. Auxiliary results

In this section we provide the main tools for the proof of Proposition 1. For the sake of completeness, we give short hints for the proofs of lemmas concerned with shifts and already proved in [10].

For $\sigma = \text{id}$, the identity in \mathfrak{S}_b , we have

$$\varphi_{b,h}^{\text{id}}(x) = \begin{cases} (b-h)x & \text{if } x \in [0, h/b], \\ h(1-x) & \text{if } x \in [h/b, 1], \end{cases} \quad (6)$$

from which one obtains (see [9, Lemma 3] for details) that for $x \in [\frac{k}{b}, \frac{k+1}{b}]$, $0 \leq k < b$, we have

$$\varphi_b^{\text{id}}(x) = \frac{b(b-2k-1)}{2} \left(x - \frac{k}{b}\right) + \frac{k(b-k)}{2}. \quad (7)$$

Considering shifts of general permutations, we will need extensions of formulas (6) and (7). First, recall from [1, Propriété 3.4] that

$$(\varphi_{b,h}^\sigma)'(k/b + 0) = (\varphi_{b,h}^{\text{id}})'(\sigma(k)/b + 0). \quad (8)$$

(Here and later on by $f'(x+0)$ we mean the right-derivative of the function f at x .) Then, using that the functions $\varphi_{b,h}^\sigma$ are continuous and piecewise linear, it is easy to see that

$$\varphi_{b,h}^\sigma(l/b) = (1/b) \sum_{k=0}^{l-1} (\varphi_{b,h}^\sigma)'(k/b + 0), \quad (9)$$

from which we deduce $\varphi_b^\sigma(l/b) = (1/b) \sum_{k=0}^{l-1} (\varphi_b^\sigma)'(k/b + 0)$, and, using (8) and (6),

$$\varphi_b^\sigma\left(\frac{l}{b}\right) = l \frac{b-1}{2} - \sum_{i=0}^{l-1} \sigma(i). \quad (10)$$

These two last properties were proved directly in [1, Propriété 3.5 (i)]. Since φ_b^σ is linear on intervals $[l/b, (l+1)/b]$ we obtain from (10) that for all $x \in [l/b, (l+1)/b]$ we have

$$\varphi_b^\sigma(x) = b \frac{b - 2\sigma(l) - 1}{2} \left(x - \frac{l}{b} \right) + l \frac{b - 1}{2} - \sum_{i=0}^{l-1} \sigma(i). \quad (11)$$

In [12, Lemma 5] it has been shown that

$$\Phi_b^\sigma = \frac{1}{b^2} \left(\sum_{i=0}^{b-1} \sigma(i) i - b \left(\frac{b-1}{2} \right)^2 \right). \quad (12)$$

In the following we assume that $\pi \in \mathfrak{S}_b$ is linear, i.e., $\pi(k) = \alpha k \pmod{b}$ for some $\alpha \in \{1, \dots, b-1\}$ with $\gcd(\alpha, b) = 1$. In this case we have that π^{-1} is linear as well, that $\pi^{-1}(l + \pi(k)) = \pi^{-1}(l) + k \pmod{b}$ and that $(\pi_l)^{-1} = (\pi^{-1})_{b-\pi^{-1}(l)} \pmod{b}$.

LEMMA 1. *For any $0 \leq k, l < b$ we have*

$$(\varphi_b^{\pi_l})' \left(\frac{k}{b} + 0 \right) = (\varphi_b^\pi)' \left(\frac{k + \pi^{-1}(l)}{b} + 0 \right).$$

Proof. With (8) and with $\varphi_b^\sigma = \sum_{h=0}^{b-1} \varphi_{b,h}^\sigma$ we obtain

$$\begin{aligned} (\varphi_b^{\pi_l})' \left(\frac{k}{b} + 0 \right) &= (\varphi_b^{\text{id}})' \left(\frac{\pi_l(k)}{b} + 0 \right) = (\varphi_b^{\text{id}})' \left(\frac{\pi(k) + l}{b} + 0 \right) \\ &= (\varphi_b^\pi)' \left(\frac{\pi^{-1}(\pi(k) + l)}{b} + 0 \right) = (\varphi_b^\pi)' \left(\frac{k + \pi^{-1}(l)}{b} + 0 \right). \end{aligned}$$

□

The following lemma gives a relation between the functions φ_b^σ and $\varphi_b^{\sigma_l}$.

LEMMA 2. *For any $0 \leq l < b$ and $x \in [0, 1]$ we have*

$$\varphi_b^{\pi_l}(x) = \varphi_b^\pi \left(x + \frac{\pi^{-1}(l)}{b} \right) - \varphi_b^\pi \left(\frac{\pi^{-1}(l)}{b} \right). \quad (13)$$

Proof. Both sides of (13) coincide in $x = 0$, are continuous and linear on intervals of the form $[k/b, (k+1)/b]$ and also their right derivatives at k/b , $0 \leq k < b$ coincide by Lemma 1. Hence (13) is a proper equation. □

REMARK 1. For an arbitrary permutation σ , it can be shown by recursion that Lemma 2 becomes: For any $x \in [\frac{k}{b}, \frac{k+1}{b}]$, $0 \leq k < b$,

$$\varphi_b^{\sigma^l}(x) = \varphi_b^\sigma \left(x - k + \frac{\sigma^{-1}(l + \sigma(k))}{b} \right) - B_k,$$

where $B_k = \sum_{h=0}^k \varphi_b^\sigma \left(\frac{\sigma^{-1}(l + \sigma(h))}{b} \right) - \sum_{h=0}^{k-1} \varphi_b^\sigma \left(\frac{\sigma^{-1}(l + \sigma(h)) + 1}{b} \right)$. Apart from the interest of linear digit scramblings π_l in applications, the complexity of that formula is a reason for choosing, at first, linear permutations.

The following lemma provides a formula for the discrepancy function of generalized Hammersley point sets.

LEMMA 3. For integers $1 \leq \lambda, N \leq b^n$ we have

$$E \left(\frac{\lambda}{b^n}; \frac{N}{b^n}; \mathcal{H}_{b,n}^\Sigma \right) = \sum_{j=1}^n \varphi_{b,\varepsilon_j}^{\sigma_{j-1}} \left(\frac{N}{b^j} \right),$$

where the $\varepsilon_j = \varepsilon_j(\lambda, n, N)$ can be given explicitly.

A proof of this result together with formulas for $\varepsilon_j = \varepsilon_j(\lambda, n, N)$ can be found in [9, Lemma 1].

REMARK 2. Let $0 \leq x, y \leq 1$ be arbitrary. Since all points from $\mathcal{H}_{b,n}^\Sigma$ have coordinates of the form αb^{-n} for some $\alpha \in \{0, 1, \dots, b^n - 1\}$, we have

$$E(x; y; \mathcal{H}_{b,n}^\Sigma) = E(x(n); y(n); \mathcal{H}_{b,n}^\Sigma) + b^n(x(n)y(n) - xy), \quad (14)$$

where for $0 \leq x \leq 1$ we define $x(n) := \min\{\alpha b^{-n} \geq x : \alpha \in \{0, \dots, b^n\}\}$.

In the following we will give a series of lemmas with further, more involved properties of the functions $\varphi_{b,h}^\sigma$ and $\varphi_b^{\sigma,(r)}$ functions. Before going on, as stated in Section 2, we briefly outline the unfolding of the proof of Proposition 1. In Section 4, after the proof of our last lemma (Lemma 8), the proof of Proposition 1 starts with the consideration of the two-dimensional integral involving (14). This term can be easily split into three sums. The first one, Σ_1 , is the most important one, with a priori n^2 terms, and needs the whole series of lemmas (excepted Lemma 7 which is required for Σ_2) culminating in Lemma 8 which can be viewed as a discrete version of Proposition 1. The third sum Σ_3 is trivial, but Σ_2 has a lot of technical complications which require a careful analysis to be overcome. Nevertheless, the exact computation of Σ_2 is necessary in view of Theorem 1, since it contains a priori n terms. Finally, putting the three sums together yields the result of Proposition 1.

LEMMA 4. For $1 \leq N \leq b^n$, $1 \leq j_1 < \dots < j_k \leq n$ and $r_1, \dots, r_k \in \mathbb{N}$ we have

$$\sum_{\lambda=1}^{b^n} \prod_{i=1}^k \left(\varphi_{b, \varepsilon_{j_i}}^{\sigma_{j_i-1}} \left(\frac{N}{b^{j_i}} \right) \right)^{r_i} = b^{n-k} \prod_{i=1}^k \varphi_b^{\sigma_{j_i-1}, (r_i)} \left(\frac{N}{b^{j_i}} \right).$$

A proof of this result can be found in [9, Lemma 2].

LEMMA 5. For $0 \leq h, k < n$, $h \neq k$ and $0 \leq l, m < b$ we have

$$\begin{aligned} & \sum_{N=1}^{b^n} \varphi_b^{\pi l} \left(\frac{N}{b^h} \right) \varphi_b^{\pi m} \left(\frac{N}{b^k} \right) \\ &= b^n \left(b \Phi_b^\pi - \varphi_b^\pi \left(\frac{\pi^{-1}(l)}{b} \right) \right) \left(b \Phi_b^\pi - \varphi_b^\pi \left(\frac{\pi^{-1}(m)}{b} \right) \right). \end{aligned}$$

Proof. We use the abbreviation $l' = \pi^{-1}(l)$ and $m' = \pi^{-1}(m)$. Using Lemma 2 we have

$$\begin{aligned} & \sum_{N=1}^{b^n} \varphi_b^{\pi l} \left(\frac{N}{b^h} \right) \varphi_b^{\pi m} \left(\frac{N}{b^k} \right) \\ &= \sum_{N=1}^{b^n} \varphi_b^\pi \left(\frac{N}{b^h} + \frac{l'}{b} \right) \varphi_b^\pi \left(\frac{N}{b^k} + \frac{m'}{b} \right) + b^n \varphi_b^\pi \left(\frac{l'}{b} \right) \varphi_b^\pi \left(\frac{m'}{b} \right) \\ & \quad - \varphi_b^\pi \left(\frac{m'}{b} \right) \sum_{N=1}^{b^n} \varphi_b^\pi \left(\frac{N}{b^h} + \frac{l'}{b} \right) - \varphi_b^\pi \left(\frac{l'}{b} \right) \sum_{N=1}^{b^n} \varphi_b^\pi \left(\frac{N}{b^k} + \frac{m'}{b} \right). \quad (15) \end{aligned}$$

From the periodicity of φ_b^π we obtain

$$\begin{aligned} \sum_{N=1}^{b^n} \varphi_b^\pi \left(\frac{N}{b^h} + \frac{l'}{b} \right) &= b^{n-h} \sum_{N=0}^{b^h-1} \varphi_b^\pi \left(\frac{N}{b^h} + \frac{l'}{b} \right) = b^{n-h} \sum_{N=0}^{b^{h-1}-1} \sum_{t=0}^{b-1} \varphi_b^\pi \left(\frac{N}{b^h} + \frac{t}{b} \right) \\ &= b^{n-h} \sum_{N=0}^{b^{h-1}-1} b \int_0^1 \varphi_b^\pi(x) dx = b^{n+1} \Phi_b^\pi, \quad (16) \end{aligned}$$

since for fixed $0 \leq N < b^{h-1}$ we have $\sum_{t=0}^{b-1} \varphi_b^\pi \left(\frac{N}{b^h} + \frac{t}{b} \right) = b \int_0^1 \varphi_b^\pi(x) dx$ as shown in [12, Proof of Lemma 4, p. 405].

Without loss of generality we may assume that $h < k$. Then we have

$$\sum_{N=1}^{b^n} \varphi_b^\pi \left(\frac{N}{b^h} + \frac{l'}{b} \right) \varphi_b^\pi \left(\frac{N}{b^k} + \frac{m'}{b} \right)$$

$$\begin{aligned}
 &= b^{n-k} \sum_{N=0}^{b^k-1} \varphi_b^\pi \left(\frac{N}{b^h} + \frac{l'}{b} \right) \varphi_b^\pi \left(\frac{N}{b^k} + \frac{m'}{b} \right) \\
 &= b^{n-k} \sum_{N=0}^{b^{k-1}-1} \varphi_b^\pi \left(\frac{N}{b^h} + \frac{l'}{b} \right) \sum_{t=0}^{b-1} \varphi_b^\pi \left(\frac{N}{b^k} + \frac{t}{b} \right) \\
 &= b^{n-k} b \int_0^1 \varphi_b^\pi(x) dx \sum_{N=0}^{b^{k-1}-1} \varphi_b^\pi \left(\frac{N}{b^h} + \frac{l'}{b} \right) \\
 &= \int_0^1 \varphi_b^\pi(x) dx \sum_{N=0}^{b^n-1} \varphi_b^\pi \left(\frac{N}{b^h} + \frac{l'}{b} \right) \\
 &= b^n \left(\int_0^1 \varphi_b^\pi(x) dx \right)^2 = b^n (b\Phi_b^\pi)^2. \tag{17}
 \end{aligned}$$

Now the result follows from inserting (16) and (17) into (15). \square

LEMMA 6. *For $1 \leq k \leq n$ we have*

$$\sum_{N=1}^{b^n} \varphi_b^{\pi_l, (2)} \left(\frac{N}{b^k} \right) = b^n \left(b\Phi_b^{\pi, (2)} + \frac{b(b^2-1)}{36b^{2k}} + \varphi_b^{\pi, (2)} \left(\frac{\pi^{-1}(l)}{b} \right) - 2F_b^\pi(l) \right),$$

where $F_b^\pi(l) := (1/b) \sum_{h,j=0}^{b-1} \varphi_{b,h}^\pi(\pi^{-1}(l)/b) \varphi_{b,h}^\pi(j/b)$.

Proof. Again we write $l' = \pi^{-1}(l)$. We have

$$\begin{aligned}
 \varphi_b^{\pi_l, (2)} \left(\frac{N}{b^k} \right) &= \sum_{h=0}^{b-1} \left(\varphi_{b,h}^{\pi_l} \left(\frac{N}{b^k} \right) \right)^2 = \sum_{h=0}^{b-1} \left(\varphi_{b,h}^\pi \left(\frac{N}{b^k} + \frac{l'}{b} \right) - \varphi_{b,h}^\pi \left(\frac{l'}{b} \right) \right)^2 \\
 &= \varphi_b^{\pi, (2)} \left(\frac{N}{b^k} + \frac{l'}{b} \right) + \varphi_b^{\pi, (2)} \left(\frac{l'}{b} \right) - 2 \sum_{h=0}^{b-1} \varphi_{b,h}^\pi \left(\frac{N}{b^k} + \frac{l'}{b} \right) \varphi_{b,h}^\pi \left(\frac{l'}{b} \right).
 \end{aligned}$$

By using the periodicity of $\varphi_b^{\pi, (2)}$ we obtain

$$\sum_{N=1}^{b^n} \varphi_b^{\pi, (2)} \left(\frac{N}{b^k} + \frac{l'}{b} \right) = \sum_{N=1}^{b^n} \varphi_b^{\pi, (2)} \left(\frac{N}{b^k} \right) = b^n \left(b\Phi_b^{\pi, (2)} + \frac{b(b^2-1)}{36b^{2j}} \right),$$

where the last equality is [12, Lemma 4, Equation (8)].

Furthermore we have

$$\sum_{N=1}^{b^n} \sum_{h=0}^{b-1} \varphi_{b,h}^\pi \left(\frac{N}{b^k} + \frac{l'}{b} \right) \varphi_{b,h}^\pi \left(\frac{l'}{b} \right) = \sum_{h=0}^{b-1} \varphi_{b,h}^\pi \left(\frac{l'}{b} \right) \sum_{N=1}^{b^n} \varphi_{b,h}^\pi \left(\frac{N}{b^k} + \frac{l'}{b} \right).$$

Using the periodicity of $\varphi_{b,h}^\pi$, we obtain for the innermost sum

$$\begin{aligned} \sum_{N=1}^{b^n} \varphi_{b,h}^\pi \left(\frac{N}{b^k} + \frac{l'}{b} \right) &= b^{n-k} \sum_{N=0}^{b^k-1} \varphi_{b,h}^\pi \left(\frac{N}{b^k} \right) \\ &= b^{n-k} \sum_{j=0}^{b-1} \sum_{N=jb^{k-1}+1}^{(j+1)b^{k-1}} \varphi_{b,h}^\pi \left(\frac{N}{b^k} \right). \end{aligned}$$

For $j/b \leq x \leq (j+1)/b$ we have

$$\varphi_{b,h}^\pi(x) = b \left(x - \frac{j}{b} \right) \left[\varphi_{b,h}^\pi \left(\frac{j+1}{b} \right) - \varphi_{b,h}^\pi \left(\frac{j}{b} \right) \right] + \varphi_{b,h}^\pi \left(\frac{j}{b} \right).$$

Therefore we have

$$\begin{aligned} &\sum_{N=1}^{b^n} \varphi_{b,h}^\pi \left(\frac{N}{b^k} + \frac{l'}{b} \right) \\ &= b^{n-k} \sum_{j=0}^{b-1} b \left[\varphi_{b,h}^\pi \left(\frac{j+1}{b} \right) - \varphi_{b,h}^\pi \left(\frac{j}{b} \right) \right] \sum_{N=jb^{k-1}+1}^{(j+1)b^{k-1}} \left(\frac{N}{b^k} - \frac{j}{b} \right) \\ &\quad + b^{n-1} \sum_{j=0}^{b-1} \varphi_{b,h}^\pi \left(\frac{j}{b} \right) \\ &= b^{n-k} \sum_{j=0}^{b-1} \left[\varphi_{b,h}^\pi \left(\frac{j+1}{b} \right) - \varphi_{b,h}^\pi \left(\frac{j}{b} \right) \right] \frac{b+b^k}{2b} + b^{n-1} \sum_{j=0}^{b-1} \varphi_{b,h}^\pi \left(\frac{j}{b} \right) \\ &= b^{n-1} \sum_{j=0}^{b-1} \varphi_{b,h}^\pi \left(\frac{j}{b} \right). \end{aligned}$$

Hence we have

$$\sum_{h=0}^{b-1} \sum_{N=1}^{b^n} \varphi_{b,h}^\pi \left(\frac{N}{b^k} + \frac{l'}{b} \right) \varphi_{b,h}^\pi \left(\frac{l'}{b} \right) = b^{n-1} \sum_{h=0}^{b-1} \varphi_{b,h}^\pi \left(\frac{l'}{b} \right) \sum_{j=0}^{b-1} \varphi_{b,h}^\pi \left(\frac{j}{b} \right) = b^n F_b^\pi(l).$$

The result follows. \square

LEMMA 7. For $0 \leq h \leq n$ and $0 \leq l < b$ we have

$$\sum_{N=1}^{b^n} N \varphi_b^\pi \left(\frac{N}{b^h} + \frac{\pi^{-1}(l)}{b} \right) = b^{2n} \frac{b\Phi_b^\pi}{2} + b^{n+h} f_b^\pi(l) + b^n g_b^\pi(l),$$

where $g_b^\pi(l) = \frac{b}{2}\Phi_b^\pi - \frac{b^2+1}{24} + \frac{l(b-1)}{4} - \frac{1}{2}\sum_{i=0}^{l-1}\pi(i)$ and

$$\begin{aligned} f_b^\pi(l) &= -\frac{b}{2}\Phi_b^\pi + \frac{(b-1)^2(1-5b)}{24b} + \frac{1}{2b^2}\sum_{i=0}^{b-1}i^2\pi(i) + \frac{(b-1)l(l-2)}{2b} \\ &\quad - \left(l - \frac{1}{2}\right)\frac{1}{b}\sum_{i=0}^{l-1}\pi(i) + \frac{1}{b}\sum_{i=0}^{l-1}i\pi(i). \end{aligned}$$

Proof. Again we write $l' = \pi^{-1}(l)$. Splitting up the range of summation yields

$$\sum_{N=1}^{b^n} N\varphi_b^\pi\left(\frac{N}{b^h} + \frac{l'}{b}\right) = \sum_{k=0}^{b^{n-h+1}-1} \sum_{N=kb^{h-1}+1}^{(k+1)b^{h-1}} N\varphi_b^\pi\left(\frac{N}{b^h} + \frac{l'}{b}\right).$$

For $0 \leq k < b^{n-h+1}$ let $k = qb + r$ with integers $0 \leq r < b$ and $0 \leq q < b^{n-h}$. Then for $kb^{h-1} + 1 \leq N \leq (k+1)b^{h-1}$ we have $r/b \leq N/b^h - q \leq (r+1)/b$. Hence,

- if $0 \leq r < b-l$, then

$$N/b^h - q + l'/b \in \left[\frac{r+l'}{b}, \frac{r+l'+1}{b}\right] \subseteq [0, 1];$$

- if $b-l' \leq r < b$, then

$$N/b^h - q + l'/b - 1 \in \left[\frac{r+l'-b}{b}, \frac{r+l'-b+1}{b}\right] \subseteq [0, 1).$$

Using the periodicity of φ_b^π and Equation (11) we therefore obtain

$$\begin{aligned} \sum_{N=1}^{b^n} N\varphi_b^\pi\left(\frac{N}{b^h} + \frac{l'}{b}\right) &= \sum_{r=0}^{b-1} \sum_{q=0}^{b^{n-h}-1} \sum_{N=qb^{h-1}+rb^{h-1}+1}^{qb^{h-1}+(r+1)b^{h-1}} N\varphi_b^\pi\left(\frac{N}{b^h} - q + \frac{l'}{b}\right) \\ &= \sum_{r=0}^{b-l'-1} \sum_{q=0}^{b^{n-h}-1} \sum_{N=qb^{h-1}+rb^{h-1}+1}^{qb^{h-1}+(r+1)b^{h-1}} N\varphi_b^\pi\left(\frac{N}{b^h} - q + \frac{l'}{b}\right) \\ &\quad + \sum_{r=b-l'}^{b-1} \sum_{q=0}^{b^{n-h}-1} \sum_{N=qb^{h-1}+rb^{h-1}+1}^{qb^{h-1}+(r+1)b^{h-1}} N\varphi_b^\pi\left(\frac{N}{b^h} - q + \frac{l'}{b} - 1\right) \\ &= \sum_{r=0}^{b-l'-1} \sum_{q=0}^{b^{n-h}-1} \sum_{N=qb^{h-1}+rb^{h-1}+1}^{qb^{h-1}+(r+1)b^{h-1}} N \left(\frac{b(b-2\pi(r+l'))-1}{2}\right) \left(\frac{N}{b^h} - q - \frac{r}{b}\right) \end{aligned}$$

$$\begin{aligned}
& + \frac{(r+l')(b-1)}{2} - \sum_{i=0}^{r+l'-1} \pi(i) \Big) \\
& + \sum_{r=b-l'}^{b-1} \sum_{q=0}^{b^n-h-1} \sum_{N=qb^h+r b^{h-1}+1}^{qb^h+(r+1)b^{h-1}} N \left(\frac{b(b-2\pi(r+l'-b)-1)}{2} \left(\frac{N}{b^h} - q - \frac{r}{b} \right) \right. \\
& \left. + \frac{(r+l'-b)(b-1)}{2} - \sum_{i=0}^{r+l'-b-1} \pi(i) \right) \\
& = \Sigma_1 + \Sigma_2 + \Sigma_3 + \Sigma_4,
\end{aligned}$$

where:

$$\begin{aligned}
\Sigma_1 & := \sum_{r=0}^{b-l'-1} \frac{(b-2\pi(r+l')-1)b^n(b^h+b)(2b-3b^{h+1}+3b^{n+1}+b^h(4+6r))}{24b^{2+h}} \\
\Sigma_2 & := \sum_{r=0}^{b-l'-1} \left(\frac{(r+l')(b-1)}{2} - \sum_{i=0}^{r+l'-1} \pi(i) \right) \frac{b^n(b-b^{h+1}+b^{n+1}+b^h(1+2r))}{2b^2} \\
\Sigma_3 & := \sum_{r=b-l'}^{b-1} \frac{(b-2\pi(r+l'-b)-1)b^n(b^h+b)(2b-3b^{h+1}+3b^{n+1}+b^h(4+6r))}{24b^{2+h}} \\
\Sigma_4 & := \sum_{r=b-l'}^{b-1} \left(\frac{(r+l'-b)(b-1)}{2} - \sum_{i=0}^{r+l'-b-1} \pi(i) \right) \\
& \quad \times \frac{b^n(b-b^{h+1}+b^{n+1}+b^h(1+2r))}{2b^2}
\end{aligned}$$

Now tedious calculations using (12) lead to the desired result. \square

4. The proofs of the main results

First we show a discrete version of Proposition 1.

LEMMA 8. *For $\Sigma = (\sigma_0, \dots, \sigma_{n-1}) \in \{\pi_l : 0 \leq l < b\}^n$ let $\lambda_l := \#\{0 \leq i < n : \sigma_i = \pi_l\}$. Then we have*

$$\frac{1}{b^{2n}} \sum_{\lambda, N=1}^{b^n} E \left(\frac{\lambda}{b^n}, \frac{N}{b^n}; \mathcal{H}_{b,n}^\Sigma \right) = \sum_{l=0}^{b-1} \frac{\lambda_l}{b} \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{\pi^{-1}(l)}{b} \right) \right) \quad (18)$$

and

$$\begin{aligned}
 & \frac{1}{b^{2n}} \sum_{\lambda, N=1}^{b^n} \left(E \left(\frac{\lambda}{b^n}; \frac{N}{b^n}; \mathcal{H}_{b,n}^\Sigma \right) \right)^2 \\
 &= \left(\sum_{l=0}^{b-1} \frac{\lambda_l}{b} \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{\pi^{-1}(l)}{b} \right) \right) \right)^2 + n\Phi_b^{\pi, (2)} + \frac{1}{36} \left(1 - \frac{1}{b^{2n}} \right) \\
 & \quad + \sum_{l=0}^{b-1} \frac{\lambda_l}{b} \left[\varphi_b^{\pi, (2)} \left(\frac{\pi^{-1}(l)}{b} \right) - 2F_b^\pi(l) - \frac{1}{b} \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{\pi^{-1}(l)}{b} \right) \right)^2 \right].
 \end{aligned} \tag{19}$$

Proof. We just give the (much more involved) proof of (19). Equation (18) can be shown in the same way.

When $\lambda_l \neq 0$, let $h_i^{(l)}$ for $1 \leq i \leq \lambda_l$ ($1 \leq h_i^{(l)} \leq n$) be the integers such that $\sigma_{h_i^{(l)}-1} = \pi_l$, i.e. $\sigma_{h_1^{(l)}-1} = \dots = \sigma_{h_{\lambda_l}^{(l)}-1} = \pi_l$. Using Lemma 3 and Lemma 4 we have

$$\begin{aligned}
 & \frac{1}{b^{2n}} \sum_{\lambda, N=1}^{b^n} \left(E \left(\frac{\lambda}{b^n}; \frac{N}{b^n}; \mathcal{H}_{b,n}^\Sigma \right) \right)^2 = \frac{1}{b^{2n}} \sum_{\lambda, N=1}^{b^n} \sum_{i,j=1}^n \varphi_{b,\varepsilon_i}^{\sigma_{i-1}} \left(\frac{N}{b^i} \right) \varphi_{b,\varepsilon_j}^{\sigma_{j-1}} \left(\frac{N}{b^j} \right) \\
 &= \frac{1}{b^{2n}} \sum_{i=1}^n \sum_{N=1}^{b^n} \sum_{\lambda=1}^{b^n} \left(\varphi_{b,\varepsilon_i}^{\sigma_{i-1}} \left(\frac{N}{b^i} \right) \right)^2 \\
 & \quad + \frac{1}{b^{2n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{N=1}^{b^n} \sum_{\lambda=1}^{b^n} \varphi_{b,\varepsilon_i}^{\sigma_{i-1}} \left(\frac{N}{b^i} \right) \varphi_{b,\varepsilon_j}^{\sigma_{j-1}} \left(\frac{N}{b^j} \right) \\
 &= \frac{1}{b^{2n}} \sum_{i=1}^n \sum_{N=1}^{b^n} b^{n-1} \varphi_b^{\sigma_{i-1}, (2)} \left(\frac{N}{b^i} \right) \\
 & \quad + \frac{1}{b^{2n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{N=1}^{b^n} b^{n-2} \varphi_b^{\sigma_{i-1}} \left(\frac{N}{b^i} \right) \varphi_b^{\sigma_{j-1}} \left(\frac{N}{b^j} \right) \\
 &= \frac{1}{b^{2n}} \sum_{l=0}^{b-1} \sum_{i=1}^{\lambda_l} \sum_{N=1}^{b^n} b^{n-1} \varphi_b^{\pi_l, (2)} \left(\frac{N}{b^{h_i^{(l)}}} \right) \\
 & \quad + \frac{1}{b^{2n}} \sum_{\substack{l,m=0 \\ l \neq m}}^{b-1} \sum_{i=1}^{\lambda_l} \sum_{j=1}^{\lambda_m} \sum_{N=1}^{b^n} b^{n-2} \varphi_b^{\pi_l} \left(\frac{N}{b^{h_i^{(l)}}} \right) \varphi_b^{\pi_m} \left(\frac{N}{b^{h_j^{(m)}}} \right)
 \end{aligned}$$

$$+ \frac{1}{b^{2n}} \sum_{l=0}^{b-1} \sum_{\substack{i,j=1 \\ i \neq j}}^{\lambda_l} \sum_{N=1}^{b^n} b^{n-2} \varphi_b^{\pi l} \left(\frac{N}{b^{h_i^{(l)}}} \right) \varphi_b^{\pi l} \left(\frac{N}{b^{h_j^{(l)}}} \right)$$

$=: A + B + C.$

Using Lemma 5 we get (again we write $l' = \pi^{-1}(l)$ and $m' = \pi^{-1}(m)$)

$$B = \frac{1}{b^2} \sum_{\substack{l,m=0 \\ l \neq m}}^{b-1} \lambda_l \lambda_m \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{l'}{b} \right) \right) \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{m'}{b} \right) \right)$$

and

$$C = \frac{1}{b^2} \sum_{l=0}^{b-1} \lambda_l (\lambda_l - 1) \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{l'}{b} \right) \right)^2.$$

For A we use Lemma 6 and the fact that $\sum_{l=0}^{b-1} \lambda_l = n$ to obtain

$$\begin{aligned} A &= \frac{1}{b^{2n}} \sum_{l=0}^{b-1} \sum_{i=1}^{\lambda_l} \sum_{N=1}^{b^n} b^{n-1} \varphi_b^{\pi l, (2)} \left(\frac{N}{b^{h_i^{(l)}}} \right) \\ &= \frac{1}{b} \sum_{l=0}^{b-1} \sum_{i=1}^{\lambda_l} \left(b\Phi_b^{\pi, (2)} + \frac{b(b^2 - 1)}{36b^{2h_i^{(l)}}} + \varphi_b^{\pi, (2)} \left(\frac{l'}{b} \right) - 2F_b^\pi(l) \right) \\ &= n\Phi_b^{\pi, (2)} + \frac{1}{36} \left(1 - \frac{1}{b^{2n}} \right) + \sum_{l=0}^{b-1} \frac{\lambda_l}{b} \left(\varphi_b^{\pi, (2)} \left(\frac{l'}{b} \right) - 2F_b^\pi(l) \right). \end{aligned}$$

Overall we obtain

$$\begin{aligned} &\frac{1}{b^{2n}} \sum_{\lambda, N=1}^{b^n} \left(E \left(\frac{\lambda}{b^n}; \frac{N}{b^n}; \mathcal{H}_{b,n}^\Sigma \right) \right)^2 \\ &= \left(\sum_{l=0}^{b-1} \frac{\lambda_l}{b} \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{l'}{b} \right) \right) \right)^2 - \sum_{l=0}^{b-1} \frac{\lambda_l}{b^2} \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{l'}{b} \right) \right)^2 \\ &\quad + n\Phi_b^{\pi, (2)} + \frac{1}{36} \left(1 - \frac{1}{b^{2n}} \right) + \sum_{l=0}^{b-1} \frac{\lambda_l}{b} \left(\varphi_b^{\pi, (2)} \left(\frac{l'}{b} \right) - 2F_b^\pi(l) \right) \\ &= \left(\sum_{l=0}^{b-1} \frac{\lambda_l}{b} \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{l'}{b} \right) \right) \right)^2 + n\Phi_b^{\pi, (2)} + \frac{1}{36} \left(1 - \frac{1}{b^{2n}} \right) \\ &\quad + \sum_{l=0}^{b-1} \frac{\lambda_l}{b} \left[\varphi_b^{\pi, (2)} \left(\frac{l'}{b} \right) - 2F_b^\pi(l) - \frac{1}{b} \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{l'}{b} \right) \right)^2 \right]. \end{aligned}$$

□

Now we give the proof of Proposition 1.

PROOF. Again we use the abbreviation $l' = \pi^{-1}(l)$. Using (14) we obtain

$$\begin{aligned}
 (L_2(\mathcal{H}_{b,n}^\Sigma))^2 &= \int_0^1 \int_0^1 (E(x(n); y(n); \mathcal{H}_{b,n}^\Sigma) + b^n(x(n)y(n) - xy))^2 dx dy \\
 &= \frac{1}{b^{2n}} \sum_{\lambda, N=1}^{b^n} \left(E\left(\frac{\lambda}{b^n}; \frac{N}{b^n}; \mathcal{H}_{b,n}^\Sigma\right) \right)^2 \\
 &\quad + 2b^n \sum_{\lambda, N=1}^{b^n} \int_{\frac{\lambda-1}{b^n}}^{\frac{\lambda}{b^n}} \int_{\frac{N-1}{b^n}}^{\frac{N}{b^n}} E\left(\frac{\lambda}{b^n}; \frac{N}{b^n}; \mathcal{H}_{b,n}^\Sigma\right) \left(\frac{\lambda}{b^n} \frac{N}{b^n} - xy\right) dx dy \\
 &\quad + b^{2n} \sum_{\lambda, N=1}^{b^n} \int_{\frac{\lambda-1}{b^n}}^{\frac{\lambda}{b^n}} \int_{\frac{N-1}{b^n}}^{\frac{N}{b^n}} \left(\frac{\lambda}{b^n} \frac{N}{b^n} - xy\right)^2 dx dy \\
 &=: \Sigma_1 + \Sigma_2 + \Sigma_3.
 \end{aligned}$$

The term Σ_1 has been evaluated in Lemma 8 and straightforward calculus shows that $\Sigma_3 = (1 + 18b^n + 25b^{2n})/(72b^{2n})$. So it remains to deal with Σ_2 .

Evaluating the integral appearing in Σ_2 we obtain

$$\begin{aligned}
 \Sigma_2 &= \frac{1}{b^{3n}} \sum_{\lambda, N=1}^{b^n} (\lambda + N) E\left(\frac{\lambda}{b^n}; \frac{N}{b^n}; \mathcal{H}_{b,n}^\Sigma\right) - \frac{1}{2b^{3n}} \sum_{\lambda, N=1}^{b^n} E\left(\frac{\lambda}{b^n}; \frac{N}{b^n}; \mathcal{H}_{b,n}^\Sigma\right) \\
 &=: \Sigma_4 - \Sigma_5.
 \end{aligned}$$

The term Σ_5 can be obtained from Lemma 8, Equation (18). For Σ_4 we have

$$\begin{aligned}
 \Sigma_4 &= \frac{1}{b^{3n}} \sum_{\lambda, N=1}^{b^n} \lambda E\left(\frac{\lambda}{b^n}; \frac{N}{b^n}; \mathcal{H}_{b,n}^\Sigma\right) + \frac{1}{b^{3n}} \sum_{\lambda, N=1}^{b^n} N E\left(\frac{\lambda}{b^n}; \frac{N}{b^n}; \mathcal{H}_{b,n}^\Sigma\right) \\
 &=: \frac{1}{b^{3n}} (\Sigma_{4,1} + \Sigma_{4,2}).
 \end{aligned}$$

As to $\Sigma_{4,2}$, with Lemma 3, Lemma 4, Equation (13) and Lemma 7 we obtain

$$\begin{aligned}
 \Sigma_{4,2} &= b^{n-1} \sum_{l=0}^{b-1} \sum_{i=1}^{\lambda_l} \sum_{N=1}^{b^n} N \left(\varphi_b^\pi \left(\frac{N}{b^{h_i^{(l)}}} + \frac{l'}{b} \right) - \varphi_b^\pi \left(\frac{l'}{b} \right) \right) \\
 &= b^{2n-1} \sum_{l=0}^{b-1} \sum_{i=1}^{\lambda_l} \left(b^{n+1} \frac{\Phi_b^\pi}{2} + b^{h_i^{(l)}} f_b^\pi(l) + g_b^\pi(l) - \frac{b^n + 1}{2} \varphi_b^\pi \left(\frac{l'}{b} \right) \right)
 \end{aligned}$$

$$= b^{3n} \frac{\Phi_b^\pi}{2} n + b^{2n-1} \sum_{l=0}^{b-1} \sum_{i=1}^{\lambda_l} \left(b^{h_i^{(l)}} f_b^\pi(l) + g_b^\pi(l) - \frac{b^n+1}{2} \varphi_b^\pi \left(\frac{l}{b} \right) \right),$$

where again the integers $h_i^{(l)}$ satisfy $\sigma_{h_1^{(l)}-1} = \dots = \sigma_{h_{\lambda_l}^{(l)}-1} = \pi_l$ if $\lambda_l \neq 0$.

We turn to $\Sigma_{4,1}$. Let $g : [0, 1]^2 \rightarrow [0, 1]^2$ be defined by $g(x, y) = (y, x)$ and for $\Sigma = (\sigma_0, \dots, \sigma_{n-1})$ define $\Gamma = (\gamma_0, \dots, \gamma_{n-1}) := (\sigma_{n-1}^{-1}, \dots, \sigma_0^{-1})$. Then it is easy to see (for details see [9, Proof of Theorem 4]) that $\mathcal{H}_{b,n}^\Sigma = g \left(\mathcal{H}_{b,n}^\Gamma \right)$. Therefore we obtain

$$\Sigma_{4,1} = \sum_{\lambda, N=1}^{b^n} \lambda E \left(\frac{\lambda}{b^n}; \frac{N}{b^n}; \mathcal{H}_{b,n}^\Sigma \right) = \sum_{\lambda, N=1}^{b^n} \lambda E \left(\frac{N}{b^n}; \frac{\lambda}{b^n}; \mathcal{H}_{b,n}^\Gamma \right),$$

which will allow us to use the result for $\Sigma_{4,2}$. To this end, we must check the correspondences between Σ and Γ : For $\Sigma \in \{\pi_l : 0 \leq l < b\}^n$ we also have $\Gamma \in \{\pi_l : 0 \leq l < b\}^n$. If $l = 0$ we get $(\pi_0)^{-1} = (\pi^{-1})_0$ and for $0 < l < b$, we get $(\pi_l)^{-1} = (\pi^{-1})_{b-\pi^{-1}(l)}$ as remarked just before Lemma 1. Hence for $l > 0$ and $\lambda_l \neq 0$ we have $\sigma_{h_1^{(l)}-1}^{-1} = \dots = \sigma_{h_{\lambda_l}^{(l)}-1}^{-1} = (\pi_l)^{-1} = (\pi^{-1})_{b-\pi^{-1}(l)}$, so that $\gamma_{n+1-h_1^{(l)}-1} = \dots = \gamma_{n+1-h_{\lambda_l}^{(l)}-1} = (\pi^{-1})_{b-\pi^{-1}(l)}$. Further $\gamma_{u_1^{(b-r)}-1} = \dots = \gamma_{u_{\lambda_{\pi(b-r)}}-1} = (\pi^{-1})_r$, where $u_i^{(b-r)} := n - h_i^{\pi(b-r)} + 1$. Since π^{-1} is also linear we may use the formula for $\Sigma_{4,2}$ and obtain

$$\begin{aligned} \Sigma_{4,1} &= b^{3n} \frac{\Phi_b^{\pi^{-1}}}{2} n + b^{2n-1} \\ &\quad \times \sum_{r=0}^{b-1} \sum_{i=1}^{\lambda_{\pi(b-r)}} \left(b^{u_i^{(b-r)}} f_b^{\pi^{-1}}(r) + g_b^{\pi^{-1}}(r) - \frac{b^n+1}{2} \varphi_b^{\pi^{-1}} \left(\frac{\pi(r)}{b} \right) \right) \\ &= b^{3n} \frac{\Phi_b^\pi}{2} n + b^{2n-1} \\ &\quad \times \sum_{r=0}^{b-1} \sum_{i=1}^{\lambda_{\pi(b-r)}} \left(b^{n-h_i^{\pi(b-r)}+1} f_b^{\pi^{-1}}(r) + g_b^{\pi^{-1}}(r) - \frac{b^n+1}{2} \varphi_b^{\pi^{-1}} \left(\frac{\pi(r)}{b} \right) \right) \\ &= b^{3n} \frac{\Phi_b^\pi}{2} n + b^{2n-1} \\ &\quad \times \sum_{l=0}^{b-1} \sum_{i=1}^{\lambda_l} \left(b^{n-h_i^{(l)}+1} f_b^{\pi^{-1}}(b-l) + g_b^{\pi^{-1}}(b-l) - \frac{b^n+1}{2} \varphi_b^{\pi^{-1}} \left(\frac{b-l}{b} \right) \right) \end{aligned}$$

where we have used that $\Phi_b^{\pi^{-1}} = \Phi_b^\pi$ as shown in [12, Lemma 5]. Hence we have

$$\begin{aligned} \Sigma_4 &= \Phi_b^\pi n - \frac{b^n + 1}{2b^{n+1}} \sum_{l=0}^{b-1} \lambda_l \left(\varphi_b^\pi \left(\frac{l'}{b} \right) + \varphi_b^{\pi^{-1}} \left(\frac{b-l}{b} \right) \right) \\ &\quad + \frac{1}{b^{n+1}} \sum_{l=0}^{b-1} \sum_{i=1}^{\lambda_l} \left(b^{h_i^{(l)}} f_b^\pi(l) + b^{n-h_i^{(l)}+1} f_b^{\pi^{-1}}(b-l') \right) \\ &\quad + \frac{1}{b^{n+1}} \sum_{l=0}^{b-1} \lambda_l \left(g_b^\pi(l) + g_b^{\pi^{-1}}(b-l') \right). \end{aligned}$$

Now we obtain

$$\begin{aligned} (L_2(\mathcal{H}_{b,n}^\Sigma))^2 &= \left(\sum_{l=0}^{b-1} \frac{\lambda_l}{b} \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{l'}{b} \right) \right) \right)^2 + n\Phi_b^{\pi,(2)} + \frac{1}{36} \left(1 - \frac{1}{b^{2n}} \right) \\ &\quad + \sum_{l=0}^{b-1} \frac{\lambda_l}{b} \left[\varphi_b^{\pi,(2)} \left(\frac{l'}{b} \right) - 2F_b^\pi(l) - \frac{1}{b} \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{l'}{b} \right) \right)^2 \right] \\ &\quad + \Phi_b^\pi n - \frac{b^n + 1}{2b^{n+1}} \sum_{l=0}^{b-1} \lambda_l \left(\varphi_b^\pi \left(\frac{l'}{b} \right) + \varphi_b^{\pi^{-1}} \left(\frac{b-l}{b} \right) \right) \\ &\quad + \frac{1}{b^{n+1}} \sum_{l=0}^{b-1} \sum_{i=1}^{\lambda_l} \left(b^{h_i^{(l)}} f_b^\pi(l) + b^{n-h_i^{(l)}+1} f_b^{\pi^{-1}}(b-l') \right) \\ &\quad + \frac{1}{b^{n+1}} \sum_{l=0}^{b-1} \lambda_l \left(g_b^\pi(l) + g_b^{\pi^{-1}}(b-l') \right) \\ &\quad - \frac{1}{2b^n} \sum_{l=0}^{b-1} \frac{\lambda_l}{b} \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{l'}{b} \right) \right) + \frac{1 + 18b^n + 25b^{2n}}{72b^{2n}} \\ &= \left(\sum_{l=0}^{b-1} \frac{\lambda_l}{b} \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{l'}{b} \right) \right) \right)^2 + n\Phi_b^{\pi,(2)} + n\Phi_b^\pi \\ &\quad + \sum_{l=0}^{b-1} \frac{\lambda_l}{b} \left[\varphi_b^{\pi,(2)} \left(\frac{l'}{b} \right) - 2F_b^\pi(l) - \frac{1}{b} \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{l'}{b} \right) \right)^2 \right] \\ &\quad - \frac{1}{2b} \sum_{l=0}^{b-1} \lambda_l \left(\varphi_b^\pi \left(\frac{l'}{b} \right) + \varphi_b^{\pi^{-1}} \left(\frac{b-l}{b} \right) \right) + O(1). \quad (20) \end{aligned}$$

From Lemma 2 it follows that

$$\Phi_b^\pi - \frac{1}{b}\varphi_b^\pi(l'/b) = \Phi_b^{\pi_l} \quad (21)$$

and from [12, Lemma 5] we know that $\Phi_b^\pi = \Phi_b^{\pi^{-1}}$ and hence

$$\begin{aligned} n\Phi_b^\pi - \frac{1}{2b} \sum_{l=0}^{b-1} \lambda_l \left(\varphi_b^\pi \left(\frac{l'}{b} \right) + \varphi_b^{\pi^{-1}} \left(\frac{b-l}{b} \right) \right) \\ &= \sum_{l=0}^{b-1} \lambda_l \left(\Phi_b^\pi - \frac{1}{2b}\varphi_b^\pi \left(\frac{l'}{b} \right) - \frac{1}{2b}\varphi_b^{\pi^{-1}} \left(\frac{b-l}{b} \right) \right) \\ &= \frac{1}{2} \sum_{l=0}^{b-1} \lambda_l \Phi_b^{\pi_l} + \frac{1}{2} \sum_{l=0}^{b-1} \lambda_l \left(\Phi_b^{\pi^{-1}} - \frac{1}{b}\varphi_b^{\pi^{-1}} \left(\frac{b-l}{b} \right) \right) \\ &= \frac{1}{2} \sum_{l=0}^{b-1} \lambda_l \Phi_b^{\pi_l} + \frac{1}{2} \sum_{l=0}^{b-1} \lambda_l \Phi_b^{(\pi^{-1})_{\pi^{-1}(b-l \pmod{b})}} \\ &= \frac{1}{2} \sum_{l=0}^{b-1} \lambda_l \Phi_b^{\pi_l} + \frac{1}{2} \sum_{l=0}^{b-1} \lambda_l \Phi_b^{(\pi_l)^{-1}} = \sum_{l=0}^{b-1} \lambda_l \Phi_b^{\pi_l}. \end{aligned}$$

The desired result follows from inserting this and (21) into (20). \square

We give the proof of Theorem 1.

Proof. If $\lambda_l = \lfloor n/b \rfloor + \theta_l$ with $\theta_l \in \{0, 1\}$, then from (20) we obtain

$$\begin{aligned} &(L_2(\mathcal{H}_{b,n}^\Sigma))^2 \\ &= n\Phi_b^{\pi, (2)} + \frac{n}{b^2} \sum_{l=0}^{b-1} \left[\varphi_b^{\pi, (2)} \left(\frac{l'}{b} \right) - 2F_b^\pi(l) - \frac{1}{b} \left(b\Phi_b^\pi - \varphi_b^\pi \left(\frac{l'}{b} \right) \right)^2 \right] + O(1) \\ &= n(\Phi_b^{\pi, (2)} - (\Phi_b^\pi)^2) + \frac{n}{b^2} A + O(1), \text{ where} \end{aligned}$$

$$A := \sum_{l=0}^{b-1} \left[\varphi_b^{\pi, (2)} \left(\frac{l'}{b} \right) - 2F_b^\pi(l) + 2\Phi_b^\pi \varphi_b^\pi \left(\frac{l'}{b} \right) - \frac{1}{b} \left(\varphi_b^\pi \left(\frac{l'}{b} \right) \right)^2 \right].$$

It remains to prove that $A = 0$. Writing everything with $\varphi_{b,h}^\pi(\frac{l}{b})$, we get

$$A = \sum_{l,h=0}^{b-1} \left(\varphi_{b,h}^\pi \left(\frac{l}{b} \right) \right)^2 - \frac{2}{b} \sum_{h,j,l=0}^{b-1} \varphi_{b,h}^\pi \left(\frac{l}{b} \right) \varphi_{b,h}^\pi \left(\frac{j}{b} \right)$$

$$+ \frac{2}{b^2} \left(\sum_{l,h=0}^{b-1} \varphi_{b,h}^\pi \left(\frac{l}{b} \right) \right)^2 - \frac{1}{b} \sum_{l=0}^{b-1} \left(\sum_{h=0}^{b-1} \varphi_{b,h}^\pi \left(\frac{l}{b} \right) \right)^2.$$

Using (9) the above expression can be written as

$$\begin{aligned} A &= \frac{1}{b^2} \sum_{l,h=0}^{b-1} \sum_{k_1=0}^{l-1} (\varphi_{b,h}^\pi)' \left(\frac{k_1}{b} + 0 \right) \sum_{k_2=0}^{l-1} (\varphi_{b,h}^\pi)' \left(\frac{k_2}{b} + 0 \right) \\ &\quad - \frac{2}{b^3} \sum_{h,j,l=0}^{b-1} \sum_{k_1=0}^{l-1} (\varphi_{b,h}^\pi)' \left(\frac{k_1}{b} + 0 \right) \sum_{k_2=0}^{j-1} (\varphi_{b,h}^\pi)' \left(\frac{k_2}{b} + 0 \right) \\ &\quad + \frac{2}{b^4} \sum_{h_1,l,h_2,j=0}^{b-1} \sum_{k_1=0}^{l-1} (\varphi_{b,h_1}^\pi)' \left(\frac{k_1}{b} + 0 \right) \sum_{k_2=0}^{j-1} (\varphi_{b,h_2}^\pi)' \left(\frac{k_2}{b} + 0 \right) \\ &\quad - \frac{1}{b^3} \sum_{l=0}^{b-1} \left(\sum_{h=0}^{b-1} \sum_{k=0}^{l-1} (\varphi_{b,h}^\pi)' \left(\frac{k}{b} + 0 \right) \right)^2. \end{aligned}$$

In [12, Section 5] it is shown that

$$\begin{aligned} \sum_{h=0}^{b-1} (\varphi_{b,h}^\pi)' \left(\frac{k_1}{b} + 0 \right) (\varphi_{b,h}^\pi)' \left(\frac{k_2}{b} + 0 \right) &= \frac{b(b-1)(2b-1)}{6} \\ &\quad + \frac{b}{2} (\pi(k_1)(\pi(k_1)+1) + \pi(k_2)(\pi(k_2)+1)) - b^2 \max(\pi(k_1), \pi(k_2)) \end{aligned}$$

and $\sum_{h=0}^{b-1} (\varphi_{b,h}^\pi)' \left(\frac{k}{b} + 0 \right) = b(b-1)/2 - b\pi(k)$.

With these formulas and using the fact that $\pi(b-k) = b-\pi(k)$, $k \neq 0$, one gets after tedious computations that $A_0 = b^4 A$ consists of sums S_i whose summands are all polynomial in $b, k, l, \pi(k), \pi(l)$ and $\max(k, l), \max(\pi(k), \pi(l))$. These polynomials will be denoted by p_i . Furthermore, we can single out polynomials in b from the sums. The sums are then dependent on π while the polynomials depend only on b but not on π . I.e., we can make an ansatz for A_0 , for which we then subsequently want to determine the p_i :

$$A_0 = A_0(\pi) = \sum_{i=0}^5 p_i(b) S_i(\pi),$$

where the p_i are polynomials in b and

$$S_0 \equiv 1, \quad S_i(\pi) := \sum_{k=0}^{b-1} k^i \pi(k)^i, \quad \text{for } i = 1, 2,$$

$$S_3(\pi) := \sum_{k,l=0}^{b-1} \max(k,l) \max(\pi(k), \pi(l)),$$

$$S_4(\pi) := \sum_{k,l=0}^{b-1} kl \max(\pi(k), \pi(l)),$$

$$S_5(\pi) := S_1(\pi)^2.$$

Any linear dependencies between the S_i (which we were not able to show directly) will appear as additional dimensions to the solution space of the ansatz.

Now, to determine the p_i , we first solve the following linear equation system for fixed b , where b is large enough such that there exist at least as many π as S_i :

$$(S_i(\pi))_{\pi,i}(x_i)_i = (A_0(\pi))_{\pi}.$$

The left hand side matrix is in $\mathbb{Z}^{\varphi(b) \times b}$, where $\varphi(b)$ is the Euler phi function.

Since by the ansatz the system is solvable, we obtain as particular solution the values of p_i at the fixed b , i.e., the vector $(p_i(b))_i$. If the left hand side matrix is of full rank we have a unique solution, otherwise we additionally have a nullspace N_b (which we consider in a canonical form).

Repeating this for sufficiently many b we can derive the p_i by interpolation. Note that the degrees of the p_i are bounded by the maximum degree of terms inside the sums plus the depth with which they are nested. (In this application the bound is 8.)

Similarly we also derive a basis of the nullspace and get

$$A_0(\pi) = \sum_i \left(p_i(b) + \sum_j \lambda_j q_{i,j}(b) \right) S_i(\pi),$$

for some polynomials $q_{i,j}$ and arbitrarily chosen $\lambda_j \in \mathbb{Q}$, where

$$0 = \sum_i q_{i,j}(b) S_i(\pi),$$

and with j ranging over the maximum of dimensions of the N_b . So at least one S_i is linearly dependent. We repeat the process with this S_i excluded and A_0 multiplied with $\text{lcm}_j(q_{i,j})$. In the iteration the dimension of the nullspaces is now reduced by one, so the algorithm terminates after finitely many steps.

By an implementation in MATHEMATICA we observe that, in fact, in this application

$$S_4 = -bS_3 + S_2 + bS_1 + \frac{b^3(b-1)^2}{2}$$

$$4S_5 = 4b^2S_3 - 5bS_2 + b^2(5b - 8)S_1 - \frac{4b^2(40b^4 - 72b^3 + 31b^2 + 1)}{72}.$$

The matrices $(S_i(\pi))_{\pi; i=0,1,2,3}$ are of full rank for $b = 20, \dots, 35$ and any choices of π . Furthermore all right hand side vectors $A_0(\pi)$ evaluate to zero. So finally we obtain that all p_i are identically zero, and thus $A_0 = 0$ and the same holds for A . \square

Acknowledgment: The comments and suggestions of an anonymous referee concerning the presentation of the results are greatly appreciated.

REFERENCES

- [1] H. Chaix and H. Faure: Discr ance et diaphonie en dimension un. *Acta Arith.* 63: 103–141, 1993.
- [2] W. W. L. Chen and M. M. Skriganov: Explicit constructions in the classical mean squares problem in irregularities of point distribution. *J. Reine Angew. Math.* 545: 67–95, 2002.
- [3] W. W. L. Chen and M. M. Skriganov: Orthogonality and digit shifts in the classical mean squares problem in irregularities of point distribution. In: *Diophantine approximation*, 141–159, *Dev. Math.*, 16, Springer Wien New York, Vienna, 2008.
- [4] H. Davenport: Note on irregularities of distribution. *Mathematika* 3: 131–135, 1956.
- [5] J. Dick and F. Pillichshammer: *Digital Nets and Sequences. Discrepancy Theory and Quasi-Monte Carlo Integration*. Cambridge University Press, 2010.
- [6] M. Drmota and R. F. Tichy: *Sequences, Discrepancies and Applications*. Lecture Notes in Mathematics 1651, Springer, Berlin, 1997.
- [7] H. Faure: Discr ance de suites associ es   un syst me de num ration (en dimension un). *Bull. Soc. Math. France* 109: 143–182, 1981.
- [8] H. Faure and C. Lemieux: Generalized Halton sequences in 2008: A comparative study, *ACM Trans. Model. Comp. Sim.* 19: Article 15, 2009.
- [9] H. Faure and F. Pillichshammer: L_p discrepancy of generalized two-dimensional Hammersley point sets. *Monatsh. Math.* 158: 31–61, 2009.
- [10] H. Faure and F. Pillichshammer: L_2 discrepancy of two-dimensional digitally shifted Hammersley point sets in base b . In: *Monte Carlo and Quasi-Monte Carlo Methods 2008* (P. L’Ecuyer and A. Owen, eds.), Springer, Wien Heidelberg New York, 2009.
- [11] H. Faure and F. Pillichshammer: L_2 discrepancy of generalized Zaremba point sets. *J. Th. Nombres de Bordeaux* 23: 121–136, 2011.
- [12] H. Faure, F. Pillichshammer, G. Pirsic and W. Ch. Schmid: L_2 discrepancy of generalized two-dimensional Hammersley point sets scrambled with arbitrary permutations. *Acta Arith.* 141: 395–418, 2010.
- [13] J. H. Halton and S. K. Zaremba: The extreme and the L^2 discrepancies of some plane sets. *Monatsh. Math.* 73: 316–328, 1969.
- [14] A. Hinrichs and L. Markhasin: On lower bounds for the L_2 -discrepancy. *J. Complexity* 27: 127–132, 2011.
- [15] P. Kritzer and F. Pillichshammer: An exact formula for the L_2 discrepancy of the shifted Hammersley point set. *Uniform Distribution Theory* 1: 1–13, 2006.

L_2 DISCREPANCY OF LINEARLY DIGIT SCRAMBLED ZAREMBA POINT SETS

- [16] L. Kuipers and H. Niederreiter: *Uniform Distribution of Sequences*. John Wiley, New York, 1974; reprint, Dover Publications, Mineola, NY, 2006.
- [17] G. Larcher and F. Pillichshammer: Walsh series analysis of the L_2 -discrepancy of symmetrized point sets. *Monatsh. Math.* 132: 1–18, 2001.
- [18] J. Matoušek: *Geometric Discrepancy: An Illustrated Guide*. Springer, Berlin Heidelberg New York, 1999.
- [19] E. Novak and H. Woźniakowski: *Tractability of Multivariate Problems. Volume I: Linear Information*. EMS Tracts in Mathematics, 6. European Mathematical Society (EMS), Zürich, 2008.
- [20] E. Novak and H. Woźniakowski: *Tractability of Multivariate Problems. Volume II: Standard Information for Functionals*. EMS Tracts in Mathematics, 12. European Mathematical Society (EMS), Zürich, 2010.
- [21] F. Pillichshammer: On the L_p -discrepancy of the Hammersley point set. *Monatsh. Math.* 136: 67–79, 2002.
- [22] P. D. Proinov: Symmetrization of the van der Corput generalized sequences. *Proc. Japan Acad.* 64, Ser. A: 159–162, 1988.
- [23] K. F. Roth: On irregularities of distribution. *Mathematika* 1: 73–79, 1954.
- [24] I. V. Vilenkin: Plane nets of Integration. *Ž. Vyčisl. Mat. i Mat. Fiz.* 7: 189–196, 1967. (English translation in: U.S.S.R. Computational Math. and Math. Phys. 7: 258–267, 1967.)
- [25] B. E. White: Mean-square discrepancies of the Hammersley and Zaremba sequences for arbitrary radix. *Monatsh. Math.* 80: 219–229, 1975.

Received December 20, 2010

Accepted May, 23 2011

Henri Faure

Institut de Mathématiques de Luminy

U.M.R. 6206 CNRS

163 avenue de Luminy, case 907

13288 Marseille Cedex 09

France

E-mail: faure@iml.univ-mrs.fr

Friedrich Pillichshammer

Gottlieb Pirsic

Institut für Finanzmathematik

Universität Linz

Altenbergerstraße 69

A-4040 Linz

Austria

E-mail: friedrich.pillichshammer@jku.at

gpirsic@gmail.com