

LOG-LIKE FUNCTIONS AND UNIFORM DISTRIBUTION MODULO ONE

MARTIN REHBERG

Technische Hochschule Mittelhessen, Friedberg, GERMANY

ABSTRACT. For a function f satisfying $f(x) = o((\log x)^K)$, $K > 0$, and a sequence of numbers $(q_n)_n$, we prove by assuming several conditions on f that the sequence $(\alpha f(q_n))_{n \geq n_0}$ is uniformly distributed modulo one for any nonzero real number α . This generalises some former results due to Too, Goto and Kano where instead of $(q_n)_n$ the sequence of primes was considered.

Communicated by Werner Georg Nowak

1. Introduction and results

Let p_n be the n th prime number in ascending order and α a nonzero real number. Then Goto and Kano [1], [2], as well as Too [7], proved by assuming several conditions on the function f that the sequence $(\alpha f(p_n))_n$ is uniformly distributed modulo one. Recall that a sequence $(x_n)_{n \in \mathbb{N}}$ of real numbers is said to be *uniformly distributed modulo one* if for every pair α, β of real numbers with $0 \leq \alpha < \beta \leq 1$ the proportion of the fractional parts of the x_n in the interval $[\alpha, \beta)$ tends to its length in the following sense:

$$\lim_{N \rightarrow \infty} \frac{\#\{1 \leq n \leq N : \{x_n\} \in [\alpha, \beta)\}}{N} = \beta - \alpha.$$

In fact Goto, Kano and Too proved their results by determining a bound for the discrepancy of the considered sequence.

DEFINITION 1. Let x_1, \dots, x_N be a finite sequence of real numbers. The number

$$D_N := D_N(x_1, \dots, x_N) := \sup_{0 \leq \alpha < \beta \leq 1} \left| \frac{\#\{1 \leq n \leq N : \{x_n\} \in [\alpha, \beta)\}}{N} - (\beta - \alpha) \right|$$

2010 Mathematics Subject Classification: 11J71, 11K38.

Keywords: Uniform Distribution, Discrepancy.

is called the **discrepancy** of the given sequence. If $\omega = (x_n)$ is an infinite sequence (or a finite sequence containing at least N terms), $D_N(\omega)$ is meant to be the discrepancy of the first N terms of ω .

It is well known that a sequence $\omega = (x_n)_{n \in \mathbb{N}}$ is uniformly distributed modulo one if and only if $\lim_{N \rightarrow \infty} D_N(\omega) = 0$. Instead of the sequence of primes, which was investigated in [1], [2] and [7], we consider a sequence of real numbers $(q_n)_{n \geq 1}$ satisfying $1 < q_1 < q_2 < \dots$ with $q_n \rightarrow \infty$ as $n \rightarrow \infty$. Further, we assume that the sequence $(q_n)_{n \geq 1}$ satisfies

$$Q(x) - c \int_2^x \frac{dt}{\log t} \ll \frac{x}{(\log x)^k} \quad (1)$$

for every positive $k > 1$, where $Q(x) := \sum_{q_n \leq x} 1$ and $c > 0$ is some constant. Note that condition (1) holds for the sequence of primes (with $c = 1$), as well as for primes in arithmetic progressions (with $c = \varphi(q)^{-1}$, where φ is Euler's function and q is the modulus).

However, the sequence $(q_n)_{n \geq 1}$ we consider satisfies (1) and for such a sequence we prove the following theorems:

THEOREM 2. *Let $a > 0$, $n_0 := \min\{n \in \mathbb{N} : q_n > a\}$ and let the function $f : [a, \infty) \rightarrow (0, \infty)$ satisfy the conditions*

- (a.) f is twice differentiable with $f' > 0$,
- (b.) $x^2 f''(x) \rightarrow -\infty$ as $x \rightarrow \infty$,
- (c.) $(\log x)^2 f''(x)$ and $x (\log x)^2 f''(x)$ are nonincreasing for sufficiently large x ,
- (d.) $f(x) = o((\log x)^K)$ for some $K > 0$ as $x \rightarrow \infty$.

Then, for any nonzero real constant α , the sequence $(\alpha f(q_n))_{n \geq n_0}$ is uniformly distributed modulo one and

$$D_N \ll \sqrt{\frac{f(q_N)}{(\log q_N)^K}} + \sqrt{\frac{1}{-q_N^2 f''(q_N)}} + \frac{1}{(\log q_N)(-q_N^2 f''(q_N))} \quad (2)$$

as $N \rightarrow \infty$.

THEOREM 3. *Let $a > 0$, $n_0 := \min\{n \in \mathbb{N} : q_n > a\}$ and let the function $f : [a, \infty) \rightarrow (0, \infty)$ satisfy the conditions:*

- (a.) f is twice differentiable with $f' > 0$,
- (b.) $x^2 f''(x) \rightarrow \infty$ as $x \rightarrow \infty$,
- (c.) $(\log x)^2 f''(x)$ is nonincreasing for sufficiently large x ,
- (d.) $f(x) = o((\log x)^K)$ for some $K > 0$ as $x \rightarrow \infty$.

LOG-LIKE FUNCTIONS

Then, for any nonzero real constant α , the sequence $(\alpha f(q_n))_{n \geq n_0}$ is uniformly distributed modulo one and

$$D_N \ll \sqrt{\frac{f(q_N)}{(\log q_N)^K}} + \sqrt{\frac{1}{q_N^2 f''(q_N)}} \quad (3)$$

as $N \rightarrow \infty$.

THEOREM 4. Let $a > 0$, $n_0 := \min\{n \in \mathbb{N} : q_n > a\}$ and let the function $f : [a, \infty) \rightarrow (0, \infty)$ satisfy the conditions:

- (a.) f is continuously differentiable,
- (b.) $x f'(x) \rightarrow \infty$ as $x \rightarrow \infty$,
- (c.) $(\log x) f'(x)$ is monotone for sufficiently large x ,
- (d.) $f(x) = o((\log x)^K)$ for some $K > 0$ as $x \rightarrow \infty$.

Then, for any nonzero real constant α , the sequence $(\alpha f(q_n))_{n \geq n_0}$ is uniformly distributed modulo one and

$$D_N \ll \sqrt{\frac{f(q_N)}{(\log q_N)^K}} + \max \left\{ \frac{1}{N}, \frac{1}{q_N f'(q_N)} \right\} \quad (4)$$

as $N \rightarrow \infty$.

In view of [1, Theorem 1] it should be remarked that in Theorem 4 a replacement of conditions (a.) and (b.) by:

- (a'.) f is continuously differentiable and $f(x) \rightarrow \infty$ as $x \rightarrow \infty$,
- (b'.) $x |f'(x)| \rightarrow \infty$ as $x \rightarrow \infty$,

would lead to the same discrepancy estimate, where only $f'(q_N)$ has to be replaced by $|f'(q_N)|$. If we compare Theorem 2 with [7, Theorem 3], one notices that the “nondecreasing” condition is replaced by “nonincreasing”. It was already remarked in [6] that this replacement is necessary.

Applying Theorem 2 to the function $f(x) = (\log x)^K$ with an arbitrary $K > 1$, we obtain that the sequence $((\log q_n)^K)_{n \geq 1}$ is uniformly distributed modulo one. This generalises the example in [7] on the uniform distribution modulo one of the sequence $((\log p_n)^K)_{n \geq 1}$.

Note that it was proved by W i n t n e r [10] that the sequence $(\log p_n)_{n \geq 1}$ is not uniformly distributed modulo one. A shorter proof can also be found in [8, Exercise 5.19]. Similarly, the sequence $(\log q_n)_{n \geq 1}$ is not uniformly distributed modulo one if (1) is replaced by $Q(x) \sim x(\log x)^{-1}$ for $x \rightarrow \infty$. The proof is analogue to the one in [8]. An example of a sequence satisfying $Q(x) \sim x(\log x)^{-1}$ for $x \rightarrow \infty$ is a sequence where each q_n fulfills $p_n \leq q_n \leq p_{n+1}$ (see [3], [5]).

2. Proofs

Except for the already pointed out change of the conditions “nondecreasing” and “nonincreasing” in Theorem 2, the theorems of this paper are generalisations of the theorems in [7]. Therefore it might not be surprising that the proofs are similar. However, for the sake of completeness we state the whole proofs and do not only point out changes in the reasoning.

In the proofs we will make use of a theorem due to Erdős and Turán to estimate the discrepancy, which was proved by them in 1940:

THEOREM 5. *For any finite sequence x_1, \dots, x_N of real numbers and any positive integer m , we have*

$$D_N \leq C \cdot \left(\frac{1}{m} + \sum_{h=1}^m \frac{1}{h} \left| \frac{1}{N} \sum_{n=1}^N e^{2\pi i h x_n} \right| \right),$$

where C is an absolute constant.

Proof of Theorem 5. can be found in [4]. In addition, we need the following estimates

LEMMA 6. *Let $F(x)$ and $G(x)$ be real functions, $\frac{G(x)}{F(x)}$ monotone and $\frac{F'(x)}{G(x)} \geq m > 0$, or $\frac{F'(x)}{G(x)} \leq -m < 0$. Then*

$$\left| \int_a^b G(x) e^{iF(x)} dx \right| \leq \frac{4}{m}.$$

LEMMA 7. *Let $G(x)$ be a positive decreasing function.*

(a.) *If $F''(x) < 0$, $F'(x) \geq 0$ and $\frac{G'(x)}{F''(x)}$ is monotone, then*

$$\left| \int_a^b G(x) e^{2\pi i F(x)} dx \right| \leq 4 \max_{a \leq x \leq b} \left\{ \frac{G(x)}{|F''(x)|^{\frac{1}{2}}} \right\} + \max_{a \leq x \leq b} \left\{ \left| \frac{G'(x)}{F''(x)} \right| \right\}.$$

(b.) *If $F''(x) > 0$ and $F'(x) \geq 0$, then*

$$\left| \int_a^b G(x) e^{2\pi i F(x)} dx \right| \leq 4 \max_{a \leq x \leq b} \left\{ \frac{G(x)}{F''(x)^{\frac{1}{2}}} \right\}.$$

The first Lemma can be found in [9, Lemma 4.3] and the second one in [11, Lemma 10.2, Lemma 10.3, p.225]. \square

LOG-LIKE FUNCTIONS

PROOF OF THEOREM 2. Note first that it is enough to prove (2), (3) and (4) for $\alpha > 0$. If we replace f by $\frac{1}{\alpha}f$ we see that it is enough to prove these statements for $\alpha = 1$. In view of condition (b.) we can assume that $f''(x) < 0$ for sufficiently large x . Further, we may assume that for $x \geq a$ we have $f''(x) < 0$ and that both $(\log x)^2 f''(x)$ and $x(\log x)^2 f''(x)$ are nonincreasing for $x \geq a$. To prove that the sequence $(f(q_n))_{n \geq n_0}$ is uniformly distributed modulo one, it is enough to prove estimation (2). In view of conditions (b.) and (d.), the term on the right side in (2) surely tends to zero as N tends to infinity. To obtain (2), our aim is to apply Theorem 5 in the form

$$D_N \ll \frac{1}{m} + \sum_{h=n_0}^m \frac{1}{h} \left| \frac{1}{N} \sum_{n=n_0}^N e^{2\pi i h f(q_n)} \right|, \quad (5)$$

where m is an arbitrary positive integer to be specified later. The essential point is to estimate the exponential sum in (5). Therefore let $q_0 := \frac{q_{n_0} + a}{2}$ and $\chi(n)$ be the characteristic function of the sequence $(q_n)_{n \geq 1}$, i.e., $\chi(n) = 1$ if n is a member of the sequence $(q_n)_{n \geq 1}$ and zero otherwise. Then $\sum_{m \leq x} \chi(n) = Q(x)$ and integration by parts yields

$$\begin{aligned} E(n_0, N; q_n) &:= \sum_{n=n_0}^N e^{2\pi i h f(q_n)} = \sum_{q_0 < m \leq q_N} e^{2\pi i h f(m)} \chi(m) \\ &= \int_{q_0}^{q_N} e^{2\pi i h f(x)} dQ(x) \\ &= Q(q_N) e^{2\pi i h f(q_N)} - Q(q_0) e^{2\pi i h f(q_0)} \\ &\quad - \int_{q_0}^{q_N} (L^*(x) + R^*(x)) d e^{2\pi i h f(x)}, \end{aligned}$$

where $R^*(x) := Q(x) - L^*(x)$, $L^*(x) := c \int_{q_0}^x \frac{dt}{\log t}$ for $c > 0$ constant, $x \geq q_0$ and $\int_a^b := \int_{(a,b]}$. Again by integration by parts,

$$\begin{aligned} \int_{q_0}^{q_N} L^*(x) d e^{2\pi i h f(x)} &= L^*(q_N) e^{2\pi i h f(q_N)} - L^*(q_0) e^{2\pi i h f(q_0)} \\ &\quad - c \int_{q_0}^{q_N} (\log x)^{-1} e^{2\pi i h f(x)} dx \end{aligned}$$

and

$$\int_{q_0}^{q_N} R^*(x) d e^{2\pi i h f(x)} = 2\pi i h \int_{q_0}^{q_N} R^*(x) f'(x) e^{2\pi i h f(x)} dx.$$

Therefore,

$$\begin{aligned} E(n_0, N; q_n) &= \left(R^*(q_N) e^{2\pi i h f(q_N)} - R^*(q_0) e^{2\pi i h f(q_0)} \right) \\ &\quad + c \int_{q_0}^{q_N} (\log x)^{-1} e^{2\pi i h f(x)} dx \\ &\quad - 2\pi i h \int_{q_0}^{q_N} R^*(x) f'(x) e^{2\pi i h f(x)} dx \\ &=: I_1 + I_2 + I_3, \end{aligned}$$

where we used $R^*(x) = Q(x) - L^*(x)$. Now we estimate each I_i ($i = 1, 2, 3$) individually: In view of our assumption (1), we get

$$R^*(x) = Q(x) - c \int_{q_0}^x \frac{dt}{\log t} \ll \frac{x}{(\log x)^k} \quad (6)$$

for every $k > 1$, implying

$$I_1 \ll \frac{q_N}{(\log q_N)^{1+K}} \quad (7)$$

for $K > 0$ as $N \rightarrow \infty$. By estimation (6) and condition (a),

$$I_3 \ll h \frac{q_N f(q_N)}{(\log q_N)^{1+K}}, \quad (8)$$

so it remains to estimate I_2 . First, let us remark that (8) for $N \rightarrow \infty$ is also an estimation for (7), since $f > 0$ and h is a positive integer. To estimate I_2 we apply Lemma 7 (a.) to get

$$\begin{aligned} I_2 &\ll \max_{q_0 \leq x \leq q_n} \left\{ \frac{4}{|h(\log x)^2 f''(x)|^{\frac{1}{2}}} + \left| \frac{1}{hx(\log x)^2 f''(x)} \right| \right\} \\ &\ll \frac{1}{(\log q_N) (-h f''(q_N))^{\frac{1}{2}}} + \frac{1}{q_N (\log q_N)^2 (-h f''(q_N))} \end{aligned}$$

as $N \rightarrow \infty$. Note that for the last estimation condition (c.) is needed in terms of “nonincreasing”. Putting our estimation for I_2 and (8) in (5) yields

$$D_N \ll \frac{1}{m} + \frac{1}{N(\log q_N) (-f''(q_N))^{\frac{1}{2}}} + \frac{1}{N q_N (\log q_N)^2 (-f''(q_N))} + \frac{m q_N f(q_N)}{N(\log q_N)^{1+K}} \quad (9)$$

for $N \rightarrow \infty$. After comparing the first and the last term in (9), we choose

$$m = \left[\left(N \cdot \frac{(\log q_N)^{1+K}}{q_N f(q_N)} \right)^{\frac{1}{2}} \right].$$

LOG-LIKE FUNCTIONS

Using this together with $Q(q_N) = N$ and $Q(x) = \frac{cx}{\log x} + O\left(\frac{x}{(\log x)^2}\right)$, we end up with our desired estimation (2). \square

Proof of Theorem 3. In respect of condition (b.) we can assume that $f''(x) > 0$ for sufficiently large x . Further, we may assume that for $x \geq a$ we have $f''(x) > 0$ and that $(\log x)^2 f''(x)$ is nonincreasing for $x \geq a$. Like in the proof of the previous theorem, we will show that the discrepancy D_N of the sequence $(f(q_n))_{n=n_0}^N$ tends to zero as $N \rightarrow \infty$. Since the estimations (7) and (8) for I_1 and I_3 still hold in the considered setting, it remains to estimate I_2 . By applying Lemma 7 (b.) we get

$$I_2 \ll \max_{q_0 \leq x \leq q_N} \left\{ \frac{1}{(h(\log x)^2 f''(x))^{\frac{1}{2}}} \right\} = \frac{1}{(h(\log q_N)^2 f''(q_N))^{\frac{1}{2}}}.$$

Combining this with (7) and (8) we obtain in (5)

$$D_N \ll \frac{1}{m} + \frac{mq_N f(q_N)}{N(\log q_N)^{1+K}} + \frac{1}{(q_N^2 f''(q_N))^{\frac{1}{2}}},$$

where we used $Q(q_N) = N$ and $Q(x) = \frac{cx}{\log x} + O\left(\frac{x}{(\log x)^2}\right)$. Choosing

$$m = \left[\left(N \cdot \frac{(\log q_N)^{1+K}}{q_N f(q_N)} \right)^{\frac{1}{2}} \right],$$

we get

$$D_N \ll \left(\frac{f(q_N)}{(\log q_N)^K} \right)^{\frac{1}{2}} + \frac{1}{(q_N^2 f''(q_N))^{\frac{1}{2}}},$$

which tends to zero as $N \rightarrow \infty$ in view of (b.) and (d.). \square

Proof of Theorem 4. With respect to condition (b.) we can assume that $f'(x) > 0$ for sufficiently large x . Further, we may assume that for $x \geq a$ we have $f'(x) > 0$ and that $(\log x)f'(x)$ is monotone for $x \geq a$. As in the previous proofs, we will show that the discrepancy D_N of the sequence $(f(q_n))_{n=n_0}^N$ tends to zero as $N \rightarrow \infty$. Since the estimations (7) and (8) for I_1 and I_3 still hold in the considered setting, it remains to estimate I_2 . Applying Lemma 6 yields

$$I_2 \ll \frac{1}{h} \cdot \max \left\{ 1, \frac{1}{\lceil \log q_N f'(q_N) \rceil} \right\}.$$

Thus, together with (7) and (8) in (5), we get

$$\begin{aligned} D_N &\ll \frac{1}{m} + \max \left\{ \frac{1}{N}, \frac{1}{N [\log q_N f'(q_N)]} \right\} + \frac{mq_N f(q_N)}{N (\log q_N)^{1+K}} \\ &\ll \frac{1}{m} + \max \left\{ \frac{1}{N}, \frac{1}{[q_N f'(q_N)]} \right\} + \frac{mq_N f(q_N)}{N (\log q_N)^{1+K}}, \end{aligned}$$

where we used in the second argument in the maximum that $Q(q_N) = N$ and $Q(x) = \frac{cx}{\log x} + O\left(\frac{x}{(\log x)^2}\right)$. After comparing the first and the last term in the discrepancy estimate, we choose

$$m = \left[\left(N \cdot \frac{(\log q_N)^{1+K}}{q_N f(q_N)} \right)^{\frac{1}{2}} \right],$$

and get

$$D_N \ll \left(\frac{f(q_N)}{(\log q_N)^K} \right)^{\frac{1}{2}} + \max \left\{ \frac{1}{N}, \frac{1}{[q_N f'(q_N)]} \right\}.$$

By condition (b.) and (d.) this tends to zero as N tends to infinity. □

ACKNOWLEDGEMENTS 1. I would like to thank Prof. J. Schoißengeier for his valuable corrections and comments.

REFERENCES

- [1] GOTO, K.—KANO, T.: *Uniform Distribution of Some Special Sequences*, Proc. Japan Acad. Ser. A **61** (1985), no. 3, 83–86.
- [2] ——— *Remarks to our Former Paper Uniform Distribution of Some Special Sequences*, Proc. Japan Acad. Ser. A **68** (1992), no. 10, 348–350.
- [3] GROSSWALD, E.—SCHNITZER, F.J.: *A Class of modified ζ and L -Functions*, Pacific Journal of Mathematics **74** (1978), no. 2, 357–364.
- [4] KUIPERS, L.—NIEDERREITER, H.: *Uniform Distribution of Sequences*. Dover Publications, New York 2006.
- [5] MÜLLER, H.: *Über eine Klasse modifizierter ζ - und L -Funktionen*, Arch. Math. **36** (1981), 157–161.
- [6] STRAUCH, O.—PORUBSKÝ, Š.: *Distribution of Sequences: A Sampler*. eBook, 2016. <https://math.boku.ac.at/udt/books/MYBASISNew.pdf>
- [7] TOO, Y.-H.: *On the Uniform Distribution Modulo One of Some Log-like Sequences*, Proc. Japan Acad. Ser. A **68** (1992), no. 9, 269–272.
- [8] PARENT, D. P.: *Exercises in Number Theory*. Springer, New York 1984.

LOG-LIKE FUNCTIONS

- [9] TITCHMARSH, E. C.: *The Theory of the Riemann Zeta-Function*. Oxford University Press, second edition, New York 1986.
- [10] WINTNER, A.: *On the cyclical distribution of the logarithms of the prime numbers*, Quart. J. Math. **6** (1935), no. 1, 65–68.
- [11] ZYGMUND, A.: *Trigonometric Series*. Cambridge University Press, third edition, London 2002.

Received October 20, 2017

Accepted February 2, 2018

Martin Rehberg

Technische Hochschule Mittelhessen,

Department MND

Wilhelm-Leuschner-Str. 13

61169 Friedberg

GERMANY

E-mail: martin.rehberg@mnd.thm.de