
UNSOLVED PROBLEMS

EDITED BY O. STRAUCH

Editorial Board welcome papers containing some progress in problems listed below. Also it welcomes open problems in the line of the aim of this UDT for possible publication in this section. ¹

Notations, definitions and basic properties should be consulted by the following monographs:

L. KUIPERS – H. NIEDERREITER: *Uniform Distribution of Sequences* published by John Wiley in 1974 (reprint edition published by Dover Publications, Inc. Mineola, New York in 2006) and hereafter referred to as [KN];

G. RAUZY: *Propriétés statistiques de suites arithmétiques* published by Presses Universitaires de France in 1976;

E. HLAWKA: *Theorie der Gleichverteilung* published in German by Bibliographisches Institut in 1979 and English under the title *The Theory of Uniform Distribution* by A B Academic Publishers in 1984;

H. NIEDERREITER: *Random Number Generation and Quasi-Monte Carlo Methods* published by SIAM in 1992 and referred to as [N];

M. DRMOTA – R.F. TICHY: *Sequences, Discrepancies and Applications* published by Springer Verlag in 1997 and referred to as [DT];

O. STRAUCH – Š. PORUBSKÝ: *Distribution of Sequences: A Sampler* published by Peter Lang in 2005 and referred to as [SP];

S. TEZUKA: *Uniform Random Numbers. Theory and Practice*, published by Kluwer Academic Publishers in 1995;

J. MATOUŠEK: *Geometric Discrepancy. An Illustrated Guide, Algorithms and Combinatorics* published by Springer Verlag in 1999;

H. NIEDERREITER: *Quasi-Monte Carlo methods and pseudo-random numbers*, Bull. Amer. Math. Soc. **84** (1978), no. 6, 957–1040 (MR 80d:65016);

E. HLAWKA: *Statistik und Gleichverteilung (Statistics and uniform distribution)*, (German), Grazer Math. Ber. **335** (1998), ii+206 pp (MR 99g:11093);

J.F. KOKSMA: *Diophantische Approximationen (Diophantine Approximations)*, (German), published by Springer Verlag in 1936;

J. DICK – F. PILLICHSHAMMER: *Digital Nets and Sequences (Discrepancy Theory and Quasi-Monte Carlo Integration)* published by Cambridge University Press in 2010.

¹Date of a new version: February 9, 2015

Contents

1	Problems	7
1.1	Extended van der Corput difference theorem.	7
1.2	Inverse modulo prime.	7
1.3	Logarithm of primes.	8
1.4	Fractional part of $n\alpha$	10
1.5	Strange recurring sequence.	11
1.6	Function $\pi(n)$	11
1.7	Glasner sets.	12
1.8	Digitally shifted Hammersley sequences.	13
1.9	Block sequence.	15
1.10	Logarithmic and trigonometric functions.	21
1.11	Euler totient function.	24
1.12	van der Corput sequence in the base q	31
1.13	Sequences of differences	37
1.14	Bernoulli numbers	39
1.15	Ratio sequences.	39
1.16	Continued fractions.	40
1.17	Strong uniform distribution.	41
1.18	Algebraic dilatations.	43
1.19	Subsequence ergodic theorems.	43
1.20	Square functions for subsequence ergodic averages.	44
1.21	Sets of integers of positive density	45
1.22	Uniform distribution of the weighted sum-of-digits function.	48
1.23	Moment problem.	49
1.23.1	Truncated Hausdorff moment problem	49
1.23.2	L^2 moment problem	50
1.24	Scalar product.	51
1.25	Determinant.	52
1.26	Ramanujan function.	52

1.27	Apéry sequence.	53
1.28	Urban's conjecture.	53
1.29	Extreme values of $\int_0^1 \int_0^1 F(x, y) d_x d_y g(x, y)$ for copulas $g(x, y)$	55
1.30	Niederreiter-Halton (<i>NH</i>) sequence	60
1.31	Gauss-Kuzmin theorem and $g(x) = g_f(x)$	62
1.32	Benford law	64
1.33	The integral $\int_{[0,1]^s} F(\mathbf{x}, \mathbf{y}) dg(\mathbf{x}) dg(\mathbf{y})$ for d.f.'s $g(\mathbf{x})$	67
1.34	Comparison of random sequences using the game theory	69
1.35	Oscillating sums	71
1.36	Discrepancy system in the unit cube	72
1.37	Extremes of $\int_0^1 \int_0^1 F(x, y) d_x d_y g(x, y)$ attained at shuffles of M	73
1.38	Two-dimensional Benford's law	78
1.39	Problem of N.M. Korobov	81
2	Open theories	83
2.1	Uniform distribution theories.	83
2.2	Distribution functions of sequences.	88
2.3	Deterministic analysis of sequences.	94
2.4	Exponential sequences.	99
2.5	Duffin-Scheaffer conjecture and related sequences.	107

Definitions and notations

• A function $g : [0, 1] \rightarrow [0, 1]$ will be called **distribution function** (abbreviated d.f.) if the following two conditions are satisfied:

- (i) $g(0) = 0, g(1) = 1,$
- (ii) g is non-decreasing.

We shall identify any two distribution functions g, \tilde{g} which coincide at common continuity points, or equivalently, if $g(x) = \tilde{g}(x)$ a.e.

• Given a sequence x_n of real numbers, a positive integer N and a subset I of the unit interval $[0, 1)$, the **counting function** $A(I; N; x_n \bmod 1)$ is defined as the number of terms of x_n with $1 \leq n \leq N$, and with x_n taken modulo one, belonging to I , i.e.

$$A(I; N; x_n \bmod 1) = \#\{n \leq N ; \{x_n\} \in I\} = \sum_{n=1}^N c_I(\{x_n\}),$$

where $c_I(t)$ is the **characteristic function** of I .

• For a sequence $x_1, \dots, x_N \bmod 1$ we define the **step distribution function** $F_N(x)$ for $x \in [0, 1)$ by

$$F_N(x) = \frac{A([0, x); N; x_n \bmod 1)}{N}$$

while $F_N(1) = 1$.

• A d.f. g is called a **distribution function** of the sequence $x_n \bmod 1$ if an increasing sequence of positive integers N_1, N_2, \dots exists such that the equality

$$g(x) = \lim_{k \rightarrow \infty} \frac{A([0, x); N_k; x_n \bmod 1)}{N_k} \left(= \lim_{k \rightarrow \infty} F_{N_k}(x) \right)$$

holds at every point $x, 0 \leq x \leq 1$, of the continuity of $g(x)$ and thus a.e. on $[0, 1]$.

• If there exists a limit $\lim_{N \rightarrow \infty} F_N(x) = g(x)$ a.e. on $[0, 1]$, then $g(x)$ is called **asymptotic distribution function** (abbreviating a.d.f.) of $x_n \bmod 1$ and if $g(x) = x$ then $x_n \bmod 1$ is called **uniformly distributed** in $[0, 1]$ (abbreviating u.d.)

- The **set of all distribution functions** of a sequence $x_n \bmod 1$ will be denoted by $G(x_n \bmod 1)$. We shall identify the notion of the **distribution** of a sequence $x_n \bmod 1$ with the set $G(x_n \bmod 1)$, i.e. the distribution of $x_n \bmod 1$ is known if we know the set $G(x_n \bmod 1)$. The set $G(x_n \bmod 1)$ has the following fundamental properties for every sequence $x_n \bmod 1$:
 - $G(x_n \bmod 1)$ is non-empty, and it is either a singleton or has infinitely many elements,
 - $G(x_n \bmod 1)$ is closed and connected in the topology of the weak convergence, and these properties are characteristic, i.e.
 - for given a non-empty set H of distribution functions, there exists a sequence x_n in $[0, 1)$ such that $G(x_n) = H$ if and only if H is closed and connected.
- Let x_1, \dots, x_N be a given sequence of real numbers from the unit interval $[0, 1)$. Then the number

$$D_N = D_N(x_1, \dots, x_N) = \sup_{0 \leq \alpha < \beta \leq 1} \left| \frac{A([\alpha, \beta); N; x_n]}{N} - (\beta - \alpha) \right|$$

is called the **(extremal) discrepancy** of this sequence. The number

$$D_N^* = \sup_{x \in [0, 1]} \left| \frac{A([0, x); N; x_n]}{N} - x \right|$$

is called **star discrepancy**, and the number

$$D_N^{(2)} = \int_0^1 \left(\frac{A([0, x); N; x_n]}{N} - x \right)^2 dx$$

is called its **L^2 discrepancy**.

- For multidimensional case see 2.2, p. 88.
- The Riemann-Stiltjes integral $\int_0^1 \int_0^1 f(x, y) d_x d_y g(x, y)$ is defined as the limit

$$\begin{aligned} & \sum_{k=1}^m \sum_{l=1}^n f(\alpha_k, \beta_l) (g(x_k, y_l) + g(x_{k+1}, y_{l+1}) - g(x_k, y_{l+1}) - g(x_{k+1}, y_l)) \\ & \rightarrow \int_0^1 \int_0^1 f(x, y) d_x d_y g(x, y) \end{aligned}$$

if diameters of $[x_k, x_{k+1}] \times [y_l, y_{l+1}]$ tend to zero for partition $0 = x_0 < x_1 < \dots < x_m = 1$ of x -axis, $0 = y_0 < y_1 < \dots < y_n = 1$ of y -axis and for $(\alpha_k, \beta_l) \in [x_k, x_{k+1}] \times [y_l, y_{l+1}]$. This integral exists for continuous

$f(x, y)$ and $g(x, y)$ with bounded variation. Let \square denote the rectangle $[x_k, x_{k+1}] \times [y_l, y_{l+1}]$ and denote

$$\square g(x, y) = g(x_k, y_l) + g(x_{k+1}, y_{l+1}) - g(x_k, y_{l+1}) - g(x_{k+1}, y_l).$$

If diameter $\square \rightarrow 0$, then we find the differential $d_x d_y g(x, y)$ as

$$d_x d_y g(x, y) = g(x, y) + g(x + dx, y + dy) - g(x, y + dy) - g(x + dx, y).$$

In some cases we shorten $d_x d_y g(x, y) = dg(x, y)$.

1. Problems

1.1 Extended van der Corput difference theorem.

Prove or disprove: If the sequence

$$k(x_{n+h} - x_n) - h(x_{n+k} - x_n) \pmod{1}, \quad n = 1, 2, \dots,$$

is u.d. for every $k, h = 1, 2, \dots, k > h$, then the original sequence

$$x_n \pmod{1}, \quad n = 1, 2, \dots,$$

is also

u.d.

NOTES: This problem was posed by M.H. Huxley at the Conference on Analytic and Elementary Number Theory, Vienna, July 18–20, 1996.

Entered by O. Strauch.

1.2 Inverse modulo prime.

Let $p > 2$ be a prime number. For an integer $0 < n < p$, define n^* by the congruence $nn^* \equiv 1 \pmod{p}$, $0 < n^* < p$. Is it true that the sequence of blocks

$$\left(\frac{n^*}{p}, \frac{(n+1)^*}{p} \right), \quad n = 1, 2, \dots, p-2,$$

is u.d. as $p \rightarrow \infty$?

NOTES: Tsz Ho, Chan (2004) proved that

$$\frac{1}{p} \sum_{n=1}^{p-2} \left| \frac{n^*}{p} - \frac{(n+1)^*}{p} \right| = \frac{1}{3} + \mathcal{O}\left(\frac{(\log p)^3}{\sqrt{p}}\right)$$

for every prime $p > 2$. Moreover, the sequence

$$\left(\frac{n}{p}, \frac{n^*}{p} \right), \quad n = 1, 2, \dots, p-1,$$

is u.d. as $p \rightarrow \infty$,

see [SP, p. 3–25, 3.7.2].

Solution: The s -dimensional sequence

$$\left(\frac{n^*}{p}, \frac{(n+1)^*}{p}, \dots, \frac{(n+s-1)^*}{p} \right), \quad n = 1, 2, \dots, p,$$

is u.d. as $p \rightarrow \infty$,

and the discrepancy bound is

$$D_p^* = \mathcal{O}\left(\frac{(\log p)^s}{\sqrt{p}}\right) \quad (1)$$

for all $s \geq 2$, and this estimate is essentially best possible up to the logarithmic factor.

NOTES: A. Winterhof sent to us that (1) was proved by H. Niederreiter (1994). A generalization is given in H. Niederreiter and A. Winterhof (2000).

Proposed by O. Strauch.

TSZ HO CHAN : *Distribution of difference between inverses of consecutive integers modulo p* , *Integers* **4** (2004), A3, 11 pp. (MR2056009(2005b:11159)).

H. NIEDERREITER: *Pseudorandom vector generation by the inverse method*, *ACM Trans. Model. Comput. Simul.* **4** (1994), 2, 191–212 (Zbl 0847.11039).

H. NIEDERREITER – A. WINTERHOF: *Incomplete exponential sums over finite fields and their applications to new inverse pseudorandom number generators*, *Acta Arith.* **XCIII** (2000), 4, 387–399.

1.3 Logarithm of primes.

See [SP, p.2–175, 2.19.8]. Let p_n be the n th prime. Find the set of all distribution functions $G(x_n)$ of the sequence

$$x_n = \log p_n \bmod 1, \quad n = 1, 2, \dots$$

NOTES: (I) A. Wintner (1935) has shown that $\log p_n \bmod 1$ is not u.d. A proof can be found in D.P. Parent (1984, pp. 282–283, Solut. 5.19).

(II) S. Akiyama (1996, 1998) proved: Let c_i , $i = 0, 1, 2, \dots, k-1$, be real numbers with $\sum_{i=0}^{k-1} c_i \neq 0$. Then the sequence $x_n = \sum_{i=0}^{k-1} c_i \log p_{n+i} \bmod 1$, $n = 1, 2, \dots$ is not almost u.d., i.e. $x \notin G(x_n)$.

(II) R.E. Whitney (1972) proved that $\log p_n \bmod 1$ is u.d. with respect to the logarithmic weighted means, i.e.

$$\lim_{N \rightarrow \infty} \left(\sum_{n=1}^N \frac{1}{n} \right)^{-1} \sum_{n=1}^N \frac{c_{[0,x]}(\{\log p_n\})}{n} = x$$

for all $x \in [0, 1]$.

(III) D.I.A. Cohen and T.M. Katz (1984) have shown the u.d. of $\log p_n \bmod 1$ with respect to the zeta distribution, i.e.

$$\lim_{\alpha \rightarrow 1^+} \frac{1}{\zeta(\alpha)} \sum_{n=1}^{\infty} \frac{c_{[0,x]}(\{\log p_n\})}{n^\alpha} = x$$

for all $x \in [0, 1]$.

Solution. Y. Ohkubo (2011) proved the following results (i)–(ix):

(i) Two sequences $\log p_n \bmod 1$ and $\log n \bmod 1$ have the same d.f.s, i.e.

$$G(\log p_n \bmod 1) = G(\log n \bmod 1).$$

(ii) Every u.d. sequence $x_n \bmod 1$ is statistically independent of $\log p_n \bmod 1$, i.e., $x_n \bmod 1$ and $(x_n + \log p_n) \bmod 1$ are u.d. simultaneously.

(iii) The result (ii) follows from that every u.d. sequence $x_n \bmod 1$ is statistically independent with $\log(n \log n) \bmod 1$ and

$$\lim_{n \rightarrow \infty} (\log p_n - \log(n \log n)) = 0.$$

(iv) The result (ii) implies that, for every irrational θ the sequence $p_n \theta + \log p_n$ is u.d. $\bmod 1$.

(v) Also, every u.d. sequence $x_n \bmod 1$ is statistically independent with $\frac{p_n}{n} \bmod 1$. It follows from the limit

$$\lim_{n \rightarrow \infty} \left(\frac{p_n}{n} - \log(n \log n) \right) = -1.$$

(vi) The result (v) implies that $p_n \theta + \frac{p_n}{n}$ is u.d. $\bmod 1$.

(vii) Theorem: Let the real-valued function $f(x)$ be strictly increasing for $x \geq 1$ and let $f^{-1}(x)$ be the inverse function of $f(x)$. Suppose that

- $\lim_{k \rightarrow \infty} f^{-1}(k+1) - f^{-1}(k) = \infty$,
- $\lim_{k \rightarrow \infty} \frac{f^{-1}(k+w_k)}{f^{-1}(k)} = \psi(u)$ for every sequence $w_k \in [0, 1]$

for which $\lim_{k \rightarrow \infty} w_k = u$, where this limit defines the function $\psi(u)$ on $[0, 1]$,

- $\psi(1) > 1$. Then

$$G(f(p_n) \bmod 1) = \left\{ \tilde{g}_u(x) = \frac{\min(\psi(x), \psi(u)) - 1}{\psi(u)} + \frac{1}{\psi(u)} \cdot \frac{\psi(x) - 1}{\psi(1) - 1} : u \in [0, 1] \right\}.$$

(viii) The result (vii) implies that $\log p_n$ and the sequences $\log(p_n \log^{(i)} p_n)$, $i = 1, 2, \dots$ have the same distribution as the sequence $\log n$.

(ix) S. Akiyama (1998) proved

$$\lim_{n \rightarrow \infty} \left(\sum_{i=0}^{\ell-1} c_i \log p_{n+i} - \left(\sum_{i=0}^{\ell-1} c_i \right) \log p_n \right) = 0.$$

Then

$$G\left(\sum_{i=0}^{\ell-1} c_i \log p_{n+i} \bmod 1\right) = G\left(\left(\sum_{i=0}^{\ell-1} c_i\right) \log p_n \bmod 1\right).$$

and by (vii) we have

$$G(c \log p_n \bmod 1) = \left\{ \frac{e^{x/c} - 1}{e^{1/c} - 1} e^{-u/c} + (e^{\min(x/c, u/c)} - 1) e^{-u/c} : u \in [0, 1] \right\},$$

where $c = \sum_{i=0}^{\ell-1} c_i$.

Entered by O. Strauch.

S. AKIYAMA: *A remark on almost uniform distribution modulo 1*, in: *Analytic number theory (Japanes)* (Kyoto, 1994), Sürikaisekikenkyūsho Kōkyūroku No. 958, 1996, pp. 49–55 (MR 99b:11081).

S. AKIYAMA: *Almost uniform distribution modulo 1 and the distribution of primes*, Acta Math. Hungar. **78** (1998), no. 1–2, 39–44 (MR 99b:11083).

D.I.A. COHEN – T.M. KATZ: *Prime numbers and the first digit phenomenon*, J. Number Theory **18** (1984), 261–268 (MR 85j:11014).

D.P. PARENT: *Exercises in Number Theory*, Problem Books in Mathematics, Springer Verlag, New York, 1984 (French original: *Exercices de théorie des nombres*, Gauthier – Villars, Paris, 1978) (86f:11002).

Y. OHKUBO: *On sequences involving primes*, Uniform Distribution Theory **6** (2011), no. 2, 221–238.

R.E. WHITNEY: *Initial digits for the sequence of primes*, Amer. Math. Monthly **79** (1972), no. 2, 150–152 (MR 46#3472).

A. WINTNER: *On the cyclical distribution of the logarithms of the prime numbers*, Quart. J. Math. Oxford (1) **6** (1935), 65–68 (Zbl 11, 149).

1.4 Fractional part of $n\alpha$.

See [SP, p. 2–86, 2.8.12]. Characterize the set $G(x_n)$ of all d.f.'s of the sequence

$$x_n = \begin{cases} \{n\alpha\}\alpha, & \text{if } \{n\alpha\} < 1 - \alpha, \\ (1 - \{n\alpha\})(1 - \alpha), & \text{if } \{n\alpha\} \geq 1 - \alpha, \end{cases}$$

for $0 < \alpha < 1$.

NOTES: A.F. Timan (1987) proved that the series $\sum_{n=1}^{\infty} \frac{x_n}{n^r}$ converges for all $\alpha \in (0, 1)$ if and only if $r > 1$.

Solution. S. Steinerberger: For irrational $0 < \alpha < 1$ we have $x_n = f(\{n\alpha\})$, where

$$f(x) = \begin{cases} x\alpha, & \text{if } x \in [0, 1 - \alpha], \\ (1 - x)(1 - \alpha), & \text{if } x \in [1 - \alpha, 1]. \end{cases}$$

Then a.d.f. $g(x)$ of x_n is

$$g(x) = |f^{-1}([0, x])| = \begin{cases} 1, & \text{if } x \in [\alpha(1 - \alpha), 1], \\ \frac{x}{\alpha(1 - \alpha)}, & \text{others.} \end{cases}$$

Proposed by O. Strauch.

A.F. TIMAN: *Distribution of fractional parts and approximation of functions with singularities by Bernstein polynomials*, J. Approx. Theory **50** (1987), no. 2, 167–174 (MR 88m:11054).

1.5 Strange recurring sequence.

[SP, p. 2–243, 2.24.10]: Characterize the $G(x_n)$ of the so-called strange recurring sequences of the form

$$(i) \quad x_n = x_{n-[x_{n-1}]} + x_{n-[x_{n-2}]},$$

$$(ii) \quad x_n = x_{n-[x_{n-1}]} + x_{[x_{n-1}]},$$

$$(iii) \quad x_n = x_{[x_{n-2}]} + x_{n-[x_{n-2}]},$$

with real initial values x_1, x_2 .

NOTES: If $x_1 = x_2 = 1$, the sequence (i) was defined by D.R. Hofstadter (1979), (ii) was defined by J.H. Conway (1988) during one of his lectures and C.L. Mallows (1991) established the regular structure of (ii) and introduced the monotone sequence (iii).

Proposed by O. Strauch.

D.R. HOFSTADTER: *Gödel, Escher, Bach: an External Golden Braid*, Basic Books, Inc., Publishers, New York, 1979 (MR 80j:03009).

C.L. MALLOWS: *Conway's challenge sequence*, Amer. Math. Monthly **98** (1991), no. 1, 5–20 (MR 92e:39007).

1.6 Function $\pi(n)$.

[SP, p. 2–193]: Riemann hypothesis implies that the sequence

$$\frac{n}{\pi(n)} \bmod 1, \quad n = 1, 2, \dots,$$

is not u.d. Find all its d.f.'s.

NOTES: Under the Riemann hypothesis $\pi(x) = \text{li}(x) + \mathcal{O}(\sqrt{x} \log x)$ which implies $\lim_{n \rightarrow \infty} (n/\pi(n)) - (n/\text{li}(n)) = 0$ the sequences $n/\pi(n) \pmod{1}$ and $n/\text{li}(n) \pmod{1}$ have the same d.f.s if we prove the continuity of all d.f.'s of $n/\text{li}(n) \pmod{1}$ at 0 and 1, cf. [SP, p. 2–24, 2.3.3]. Niederreiter's theorem: If $x_n, n = 1, 2, \dots$, is a monotone sequence that is u.d. mod 1, then

$$\lim_{n \rightarrow \infty} \frac{|x_n|}{\log n} = \infty.$$

implies that the sequence $n/\pi(n) \pmod{1}$ is not u.d. (probably without the Riemann hypothesis).

Solution: Without the Riemann hypothesis sequences $n/\pi(n)$ and $\log n$ have the same d.f.'s mod 1.

NOTES: According to F. Luca this follows from

$$\left| \frac{n}{\pi(n)} - \frac{n}{\text{li}(n)} \right| = \mathcal{O}((\log n)^2 \exp(-c\sqrt{\log n})) = o(1);$$

$$\left| \frac{n}{\text{li}(n)} - \frac{n}{f(n)} \right| = \mathcal{O}((\log n)^{-1}) = o(1), \text{ where } f(n) = \frac{n}{\log n} + \frac{n}{(\log n)^2};$$

$$\frac{n}{f(n)} = \log(n) - 1 + o(1);$$

immediately.

Proposed by O. Strauch.

H. NIEDERREITER: *Distribution mod 1 of monotone sequences*, Nederl. Akad. Wetensch. Indag. Math. **46** (1984), no. 3, 315–327 (MR 86i:11041).

1.7 Glasner sets.

A strictly increasing sequence of positive integers k_n , $n = 1, 2, \dots$ is called a **Glasner set** if for every infinite set $A \subset [0, 1)$ and every $\varepsilon > 0$ there exists k_n such that the dilation $k_n A \bmod 1 = \{k_n x \bmod 1 : x \in A\}$ is ε -dense in $[0, 1]$, i.e., $k_n A \bmod 1$ intersects every subinterval of $[0, 1]$ of the length ε . The following sequences k_n , $n = 1, 2, \dots$, are Glasner sets:

- (i) $k_n = n$.
- (ii) $k_n = P(n)$, where $P(x)$ is a non-constant polynomial with integer coefficients.
- (iii) $k_n = P(p_n)$, where p_n is the increasing sequence of all primes and polynomial $P(x)$ is as in (ii).

A strictly increasing sequence of positive integers k_n , $n = 1, 2, \dots$, has **quantitative Glasner property** if for every given $\varepsilon > 0$ there exists an integer $s(\varepsilon)$ such that for any finite set $A \subset [0, 1)$ of cardinality at least $s(\varepsilon)$ there exists k_n such that the dilation $k_n A \bmod 1$ is ε -dense in $[0, 1)$. The following sequences k_n , $n = 1, 2, \dots$, have this property:

- (iv) $k_n = n$ as in (i) with $s(\varepsilon) = \lceil \varepsilon^{-2-\gamma} \rceil$, where $\gamma > 0$ is arbitrary and $\varepsilon \leq \varepsilon_0(\gamma)$.
- (v) $k_n = P(n)$, where $P(x)$ is a non-constant polynomial with integer coefficients.
- (vi) $k_n = P(p_n)$ as in (iii) with $s(\varepsilon) = \lceil \varepsilon^{-2d-\delta} \rceil$, where $d = \deg P(x)$, $\delta > 0$ arbitrary and $\varepsilon < \varepsilon_0(P(x), \delta)$.
- (vii) k_n , $n = 1, 2, \dots$, is: (*) uniformly distributed for each positive integer m (i.e., for each $i = 0, 1, \dots, m-1$ the relative density of $k_m \equiv i \pmod{m}$ is $1/m$), and (**) for each irrational α , the sequence $k_n \alpha \bmod 1$ is uniformly distributed in $[0, 1]$. Here $s(\varepsilon) =$

$[\varepsilon^{-2-3(\log \log(1/\varepsilon))^{-1}}] + 1$, for every $\varepsilon < \varepsilon_0$, where ε_0 depends on the sequence k_n , $n = 1, 2, \dots$.

(viii) $k_n = [f(n)]$, where $f(x)$ denotes a non-polynomial entire function that is real on the real numbers and such that $|f(z)| = O(e^{(\log |z|)^\alpha})$ with $\alpha < 4/3$ and $s(\varepsilon)$ is as in (vii).

(ix) $k_n = [f(p_n)]$, where f is as in (viii) and $s(\varepsilon)$ is as in (vii).

(x) $k_n = [n^\alpha]$ for any $\alpha \geq 1$ not an integer ≥ 2 and $s(\varepsilon)$ is as in (vii).

Open problem: For $\mathbf{x} = (x_1, \dots, x_n) \in [0, 1]^N$ and positive integers $k_1 < k_2 < \dots < k_K$ define the N -dimensional sequence $k_1\mathbf{x}, k_2\mathbf{x}, \dots, k_K\mathbf{x}$ and let $D_K^{(2)}(k_n\mathbf{x})$ be its L^2 discrepancy. Generalizing O. Strauch (1989), H. Albrecher (2002) (cf. [SP, p. 3–14]) proved, for the mean value of the L^2 discrepancy $D_N^{(2)}(k_n\mathbf{x})$, that

$$\int_{[0,1]^s} D_N^{(2)}(k_n\mathbf{x}) \, d\mathbf{x} = \sum_{m,n=1}^K \left(\frac{1}{3} + \frac{1}{12} \frac{(k_m, k_n)^2}{k_m k_n} \right)^N + \left(\frac{1}{2^N} - \left(\frac{5}{12} \right)^N \right) - \frac{1}{3^N},$$

where (k_m, k_n) is a g.c.d. of k_m and k_n . Find some connection between Glasner sets and mean values of such L^2 discrepancy.

Proposed by O. Strauch.

H. ALBRECHER: *Metric distribution results for sequences* ($\{q_n \vec{\alpha}\}$), *Math. Slovaca* **52** (2002), no. 2, 195–206 (MR 2003h:11083; Zbl 1005.11036).

N. ALON – Y. PERES: *Uniform dilations*, *Geom. Funct. Anal.* **2** (1992), no. 1, 1–28 (MR 93a:11061; Zbl 0756.11020).

D. BEREND – Y. PERES: *Asymptotically dense dilations of sets on the circle*, *J. Lond. Math. Soc.*, II. Ser. **47** (1993), no. 1, 1–17 (MR 94b:11068; Zbl 0788.11028).

S. GLASNER: *Almost periodic sets and measures on the torus*, *Israel J. Math* **32** (1979), no. 2–3, 161–172 (MR 80f:54038; Zbl 0406.54023).

H.H. KAMARUL – R. NAIR: *On certain Glasner sets*, *Proc. R. Soc. Edinb.*, Sect. A, *Math.* **133** (2003), no. 4, 849–853 (Zbl prev02064747).

R. NAIR: *On asymptotic distribution on the a -adic integers*, *Proc. Indian Acad. Sci.*, *Math. Sci.* **107** (1997), no. 4, 363–376 (MR 98k:11110; Zbl 0908.11036).

R. NAIR – S.L. VELANI: *Glasner sets and polynomials in primes*, *Proc. Amer. Math. Soc.* **126** (1998), no. 10, 2835–2840 (MR 99a:11095; Zbl 0913.11031).

O. STRAUCH: *Some applications of Franel – Klyuyver’s integral, II*, *Math. Slovaca* **39** (1989), 127–140 (MR 90j:11079).

1.8 Digitally shifted Hammersley sequences.

Let $x = 0.x_1x_2\dots x_m$ and $y = 0.y_1y_2\dots y_m$ be two real numbers written in dyadic expansion. Define $x \oplus y = z = 0.z_1z_2\dots z_m$, where $z_i = x_i + y_i$

(mod 2), $i = 1, 2, \dots, m$. Let $\gamma_2(n)$ be the van der Corput radical inverse function defined by $\gamma_2(n) = 0.a_0a_1 \dots a_{m-1}$, where $n = a_{m-1}a_{m-2} \dots a_0$ is a positive integer (again in dyadic expansion). Then for the L^2 discrepancy $D_N^{(2)}$ of the sequence

$$\left(\frac{n}{N}, \gamma_2(n) \oplus x \right), \quad n = 0, 1, \dots, N-1, \quad \text{with } N = 2^m$$

we have

$$\frac{m^2}{64} - \frac{19m}{192} - \frac{lm}{16} + \frac{l^2}{16} + \frac{5}{16} + \frac{m}{8 \cdot 2^m} - \frac{l}{4 \cdot 2^m} + \frac{5}{16 \cdot 2^m} - \frac{1}{72 \cdot 4^m} \leq N^2 D_N^{(2)},$$

$$N^2 D_N^{(2)} \leq \frac{m^2}{64} - \frac{19m}{192} - \frac{lm}{16} + \frac{l^2}{16} + \frac{l}{4} + \frac{7}{16} + \frac{m}{8 \cdot 2^m} - \frac{l}{4 \cdot 2^m} + \frac{3}{16 \cdot 2^m} - \frac{1}{72 \cdot 4^m},$$

where l denotes the number of zeros in the dyadic expansion of x . If m is even and $l = m/2$, then

$$D_N^{(2)} = \mathcal{O} \left(\frac{\log N}{N^2} \right)$$

which is the best possible. A similar situation holds in the case of odd m and $l = (m-1)/2$.

Problem: Find an exact formula for $N^2 D_N^{(2)}$.

NOTES: (I) See P. Kritzer and F. Pillichshammer (2005, Th. 2 and 3) for L^2 discrepancy bounds.

(II) For the L^2 discrepancy of the 2-dimensional Hammersley sequence (also called Roth sequence)

$$\left(\frac{n}{N}, \gamma_2(n) \right), \quad n = 0, 1, \dots, N-1, \quad N = 2^m$$

the following exact formula

$$N^2 D_N^{(2)} = \frac{m^2}{64} + \frac{29m}{192} + \frac{3}{8} - \frac{m}{16 \cdot 2^m} + \frac{1}{4 \cdot 2^m} - \frac{1}{72 \cdot 2^{2m}}.$$

was proved by I.V. Vilenkin (1967) and independently by J.H. Halton and S.K. Zaremba (1969).

Solution: According to P. Kritzer and F. Pillichshammer (2006)

$$N^2 D_N^{(2)} = \frac{m^2}{64} - \frac{19m}{192} - \frac{lm}{16} + \frac{l^2}{16} + \frac{l}{4} + \frac{3}{8} + \frac{m}{16 \cdot 2^m} - \frac{l}{8 \cdot 2^m} + \frac{1}{4 \cdot 2^m} - \frac{1}{72 \cdot 4^m}.$$

Proposed by O. Strauch.

J.H. HALTON – S.K. ZAREMBA: *The extremal and L^2 discrepancies of some plane set*, *Monatsh. Math.* **73** (1969), 316–328 (MR **40**#5550).

P. KRITZER – F. PILLICHSHAMMER : *Point sets with low L_p -discrepancy*, *Math. Slovaca* **57** (2007), no. 1, 11–32 (MR2357804(2009a:11155)).

P. KRITZER – F. PILLICHSHAMMER : *An exact formula for the L_2 of the shifted Hammersley point set*, *Uniform Distribution Theory* **1** (2006), no. 1, 1–13 (MR2314263(2008d:11084)).

I.V. VILENKIN: *Plane sets of integration*, (Russian), *Z. Včisl. Mat. i Mat. Fiz.* **7** (1967), 189–196; English translation: *USSR Comp. Math. and Math. Phys.* **7** (1967), 258–267 (MR 0205464(34#5291)).

1.9 Block sequence.

Let $x_n, n = 1, 2, \dots$ be an increasing sequence of positive integers, $\underline{d}(x_n)$ be the lower asymptotic density, $\bar{d}(x_n)$ be the upper asymptotic density of $x_n, n = 1, 2, \dots$, and $X_n = \left(\frac{x_1}{x_n}, \frac{x_2}{x_n}, \dots, \frac{x_n}{x_n}\right)$. Let $G(X_n)$ be the set of all d.f.'s of the block sequence $X_n, n = 1, 2, \dots$, i.e., the set of all possible weakly limits $F(X_{n_k}, x) \rightarrow g(x)$ as $k \rightarrow \infty$, where

$$F(X_{n_k}, x) = \frac{\#\{i \leq n_k; x_i/x_{n_k} < x\}}{n_k}.$$

$G(X_n)$ has the following properties:

- (i) If $g(x) \in G(X_n)$ increases and is continuous at $x = \beta$ and $g(\beta) > 0$, then there exists $1 \leq \alpha < \infty$ such that $\alpha g(x\beta) \in G(X_n)$. If every d.f. of $G(X_n)$ is continuous at 1, then $\alpha = 1/g(\beta)$.
- (ii) Assume that all d.f.'s in $G(X_n)$ are continuous at 0 and $c_1(x) \notin G(X_n)$. Then for every $\tilde{g}(x) \in G(X_n)$ and every $1 \leq \alpha < \infty$ there exists $g(x) \in G(X_n)$ and $0 < \beta \leq 1$ such that $\tilde{g}(x) = \alpha g(x\beta)$ a.e.
- (iii) Assume that all d.f.s in $G(X_n)$ are continuous at 1. Then all d.f.'s in $G(X_n)$ are continuous on $(0, 1]$, i.e., only possible discontinuity is in 0.
- (iv) If $\underline{d}(x_n) > 0$, then for every $g(x) \in G(X_n)$ we have $(\underline{d}(x_n)/\bar{d}(x_n)).x \leq g(x) \leq (\bar{d}(x_n)/\underline{d}(x_n)).x$ for every $x \in [0, 1]$. Thus $\underline{d}(x_n) = \bar{d}(x_n) > 0$ implies u.d. of the block sequence $X_n, n = 1, 2, \dots$.
- (v) If $\underline{d}(x_n) > 0$, then every $g(x) \in G(X_n)$ is continuous on $[0, 1]$.
- (vi) If $\underline{d}(x_n) > 0$, then there exists $g(x) \in G(X_n)$ such that $g(x) \geq x$ for every $x \in [0, 1]$.
- (vii) If $\bar{d}(x_n) > 0$, then there exists $g(x) \in G(X_n)$ such that $g(x) \leq x$ for every $x \in [0, 1]$.
- (viii) Assume that $G(X_n)$ is singleton, i.e., $G(X_n) = \{g(x)\}$. Then either $g(x) = c_0(x)$ for $x \in [0, 1]$; or $g(x) = x^\lambda$ for some $0 < \lambda \leq 1$ and $x \in [0, 1]$. Moreover, if $\bar{d}(x_n) > 0$, then $g(x) = x$.

- (ix) $\max_{g \in G(X_n)} \int_0^1 g(x) dx \geq \frac{1}{2}$.
- (x) Assume that every d.f. $g(x) \in G(X_n)$ has a constant value on the fixed interval $(u, v) \subset [0, 1]$ (maybe different). If $\underline{d}(x_n) > 0$, then all d.f.'s in $G(X_n)$ has infinitely many intervals with constant values.
- (xi) There exists an increasing sequence $x_n, n = 1, 2, \dots$, of positive integers such that $G(X_n) = \{h_\alpha(x); \alpha \in [0, 1]\}$, where $h_\alpha(x) = \alpha, x \in (0, 1)$ is the constant d.f.
- (xii) There exists an increasing sequence $x_n, n = 1, 2, \dots$, of positive integers such that

$$c_1(x) \in G(X_n) \text{ but } c_0(x) \notin G(X_n)$$

where $c_0(x)$ and $c_1(x)$ are one-jump d.f.'s with the jump of height 1 at $x = 0$ and $x = 1$, respectively.

- (xiii) There exists an increasing sequence $x_n, n = 1, 2, \dots$, of positive integers such that $G(X_n)$ is non-connected.
- (xiv) $G(X_n) = \{x^\lambda\}$ if and only if $\lim_{n \rightarrow \infty} (x_{k.n}/x_n) = k^{1/\lambda}$ for every $k = 1, 2, \dots$. Here as in (viii) we have $0 < \lambda \leq 1$.
- (xv) If $\underline{d}(x_n) > 0$, then all d.f.s $g(x) \in G(X_n)$ are continuous, nonsingular and bounded by $h_1(x) \leq g(x) \leq h_2(x)$, where

$$h_1(x) = \begin{cases} x \frac{\underline{d}}{\bar{d}} & \text{if } x \in \left[0, \frac{1-\bar{d}}{1-\underline{d}}\right], \\ \frac{\underline{d}}{\frac{1}{x} - (1-\underline{d})} & \text{otherwise,} \end{cases} \quad h_2(x) = \min \left(x \frac{\bar{d}}{\underline{d}}, 1 \right).$$

Furthermore $h_1(x)$ and $h_2(x)$ are optimal and $h_1(x) \notin G(X_n)$.

NOTES: The properties (i)–(x) can be found in O. Strauch and J.T. Tóth (2001, 2002); (xi), (xiii) in G. Grekos and O. Strauch (2007); (xii) was found by L. Mišík (2004, personal communication); (xiv) is in F. Filip and J.T. Tóth (2006); (xv) is in V. Baláž, L. Mišík, J.T. Tóth and O. Strauch (2009). For concrete examples, cf. [SP, p. 2–217, 2.22.6; p. 2–219, 2.22.7; p. 2–222, 2.22.8; p. 2–225, 2.22.9, 2.22.10; p. 2–226, 2.22.11].

Methods:

(I) **Z-transform.** For positive integers $x_0 < x_1 < x_2 < \dots$ we can assign the complex function $f(z) = \sum_{n=0}^{\infty} \frac{x_n}{z^n}$. The following holds:

- (i) $x_n \rightarrow f(z) = \sum_{n=0}^{\infty} \frac{x_n}{z^n}$;
- (ii) $x_0 + x_1 + \dots + x_{n-1} \rightarrow \frac{f(z)}{z-1}$;
- (iii) $x_{n+1} - x_n \rightarrow (z-1)f(z) - zx_0$;
- (iv) $\frac{x_n}{n} \rightarrow \int_z^{\infty} \frac{f(\xi)}{\xi} d\xi$;
- (v) $nx_n \rightarrow -z \frac{d}{dz} f(z)$;
- (vi) $(n-1)x_{n-1} \rightarrow -\frac{d}{dz} f(z)$;
- (vii) If $x_n \rightarrow f(z)$ and $y_n \rightarrow g(z)$, then for convolution $x_n * y_n = z_n$, where $z_n = x_0 y_n + x_1 y_{n-1} + \dots + x_n y_0$ we have

$$x_n * y_n \longrightarrow f(z) \cdot g(z);$$

If $f(z)$ is known, then there exists inverse transform

$$(viii) \quad x_n = \frac{1}{2\pi i} \oint_C f(z) z^{n-1} dz = \sum_{i=1}^k \operatorname{res}_{z=z_i} f(z) z^{n-1};$$

$$(ix) \quad \frac{x_0 + x_1 + \dots + x_{n-1}}{(n-1)x_{n-1}} = \frac{\sum_{i=1}^k \operatorname{res}_{z=z_i} \frac{f(z)}{z-1} z^{n-1}}{\sum_{i=1}^k \operatorname{res}_{z=z_i} \left(-\frac{d}{dz} f(z)\right) z^{n-1}} = \int_0^1 x dF(X_n, x).$$

Problem: Using \mathcal{Z} -transform (vii) and (ix) for a study of $G(Z_n)$, where $z_n = x_n * y_n$.

(II) **Algorithm.** [V. Baláž, L. Mišík, O. Strauch and J.T. Tóth (2008)]: Let x_n , $n = 1, 2, \dots$ be an increasing sequence of positive integers. Put $x_0 = 0$ and $t_n = x_n - x_{n-1}$, $n = 1, 2, \dots$. For every $n = 1, 2, \dots$, from t_n we compute the finite sequence $t_1^{(n)}, t_2^{(n)}, \dots, t_n^{(n)}$ by the following procedure:

1⁰. For $n = 1$, $t_1^{(1)} = t_1 = x_1$;

2⁰. For $n = 2$, $t_1^{(2)} = t_1 + t_2 - 1 = x_2 - 1$ and $t_2^{(2)} = 1$;

3⁰. Assume that for $n - 1$ we have $t_i^{(n-1)}$, $i = 1, 2, \dots, n - 1$, for n we put $t'_i = t_i^{(n-1)}$, $i = 1, 2, \dots, n - 1$, and $t'_n = t_n$. The following steps (a) and (b) produce new t'_1, \dots, t'_n .

(a) If there exists k , $1 \leq k < n$, such that $t'_1 = t'_2 = \dots = t'_{k-1} > t'_k$ and $t'_n > 1$, then we put $t'_k := t'_k + 1$, $t'_n := t'_n - 1$ and $t'_i := t'_i$ in all other cases.

(b) If such k does not exist and $t'_n > 1$, then we put $t'_1 := t'_1 + 1$, $t'_n := t'_n - 1$ and $t'_i := t'_i$ in all other cases.

In the n th step we will repeat (a) and (b) and the algorithm ends, if $t'_n = 1$ and which gives the resulting $t_1^{(n)} := t'_1, \dots, t_n^{(n)} := t'_n$.

Assuming that $t_n \neq 1$ for infinitely many n , these $t_i^{(n)}$, $i = 1, 2, \dots, n$ can have two possible forms:

$$(A) t_1^{(n)} = \dots = t_m^{(n)} = D_n > t_{m+1}^{(n)} \geq t_{m+2}^{(n)} = t_{m+3}^{(n)} = \dots = t_n^{(n)} = 1,$$

$$(B) t_1^{(n)} = \dots = t_m^{(n)} = D_n > t_{m+1}^{(n)} = \dots = t_{m+s}^{(n)} = D_n - 1 \geq t_{m+s+1}^{(n)} = \dots = t_n^{(n)} = 1,$$

where $m = m(n)$, $s = s(n)$, $D_1 \leq D_2 \leq \dots$ and $D_n \geq 2$ starting from n with $t_n > 1$. Thus there are two possibilities:

(I) D_n is bounded;

(II) $D_n \rightarrow \infty$.

In the case (I) we have only the form (A) and $D_n = \text{const.} = c \geq 2$ for all sufficiently large n .

In the case (II) both cases (A) and (B) are possible. Further properties:

- $x_n = \sum_{i=1}^n t_i^{(n)}$ for $n = 1, 2, \dots$

- Denoting $x_j^{(n)} = \sum_{i=1}^j t_i^{(n)}$, then we have $x_j \leq x_j^{(n)}$ for $j = 1, 2, \dots, n$.

- Putting $X_n^{(n)} = \left(\frac{x_1^{(n)}}{x_n^{(n)}}, \frac{x_2^{(n)}}{x_n^{(n)}}, \dots, \frac{x_n^{(n)}}{x_n^{(n)}} \right)$ then $F(X_n^{(n)}, x) \leq F(X_n, x)$ for all $x \in$

$[0, 1]$ and $n = 1, 2, \dots$

• Selecting a sequence of indices n_k such that $F(X_{n_k}, x) \rightarrow g(x)$ and $F(X_{n_k}^{(n_k)}, x) \rightarrow \tilde{g}(x)$, then we have $\tilde{g}(x) \leq g(x)$ for all $x \in [0, 1]$.

Open problem is to execute Algorithm on a some number-theoretic sequence.

Examples:

(I) O. Strauch and J.T. Tóth (2001): Put $x_n = p_n$, the n th prime and denote

$$X_n = \left(\frac{2}{p_n}, \frac{3}{p_n}, \dots, \frac{p_{n-1}}{p_n}, \frac{p_n}{p_n} \right).$$

The sequence of blocks X_n is u.d. and therefore the ratio sequence p_m/p_n , $m = 1, 2, \dots, n$, $n = 1, 2, \dots$ is u.d. in $[0, 1]$. This generalizes a result of A. Schinzel (cf. W. Sierpiński (1964, p. 155)). Note that from u.d. of X_n applying the L^2 discrepancy of X_n we get the following interesting limit

$$\lim_{n \rightarrow \infty} \frac{1}{n^2 p_n} \sum_{i,j=1}^n |p_i - p_j| = \frac{1}{3}.$$

(II) O. Strauch and J.T. Tóth (2001): Let γ , δ , and a be given real numbers satisfying $1 \leq \gamma < \delta \leq a$. Let x_n be an increasing sequence of all integer points lying in the intervals

$$(\gamma, \delta), (\gamma a, \delta a), \dots, (\gamma a^k, \delta a^k), \dots$$

Then $G(X_n) = \{g_t(x); t \in [0, 1]\}$, where $g_t(x)$ has constant values

$$g_t(x) = \frac{1}{a^i(1+t(a-1))} \text{ for } x \in \frac{(\delta, a\gamma)}{a^{i+1}(t\delta + (1-t)\gamma)}, \quad i = 0, 1, 2, \dots$$

and on the component intervals it has a constant derivative

$$g'_t(x) = \frac{t\delta + (1-t)\gamma}{(\delta - \gamma)(\frac{1}{a-1} + t)} \text{ for } x \in \frac{(\gamma, \delta)}{a^{i+1}(t\delta + (1-t)\gamma)}, \quad i = 0, 1, 2, \dots$$

and $x \in \left(\frac{\gamma}{t\delta + (1-t)\gamma}, 1 \right)$.

Here we write $(xz, yz) = (x, y)z$ and $(x/z, y/z) = (x, y)/z$. From it follows that the set $G(X_n)$ has the following properties:

- (i) Every $g \in G(X_n)$ is continuous.
- (ii) Every $g \in G(X_n)$ has infinitely many intervals with constant values, i.e., with $g'(x) = 0$, and in the infinitely many complement intervals it has a constant derivative $g'(x) = c$, where $\frac{1}{a} \leq c \leq \frac{1}{\underline{a}}$ and for lower \underline{a} and upper \bar{a} asymptotic density of x_n we have $\underline{a} = \frac{(\delta - \gamma)}{\gamma(a-1)}$, $\bar{a} = \frac{(\delta - \gamma)a}{\delta(a-1)}$.
- (iii) The graph of every $g \in G(X_n)$ lies in the intervals $[\frac{1}{a}, 1] \times [\frac{1}{a}, 1] \cup [\frac{1}{a^2}, \frac{1}{a}] \times [\frac{1}{a^2}, \frac{1}{a}] \cup \dots$. Moreover, the graph g in $[\frac{1}{a^k}, \frac{1}{a^{k-1}}] \times [\frac{1}{a^k}, \frac{1}{a^{k-1}}]$ is similar to the graph of g in $[\frac{1}{a^{k+1}}, \frac{1}{a^k}] \times [\frac{1}{a^{k+1}}, \frac{1}{a^k}]$ with coefficient $\frac{1}{a}$. Using the parametric expression, it can be written for all $x \in (\frac{1}{a^{i+1}}, \frac{1}{a^i})$ that $g_t(x) = \frac{g_t(a^i x)}{a^i}$, $i = 0, 1, 2, \dots$

- (iv) $G(X_n)$ is connected and the upper distribution function $\bar{g}(x) = g_0(x) \in G(X_n)$ and the lower distribution function $\underline{g}(x) \notin G(X_n)$. The graph of $\underline{g}(x)$ on $[\frac{1}{a}, 1] \times [\frac{1}{a}, 1]$ coincides with the graph of $y(x) = \left(1 + \frac{1}{d} \left(\frac{1}{x} - 1\right)\right)^{-1}$ on $[\frac{\gamma}{\delta}, 1]$, further, on $[\frac{1}{a}, \frac{\gamma}{\delta}]$ we have $\underline{g}(x) = \frac{1}{a}$.
- (v) $G(X_n) = \left\{ \frac{g_0(x\beta)}{g_0(\beta)}; \beta \in \left[\frac{1}{a}, \frac{\delta}{a\gamma}\right] \right\}$.

(III) O. Strauch and G. Grekos (2007): Let x_n and y_n , $n = 1, 2, \dots$, be two strictly increasing sequences of positive integers such that for the related block sequences $X_n = \left(\frac{x_1}{x_n}, \dots, \frac{x_n}{x_n}\right)$ and $Y_n = \left(\frac{y_1}{y_n}, \dots, \frac{y_n}{y_n}\right)$, we have singleton $G(X_n) = \{g_1(x)\}$ and $G(Y_n) = \{g_2(x)\}$. Furthermore, let n_k , $k = 1, 2, \dots$, be an increasing sequence of positive integers such that $N_k = \sum_{i=1}^k n_i$ satisfies $\frac{n_k}{N_k} \rightarrow 1$. Denote by z_n the following increasing sequence of positive integers composed by blocks (here we use the notation $a(b, c, d, \dots) = (ab, ac, ad, \dots)$)

$$(x_1, \dots, x_{n_1}), x_{n_1}(y_1, \dots, y_{n_2}), x_{n_1}y_{n_2}(x_1, \dots, x_{n_3}), x_{n_1}y_{n_2}x_{n_3}(y_1, \dots, y_{n_4}), \dots$$

Then the sequence of blocks $Z_n = \left(\frac{z_1}{z_n}, \dots, \frac{z_n}{z_n}\right)$ has the set of d.f.s

$$G(Z_n) = \{g_1(x), g_2(x), c_0(x)\} \cup \{g_1(xy_n); n = 1, 2, \dots\} \cup \{g_2(xx_n); n = 1, 2, \dots\} \\ \cup \left\{ \frac{1}{1+\alpha}c_0(x) + \frac{\alpha}{1+\alpha}g_1(x); \alpha \in [0, \infty) \right\} \cup \left\{ \frac{1}{1+\alpha}c_0(x) + \frac{\alpha}{1+\alpha}g_2(x); \alpha \in [0, \infty) \right\},$$

where $g_1(xy_n) = 1$ if $xy_n \geq 1$, similarly for $g_2(xx_n)$.

Open problems:

1. Characterize a nonempty set H of d.f.s for which there exists an increasing sequence of positive integers x_n such that $G(X_n) = H$.

2. Probably $\frac{x_n}{x_{n+1}} \rightarrow 1$ implies that $G(X_n)$ is singleton.

Solution of 2. By F. Filip, L. Mišík and J.T. Tóth (2007) the solution is negative. They found counterexample:

Let a_k , n_k , $k = 1, 2, \dots$, and x_n , $n = 1, 2, \dots$ be three increasing integer sequences and $h_1 < h_2$ be two positive integers. Assume that

(i) $\frac{n_k}{n_{k+1}} \rightarrow 0$ for $k \rightarrow \infty$;

(ii) $\frac{a_k}{n_{k+1}} \rightarrow 0$ for $k \rightarrow \infty$;

(iii) for odd k we have

$$a_k^{h_2} \leq x_{n_k} = (a_{k-1} + n_k - n_{k-1})^{h_1} \leq (a_k + 1)^{h_2} \text{ and}$$

$$x_i = (a_k + i - n_k)^{h_2} \text{ for } n_k < i \leq n_{k+1};$$

(iv) for even k we have

$$a_k^{h_1} \leq x_{n_k} = (a_{k-1} + n_k - n_{k-1})^{h_2} \leq (a_k + 1)^{h_1} \text{ and}$$

$x_i = (a_k + i - n_k)^{h_1}$ for $n_k < i \leq n_{k+1}$.

Then $\frac{x_n}{x_{n+1}} \rightarrow 1$ and the set $G(X_n)$ of all distribution functions of the sequence of blocks X_n is $G(X_n) = G_1 \cup G_2 \cup G_3 \cup G_4$, where

$$G_1 = \{x^{\frac{1}{h_2}} \cdot t; t \in [0, 1]\},$$

$$G_2 = \{x^{\frac{1}{h_2}}(1-t) + t; t \in [0, 1]\},$$

$$G_3 = \{\max(0, x^{\frac{1}{h_1}} - (1 - x^{\frac{1}{h_1}})u); u \in [0, \infty)\} \text{ and}$$

$$G_4 = \{\min(1, x^{\frac{1}{h_1}} \cdot v); v \in [1, \infty)\}.$$

F. Filip, L. Mišík and J.T. Tóth (2007) also proved: If $G(X_n) = \{g(x)\}$ such that $g(x) < 1$ for $x \in [0, 1)$, then $\frac{x_n}{x_{n+1}} \rightarrow 1$. This implies that for u.d. sequence X_n we have $\frac{x_n}{x_{n+1}} \rightarrow 1$.

3. Characterize increasing sequences x_n , $n = 1, 2, \dots$, of positive integers for which $G(X_n)$ is connected.

NOTES: Some criterion of connectivity of $G(X_n)$ is given in G. Grekos and O. Strauch (2007, Th. 2) which is based on the relation $\tilde{g}(x) \prec g(x)$ defined on $G(X_n)$ if there exist α, β such that $\tilde{g}(x) = \alpha g(x\beta)$.

4. Prove or disprove: $G(X_n) \subset \{c_\alpha(x); \alpha \in [0, 1]\} \implies G(X_n) = \{c_0(x)\}$, for every increasing sequence x_n , $n = 1, 2, \dots$ of positive integers.

NOTES: G. Grekos and O. Strauch (2007) proved that if $G(X_n) \subset \{c_\alpha(x); \alpha \in [0, 1]\}$, then $c_0(x) \in G(X_n)$ and if $G(X_n)$ contains two different d.f.s, then also $c_1(x) \in G(X_n)$. Furthermore, we have that $\underline{d}(x_n) = 0$ and $\bar{d}(x_n) > 0$ implies $c_1(x) \in G(X_n)$.

5. Prove or disprove: $\lim_{n \rightarrow \infty} \frac{x_n}{x_1 + \dots + x_n} = 0 \iff c_0(x) \notin G(X_n)$. If it is true, then $c_0(x) \notin G(X_n)$ gives necessary and sufficient conditions that the sequence Y_n , $n = 1, 2, \dots$ of blocks

$$Y_n = \left(\frac{1}{x_n}, \frac{2}{x_n}, \dots, \frac{x_n}{x_n} \right)$$

is u.d. For a theory of blocks sequences Y_n , see Š. Porubský, T. Šalát and O. Strauch (1988).

6. There is open the theory of d.f. $G(X_n, Y_n)$ for two-dimensional blocks

$$(X_n, Y_n) = \left(\left(\frac{x_1}{x_n}, \frac{y_1}{y_n} \right), \left(\frac{x_2}{x_n}, \frac{y_2}{y_n} \right), \dots, \left(\frac{x_n}{x_n}, \frac{y_n}{y_n} \right) \right),$$

where x_n , $n = 1, 2, \dots$, and y_n , $n = 1, 2, \dots$ are increasing sequences of positive integers. It can be proved that the sequence $\left(\frac{p_i}{p_n}, \frac{i}{n} \right)$, $i = 1, 2, \dots, n$

is not u.d. in $[0, 1]^2$. Here p_n , $n = 1, 2, \dots$, is the increasing sequence of all primes.

7. L. Mišík: For every increasing sequence x_n , $n = 1, 2, \dots$, of positive integers there exists $g(x) \in G(X_n)$ such that $g(x) \geq x$ for all $x \in [0, 1]$. For $\underline{d}(x_n) > 0$ it holds by (vi).

Solution of 7. V. Baláž, L. Mišík, J.T. Tóth and O. Strauch (2013): If $\underline{d}(x_n) = 0$ and we select n_k such that $\frac{n_k}{x_{n_k}} = \min_{i \leq n_k} \frac{i}{x_i}$ and that $F(X_{n_k}, x) \rightarrow g(x)$, then $g(x) \geq x$ for $x \in [0, 1]$.

Proposed by O. Strauch.

V. BALÁŽ – L. MIŠÍK – J.T. TÓTH – O. STRAUCH: *Distribution functions of ratio sequences, III*, Publ. Math. Debrecen **82** (2013), no. 3-4, 511–529 (DOI:10.5486/PMD.2013.4770).

G. GREKOS – O. STRAUCH: *Distribution functions of ratio sequences, II*, Uniform Distribution Theory **2** (2007), no. 1, 53–77.

F. FILIP – J.T. TÓTH: *Distribution functions of ratio sequences*, (2006), (prepared).

F. FILIP – L. MIŠÍK – J.T. TÓTH: *On distribution functions of certain block sequences*, Uniform Distribution Theory **2** (2007), no. 2, 115–126.

Š. PORUBSKÝ – T. ŠALÁT – O. STRAUCH: *Transformations that preserve uniform distribution*, Acta Arith. **49** (1988), 459–479.

W. SIERPIŃSKI: *Elementary Theory of Numbers*, PAN Monografie Matematyczne, Vol. 42, PWN, Warszawa, 1964.

O. STRAUCH – J.T. TÓTH: *Distribution functions of ratio sequences*, Publ. Math. Debrecen **58** (2001), 751–778 (MR 2002h:11068).

O. STRAUCH – J.T. TÓTH: *Corrigendum to Theorem 5 of the paper "Asymptotic density of $A \subset \mathbb{N}$ and density of ratio set $R(A)$ " (Acta Arith. 87 (1998), 67–78)*, Acta Arith. **103.2** (2002), 191–200 (MR 2003f:11015).

1.10 Logarithmic and trigonometric functions.

[SP, p.2–131 and 2–132]: Find the set $G(x_n)$ for the following sequences x_n , $n = 1, 2, \dots$:

(i) $x_n = (\log n) \cos(n\alpha) \pmod{1}$

(ii) $x_n = (\cos n)^n$

(iii) $x_n = \cos(n + \log n) \pmod{1}$

NOTES:

(I) D. Berend, M.D. Boshernitzan and G. Kolesnik (1995) proved that (i) is everywhere dense in $[0, 1]$. They showed that there are uncountably many α 's for which every of these sequences (i) is not u.d.

(II) The original problem of everywhere density in $[-1, 1]$ of (ii) was posed by M. Benze and F. Popovici (1996) and was solved by J. Bukor (1997). This problem was also solved (in some generality) in F. Luca (1999). S. Hartman (1949) proved that if $\frac{\alpha}{\pi}$ is irrational, then

$$\liminf_{n \rightarrow \infty} (\cos \alpha n)^n = \liminf_{n \rightarrow \infty} (\sin \alpha n)^n = -1.$$

(III) The sequence (iii) is not u.d., which was proved by L. Kuipers (1953).

Solution of (iii). S. Steinerberger: The sequence $x_n = \cos(n + \log n) \bmod 1$ has the same a.d.f $g(x)$ as the sequence $\cos n \bmod 1$. It follows from that

(a) $\cos(n + \log n) = \cos 2\pi\left(\frac{n}{2\pi} + \frac{1}{2\pi} \log n\right) = \cos 2\pi z_n$, where

(b) $z_n = \frac{n}{2\pi} + \frac{1}{2\pi} \log n \bmod 1$ is u.d. sequence since $\frac{n}{2\pi}$ and $\frac{n}{2\pi} + \frac{1}{2\pi} \log n$ are u.d. simultaneously, see [SP, p. 2–27, 2.3.6.]

(c) Put $f(x) = \cos 2\pi x \bmod 1$ Then a.d.f. $g(x)$ of x_n is

$$g(x) = |f^{-1}([0, x])| = \frac{1}{2} - \frac{1}{\pi} \arccos x + 1 - \frac{1}{\pi} \arccos(x - 1).$$

(IV) D. Berend and G. Kolesnik (2011): The sequence

$$P(n) \cos n\alpha \bmod 1, n = 1, 2, \dots,$$

is completely u.d. for any non-constant polynomial $P(x)$ and α with $\cos \alpha$ transcendental. If $\cos \alpha$ is not transcendental Berend and Kolesnik (2011) also proved: Let α be such that $e^{i\alpha}$ is an algebraic number of degree d which is not a root of unity. Then the sequence

$$(1) (P(n) \cos n\alpha, P(n+1) \cos(n+1)\alpha, \dots, P(n+d-1) \cos(n+d-1)\alpha) \bmod 1,$$

$n = 1, 2, \dots$, is u.d. for any non-constant polynomial $P(x)$.

Open problem. Berend and Kolesnik (2011): Let $P(x) = x$, $\alpha = \arccos 3/5$, i.e. $e^{i\alpha} = (3 + 4i)/5$ and denote $x_n = P(n) \cos n\alpha = n \frac{(3+4i)^n - (3-4i)^n}{2 \cdot 5^n}$. Then by (1) the sequence $(x_n, x_{n+1}) \bmod 1$ is u.d., but $(x_n, x_{n+1}, x_{n+2}, x_{n+3}, x_{n+4}) \bmod 1$ is not u.d. The authors ask whether the sequences $(x_n, x_{n+1}, x_{n+2}) \bmod 1$ and $(x_n, x_{n+1}, x_{n+2}, x_{n+3}) \bmod 1$ are u.d.

Solution of (ii). Ch. Aistleitner, M. Hofer and M. Madritsch (2013): Let $x_n = \cos(\alpha n)^n \bmod 1$, $n = 1, 2, \dots$. For $\frac{\alpha}{2\pi} \notin \mathbb{Q}$ we set $a = 3/4$, in the case $\frac{\alpha}{2\pi} = \frac{p}{q} \in \mathbb{Q}$ for p, q co-prime let

$$a = \begin{cases} \frac{q+1}{2q} + \frac{q-1}{4q}, & \text{if } 4 \mid (q-1), \\ \frac{q-1}{2q} + \frac{q+1}{4q}, & \text{if } 4 \nmid (q-1) \end{cases}$$

for q odd and let

$$a = \begin{cases} \frac{1}{2} + \frac{q-2}{4q}, & \text{if } 4 \nmid q \text{ and } 8 \mid (q-2), \\ \frac{1}{2} + \frac{q+2}{4q}, & \text{if } 4 \nmid q \text{ and } 8 \nmid (q-2), \\ \frac{q+2}{2q} + \frac{1}{4}, & \text{if } 4 \mid q \text{ and } 8 \nmid q, \\ \frac{q+2}{2q} + \frac{q-4}{4q}, & \text{if } 8 \mid q \end{cases}$$

for q even. Then a.d.f. of x_n is given by

$$g_a(x) = \begin{cases} 0, & \text{if } x = 0, \\ a, & \text{if } 0 < x < 1, \\ 1, & \text{if } x = 1. \end{cases}$$

Partial solution of (i). 1⁰. Ch. Aistleitner, M. Hofer and M. Madritsch (2013) proved: Let α be such that the discrepancy D_N of the sequence

$$\frac{\alpha}{2\pi}n \bmod 1, \quad n = 1, 2, \dots, N$$

is of asymptotic order $D_N = o\left(\frac{1}{\log N}\right)$. Then the sequence $(\log n) \cos(n\alpha) \bmod 1$ is u.d. in $[0, 1]$.

2⁰. Let $x_n = (\log n) \cos(n\alpha) \bmod 1$, $n = 1, 2, \dots$, $\frac{\alpha}{2\pi} = \frac{p}{q}$, where p, q are co-prime and let $N_1 < N_2 < \dots$ be fixed integer sequence such that

$$\lim_{k \rightarrow \infty} \{\cos(\alpha i) \log N_k\} = \beta_i, \quad \text{for } i = 1, \dots, q. \quad (1)$$

Then there exists d.f. $g(x) = \lim_{k \rightarrow \infty} F_{N_k}(x)$, $F_N(x) = \frac{\{n \leq N; x_n \in [0, x]\}}{N}$, such that

$$g(x) = \frac{1}{q} \sum_{i=1}^q h_{q, \beta_i, c_i}(x), \quad (2)$$

where $h_{q, \beta_i, c_i}(x) =$

$$\begin{cases} f_{\beta_i, c_i}(x + 1 - \nu_i) - f_{\beta_i, c_i}(1 - \nu_i), & \text{if } 0 \leq x \leq \nu_i \text{ and } c_i > 0, \\ f_{\beta_i, c_i}(x - \nu_i) + 1 - f_{\beta_i, c_i}(1 - \nu_i), & \text{if } \nu_i \leq x \leq 1 \text{ and } c_i > 0, \\ f_{\beta_i, c_i}(x + \nu_i) - f_{\beta_i, c_i}(\nu_i), & \text{if } 0 \leq x \leq 1 - \nu_i \text{ and } c_i < 0, \\ f_{\beta_i, c_i}(x - (1 - \nu_i)) + 1 - f_{\beta_i, c_i}(\nu_i), & \text{if } 1 - \nu_i \leq x \leq 1 \text{ and } c_i < 0, \\ \mathbf{1}_{\{(0,1]\}}(x) & \text{if } c_i = 0, \end{cases}$$

where $\nu_i = \{|c_i| \log(q)\}$, $c_i = \cos(\alpha i)$ and

$$f_{\beta, c}(x) = \begin{cases} g_{\beta, c}(x), & \text{if } c > 0, \\ 1 - g_{\beta, |c|}(1 - x) & \text{if } c < 0, \\ \mathbf{1}_{\{(0,1]\}}(x) & \text{if } c = 0, \end{cases}$$

and

$$g_{\beta, c}(x) = \frac{e^{\frac{\min(x, \beta)}{c}} - 1}{e^{\frac{\beta}{c}} - 1} + \frac{1}{e^{\frac{\beta}{c}}} \frac{e^{\frac{x}{c}} - 1}{e^{\frac{1}{c}} - 1}.$$

Moreover, the set $G(x_n)$ is the set of all d.f. of the form (2) for those $(\beta_1, \dots, \beta_q)$ for which a subsequence $(N_k)_{k \geq 1}$ satisfying (1) exists.

The authors note that for arbitrary q , it is a difficult problem to determine all possible vectors $(\beta_1, \dots, \beta_q)$ for which there exists $N_1 < N_2 < \dots$ such that (1) holds, referred to K. Gristmair (1997).

Proposed by O. Strauch.

CH. AISTLEITNER – M. HOFER – M. MADRITSCH: *On the distribution functions of two oscillating sequences*, Uniform Distribution Theory **8** (2013), no. 2, 157–169.

M. BENCZE – F. POPOVICI: *OQ. 45*, Octogon Math. Mag.(Brasov) **4** (1996), 77.

- D. BEREND – G. KOLESNIK: *Complete uniform distribution of some oscillating sequences*, J. Ramanujan Math. Soc. **26** (2011), no. 2, 127–144.
- D. BEREND – M.D. BOSHERNITZAN – G. KOLESNIK: *Distribution modulo 1 of some oscillating sequences. II*, Israel J. Math. **92** (1995), no. 1–3, 125–147 (MR 96j:11105).
- J. BUKOR: *On a certain density problem*, Octogon Mathematical Magazine (Brasov) **5** (1997), no. 2, 73–75.
- K. GIRSTMAIR: *Some linear relations between values of trigonometric functions at $k\pi/n$* , Acta Arith. **81** (1997), no. 4, 387–398.
- S. HARTMAN: *Sur une condition supplémentaire dans les approximations diophantiques*, Colloq. Math. **2** (1949), no. 1, 48–51 (MR **12**, 807a).
- F. LUCA: $\{(\cos(n))^n\}_{n \geq 1}$ is dense in $[-1, 1]$, Bull. Math. Soc. Sci. Math. Roumanie (N.S.) **42(90)** (1999), no. 4, 369–376 (MR1879621(MR 2002k:11118)).
- L. KUIPERS: *Continuous and discrete distribution modulo 1*, Indag. Math. **15** (1953), 340–348 (MR **15**, 410e).

1.11 Euler totient function.

(cf. [SP, p. 2–191, 2.20.11]). If φ is the Euler totient function then the sequence

$$\frac{\varphi(n)}{n}, \quad n = 1, 2, 3, \dots,$$

has in $[0, 1]$ singular a.d.f.

$$g_0(x).$$

NOTES: (I) I.J. Schoenberg (1928, 1936) proved that this sequence has continuous and strictly increasing a.d.f.

(II) P. Erdős (1939) showed that this a.d.f. is singular. Here a function is **singular**, if it is continuous, strictly monotone and has vanishing derivative almost everywhere on the interval of its definition.

(III) H. Davenport (1933) expressed

$$g_0(x) = \sum_{n=1}^{\infty} A_n(x),$$

where

$$A_n(x) = \frac{1}{a_n(x)} - \sum_{i < n} \frac{1}{[a_i(x), a_n(x)]} + \sum_{i < j < n} \frac{1}{[a_i(x), a_j(x), a_n(x)]} - \dots$$

and $[a, b]$ is the least common multiple of a and b . Here $a_1(x) < a_2(x) < \dots$ is the sequence of the all positive integers n for which $\frac{\varphi(n)}{n} \leq x$ and for every divisor $d|n$, $d \neq n$ we have $\frac{\varphi(d)}{d} > x$. These n are called x -numbers. Directly from definition we have

(i) Every x -number is square-free.

(ii) Every square-free a is an x -number for some x . Concretely, if $a = p_1 p_2 \dots p_m$,

$$p_1 < p_2 < \dots < p_m, \quad p_i \text{ are primes, then } a \text{ is } x\text{-number for every } x \in \left[\prod_{i=1}^m \left(1 - \frac{1}{p_i}\right), \prod_{i=1}^{m-1} \left(1 - \frac{1}{p_i}\right) \right)$$

- (iii) For every $i < j$ we have $a_i(x) \nmid a_j(x)$.
- (iv) Let $p_1 < p_2 < \dots$ be an increasing sequence of all primes and let $x \in \left[1 - \frac{1}{p_s}, 1\right)$. Then $a_1(x) = p_1 = 2, a_2(x) = p_2 = 3, \dots, a_s(x) = p_s$. If furthermore $x < 1 - \frac{1}{p_{s+1}}$, then for every $j > s$, the $a_j(x)$ cannot be a prime and $p_i \nmid a_j(x), i = 1, 2, \dots, s$.
- (v) If $x \in \left[\prod_{i=1}^s \left(1 - \frac{1}{p_i}\right), \prod_{i=1}^{s-1} \left(1 - \frac{1}{p_i}\right)\right)$, then $a_1(x) = \prod_{i=1}^s p_i$ (p_i as in (iv)).
- (vi) For every $n = 1, 2, \dots$ and every $x \in (0, 1)$ we have

$$\frac{\varphi(n)}{n} \leq x \iff \exists_{i=1,2,\dots} a_i(x) | n.$$

- (vii) Assume that $x < x'$. Then for every x -number $a_i(x)$ there exists x' -number $a_j(x')$ such that $a_j(x') | a_i(x)$.

(III') Applying Davenport's method B.A. Venkov (1949) proved that

(i) $(1 - g_0(x)) \log \frac{1}{1-x} \rightarrow e^{-c}$ as $x \rightarrow 1$, where c is Euler's constant.

(ii) $x \log \log \frac{1}{g_0(x)} \rightarrow e^{-c}$ as $x \rightarrow 0$.

(iii) Let p be a prime. Then for every $1 - \frac{1}{p} \leq x$ we have

$$\frac{1}{p} = g_0(x) - (p-1)g_0\left(x\left(1 - \frac{1}{p}\right)\right) + (p-1)^2 g_0\left(x\left(1 - \frac{1}{p}\right)^2\right) - \dots$$

(iv) The function $g_0(x)$ at every $x = \frac{\varphi(n)}{n}, n = 1, 2, \dots$, has infinite left derivative.

(v) $\left(\int_0^1 x^s dg_0(x)\right) \log s \rightarrow e^{-c}$ as $s \rightarrow \infty$ (s are positive integers).

(IV) A.S. Faïnleïb (1967) proved that

$$\frac{A([0, x]; N; \varphi(n)/n)}{N} = g_0(x) + \mathcal{O}\left(\frac{1}{\log \log N}\right).$$

(V) W. Schwarz (1962)(c.f. A.G. Postnikov (1971, p. 267)) proved: Let $f(x)$ be a polynomial with integer coefficients having non-zero discriminant. Assume that g.c.d of coefficients of $f(x)$ is 1 and $f(n) > 0$ for $n = 1, 2, \dots$. Let $L(d)$ denote the number of solutions $f(n) \equiv 0 \pmod{d}$. Then

$$\frac{1}{N} \sum_{n=1}^N \frac{\varphi(f(n))}{f(n)} = \prod_{\substack{p=2 \\ p\text{-prime}}}^{\infty} \left(1 - \frac{L(p)}{p^2}\right) + \mathcal{O}(\log^c N),$$

where $c > 0$ is a constant. This leads to

Open problem 1: Find a.d.f (if exists) of the sequence

$$\frac{\varphi(f(n))}{f(n)}, \quad n = 1, 2, \dots$$

(VI) O. Strauch (1996) proved that

$$\int_0^1 g_0^2(x) dx = 1 - \frac{6}{\pi^2} - \frac{1}{2} \lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{m,n=1}^N \left| \frac{\varphi(m)}{m} - \frac{\varphi(n)}{n} \right|$$

and he gave an estimate

$$\frac{2}{\pi^4} \leq \lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{m,n=1}^N \left| \frac{\varphi(m)}{m} - \frac{\varphi(n)}{n} \right| \leq 2 \frac{6}{\pi^2} \left(1 - \frac{6}{\pi^2} \right). \quad (1)$$

Open problem 2: Find an estimation of the limit

$$\lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{m,n=1}^N \left| \frac{\varphi(m)}{m} - \frac{\varphi(n)}{n} \right| = L$$

better as (1), where $L \in [0.021, 0.392]$.

• J.-Ch. Schlage-Puchta (2009) send a method which gives $L \in [0.27425, 0.274465]$.
(VI) The aim of this problem is to find $\int_0^1 g_0^2(x) dx$. It is motivated by the paper of O. Strauch (1994) about three dimensional body Ω of points of the form

$$\left(\int_0^1 g(x) dx, \int_0^1 xg(x) dx, \int_0^1 g^2(x) dx \right),$$

where $g(x)$ runs the set of all d.f.'s. The points achieved for singulars $g(x)$ are interior points of Ω . Using an expression of the boundary of Ω in Problem 1.23.2 we can find

$$0.250 < \int_0^1 g_0^2(x) dx < 0.307 .$$

(VII) F. Luca ([a]2003) proved that, if $M_n = 2^n - 1$ is the n th Mersenne number then the subsequence $\varphi(M_n)/M_n$ is dense in $[0, 1]$ and has an a.d.f. ([b]2005).

(VII') F. Luca and I.E. Shparlinski (2007) proved the existence of the moment

$$\frac{1}{N} \sum_{n=0}^{N-1} \left(\frac{\varphi(F_n)}{F_n} \right)^k = \Gamma_k + O_k \left(\frac{(\log N)^k}{N} \right)$$

for all $k = 1, 2, \dots$ with some positive constant Γ_k . Thus the sequence

$$\frac{\varphi(F_n)}{F_n}, \quad n = 0, 1, 2, \dots$$

has an a.d.f. Luca in ([a]2003) also proved that $\varphi(F_n)/F_n$ is dense in $[0, 1]$.

(VIII) (See [SP, p. 1–13]). I.J. Schoenberg (1959) introduced the following summation method: the sequence x_n is called φ -**convergent** to α if the sequence $y_n = \frac{1}{n} \sum_{d|n} \varphi(d)x_d$ converges to α . Schoenberg's Theorem 2 (1959) shows that the φ -convergence of x_n implies the classical convergence of x_{n_k} (to the same limit) for every sequence n_k for which $\liminf_{k \rightarrow \infty} \frac{\varphi(n_k)}{n_k} > 0$. Since a 0–1 φ -convergent sequence has the φ -limit 0 or 1, no φ -u.d. sequence exists (E. Kováč (2005)).

Open problem 3: Find a sequence x_n for which $y_n = \frac{1}{n} \sum_{d|n} \varphi(d)x_d \pmod{1}$ is u.d. in $[0, 1]$.

In connection of this we mentioned (cf. A.G. Postnikov (1971, p. 219, Th. 6b), [SP, p. 2–189, 2.20.8]): Let $f(n)$ be an arithmetical function which satisfies

$$(i) \quad f(n) = \sum_{d|n} \Phi(d),$$

$$(ii) \quad \sum_{n=1}^{\infty} \frac{|\Phi(d)|}{d} < \infty$$

for some arithmetical function Φ . Then the sequence

$$f(n), \quad n = 1, 2, \dots,$$

has the a.d.f.

$$g(x)$$

defined on $(-\infty, \infty)$.

• For $x_n = \frac{\varphi(n)}{n}$ and an interval $(k, k + N]$ define the step d.f.

$$F_{(k, k+N]}(x) = \frac{\#\{n \in (k, k + N]; x_n \in [0, x]\}}{N}.$$

Open problem 4: Find all possible limits of step d.f.s $F_{(k, k+N]}(x)$, for sequences of intervals $(k, k + N]$.

(IX) P. Erdős (1946) proved:

(i) If $\frac{\log \log \log k}{N} \rightarrow 0$ as $N \rightarrow \infty$, then $F_{(k, k+N]}(x) \rightarrow g_0(x)$ for $x \in [0, 1]$ and by Chinese remainder theorem he found k and N such that $\frac{\log \log \log k}{N} \rightarrow \frac{1}{2}$ and $\frac{1}{N} \sum_{k < n \leq k+N} \frac{\varphi(n)}{n} < \frac{1}{2} < \frac{1}{N} \sum_{n=1}^N \frac{\varphi(n)}{n} = \frac{6}{\pi^2} + O\left(\frac{\log N}{N}\right)$, thus $F_{(k, k+N]}(x) \not\rightarrow g_0(x)$.

(ii) For a proof of (i) he used

$$\left(\frac{1}{N} \sum_{k < n \leq k+N} \left(\frac{\varphi(n(t))}{n(t)} \right)^s - \frac{1}{N} \sum_{n=1}^N \left(\frac{\varphi(n)}{n} \right)^s \right) \rightarrow 0,$$

where $n(t) = \prod_{p|n, p \leq t} p$, p are primes and $t = N$.

(X) V. Baláž, P. Liardet and O. Strauch (2007) proved:

(i) Necessary and sufficient condition: For any two sequences N and k of positive sequences, $N \rightarrow \infty$, we have $F_{(k, k+N]}(x) \rightarrow g_0(x)$, for every $x \in [0, 1]$, if and only if, for every $s = 1, 2, \dots$, $\frac{1}{N} \sum_{k < n \leq k+N} \sum_{N < d|n} \Phi(d) \rightarrow 0$, were (cf. A.G. Postnikov (1971, p. 360)) $\Phi(d) = \prod_{p|d} \left(\left(1 - \frac{1}{p}\right)^s - 1 \right)$ for the squarefree d and $\Phi(d) = 0$ in others, where p denotes a prime. In quantitative form:

$$\begin{aligned} \frac{1}{N} \sum_{k < n \leq k+N} \sum_{N < d|n} \Phi(d) &= \frac{1}{N} \sum_{k < n \leq k+N} \left(\frac{\varphi(n)}{n} \right)^s - \frac{1}{N} \sum_{n=1}^N \left(\frac{\varphi(n)}{n} \right)^s \\ &\quad + O\left(\frac{3^s (1 + \log N)^s}{N} \right). \end{aligned}$$

Using this they found that for $k = \prod_{p \leq e e^e N} p$ we have $F_{(k, k+N]}(x) \rightarrow g_0(x)$ as

$N \rightarrow \infty$ and contrary to (IX)(i) we have $\frac{\log \log \log k}{N} \rightarrow \infty$.

(ii) A quantitative form of Erdős' (IX)(ii): For every integer k, N and $t = N$ we have

$$\frac{1}{N} \sum_{k < n \leq k+N} \left(\frac{\varphi(n(t))}{n(t)} \right)^s = \frac{1}{N} \sum_{n=1}^N \left(\frac{\varphi(n)}{n} \right)^s + O\left(\frac{3^s(1 + \log N)^s}{N} \right)$$

for $s = 1, 2, \dots$

(iii) This implies that every d.f. $g(x)$, $F_{(k, k+N]} \rightarrow g(x)$ on $(0, 1)$ must satisfies

$$\int_0^1 x^s dg(x) \leq \int_0^1 x^s dg_0(x),$$

for every $s = 1, 2, \dots$

(iv) By Chinese theorem it can be found a sequence of intervals $(k, k + N]$ such that $F_{(k, k+N]}(x) \rightarrow c_0(x)$, where d.f. $c_0(x)$ has a step 1 in $x = 0$.

(v) Assume that $F_{(k, k+N]}(x) \rightarrow g(x)$ for all $x \in (0, 1)$. Then

$$g_0(x) \leq g(x) \leq g_0(x) + \prod_{i=1}^s \left(1 - \frac{1}{p_i} \right) \quad (1)$$

for $x \in (0, 1)$, where p_1, p_2, \dots is the increasing sequence of all primes and $1 - \frac{1}{p_s} \leq x$. By an anonymous referee in all cases the right hand side of (1) is ≥ 1 .

(XI) A. Schinzel and Y. Wang (1958) proved that for any given $(\alpha_1, \alpha_2, \dots, \alpha_{N-1}) \in [0, \infty)^{N-1}$ we can select a sequence of k such that

$$\left(\frac{\varphi(k+2)}{\varphi(k+1)}, \frac{\varphi(k+3)}{\varphi(k+2)}, \dots, \frac{\varphi(k+N)}{\varphi(k+N-1)} \right) \rightarrow (\alpha_1, \alpha_2, \dots, \alpha_{N-1}).$$

Select a subsequence of k such that $\frac{\varphi(k+1)}{k+1} \rightarrow \alpha$. Then

$$\left(\frac{\varphi(k+1)}{k+1}, \frac{\varphi(k+2)}{k+2}, \dots, \frac{\varphi(k+N)}{k+N} \right) \rightarrow (\alpha, \alpha\alpha_1, \alpha\alpha_1\alpha_2, \dots, \alpha\alpha_1\alpha_2 \dots \alpha_{N-1}).$$

Now, for arbitrary d.f. $\tilde{g}(x)$ there exists a sequence α_n , $n = 1, 2, \dots$ in $(0, \infty)$ such that for every $n = 1, 2, \dots$ we have $\alpha_1\alpha_2 \dots \alpha_n \in (0, 1)$ and that the sequence $\alpha_1\alpha_2 \dots \alpha_n$, $n = 1, 2, \dots$, has asymptotic d.f. $\tilde{g}(x)$. Then there exists $\alpha \in (0, 1]$ and a sequence of intervals $(k, k + N]$ such that $F_{(k, k+N]} \rightarrow g(x)$ and for $x \in (0, 1)$ we have

$$g(x) = \begin{cases} \tilde{g}\left(\frac{x}{\alpha}\right) & \text{if } x \in [0, \alpha), \\ 1 & \text{if } x \in [\alpha, 1]. \end{cases}$$

Open problem 5: Find a distribution of the sequence

$$\left(\frac{\varphi(n)}{n}, \frac{\varphi(n+1)}{n+1} \right), \quad n = 1, 2, \dots$$

Problems 1–5 proposed by O. Strauch.

Open problem 6 proposed by F. Luca:

- (i) Is the sequence of general term $(\varphi(1) + \cdots + \varphi(n))/n$ uniformly distributed modulo 1?
- (ii) Is the sequence of general term $(\varphi(1)\varphi(2)\cdots\varphi(n))^{1/n}$ uniformly distributed modulo 1?

Regarding (i) above, R. Balasubramanian and F. Luca (2007) have shown that the set of n such that $(\varphi(1) + \cdots + \varphi(n))/n$ is an integer is of asymptotic density zero.

Solution of 6. J.-M. Deshouillers and H. H. Iwaniec (2008) gave positive answer to (i) and conditional positive answer to (ii), they proved:

(XII) Let $\nu(n)$ be an arithmetic function which is completely multiplicative and satisfies the conditions

(i) $|\nu(p)| \leq \nu$ for some positive number ν and every prime p ,

(ii) $\sum_{d \leq x} \mu(d)\nu(d) \ll x(\log x)^{-A}$ for every positive A ,

where the implied constant depends only on ν and A . Define the arithmetic function ϕ by $\phi(m) = m \prod_{p|m} \left(1 - \frac{\nu(p)}{p}\right)$. Then, if the number $\alpha =$

$\frac{1}{2} \prod_p \left(1 - \frac{\nu(p)}{p^2}\right)$ is irrational, the sequence $\frac{1}{n} \sum_{m \leq n} \phi(m)$, $n = 1, 2, \dots$, is u.d. modulo one.

NOTES: For classical Euler $\varphi(n)$ function corresponding $\alpha = \frac{3}{\pi^2}$.

(XIII) Let $\nu(n)$ be a completely multiplicative function such that

(i) $-\nu \leq \nu(p) < \min\{p, \nu\}$ for some positive ν and every prime p

(ii) that there exist real numbers β and λ such that $\prod_{p \leq n} \left(1 - \frac{\nu(p)}{p}\right) = \beta(\log n)^{-\lambda} \left(1 + O\left(\frac{1}{\log n}\right)\right)$, where the implied constant depends only on ν .

Again as in (I), we define the strongly multiplicative function ϕ by $\phi(m) = m \prod_{p|m} \left(1 - \frac{\nu(p)}{p}\right)$, and we let $\alpha = \frac{1}{e} \prod_p \left(1 - \frac{\nu(p)}{p}\right)^{\frac{1}{p}}$. If α is irrational, then

the sequence $\left(\prod_{m \leq n} \phi(m)\right)^{\frac{1}{n}}$, $n = 1, 2, \dots$, is u.d. modulo one.

(XIV) Let the arithmetical function ν satisfy (i) and (ii) in (XIII). If α is rational and ν takes only algebraic values, then the sequence $\left(\prod_{m \leq n} \phi(m)\right)^{\frac{1}{n}}$, $n = 1, 2, \dots$, is not uniformly distributed modulo one. By the authors comments, for the classical Euler $\varphi(n)$ the arithmetic property of corresponding $\alpha = \frac{1}{e} \prod_p \left(1 - \frac{1}{p}\right)^{\frac{1}{p}}$ is an **open problem**. This constant is very likely to be irrational: Richard Bumby showed that if α is rational, then its denominator has at least 20 decimal digits. A special case of (XIV) shows that if

the constant α is rational, then the sequence $\left(\prod_{m \leq n} \varphi(m)\right)^{\frac{1}{n}}$, $n = 1, 2, \dots$, is not u.d. modulo 1.

(XV) F. Luca, V.J. Mejía Huguet and F. Nicolae (2009) show that

$$\left(\frac{\varphi(F_{n+1})}{\varphi(F_n)}, \frac{\varphi(F_{n+2})}{\varphi(F_n)}, \dots, \frac{\varphi(F_{n+k})}{\varphi(F_n)}\right), \quad n = 1, 2, \dots$$

is dense in $[0, \infty)^k$, $k = 1, 2, \dots$. The authors have the following comments:

- for any positive integer k and every permutation (i_1, \dots, i_k) there exist infinitely many integers n such that $\varphi(F_{n+i_1}) < \varphi(F_{n+i_2}) < \dots < \varphi(F_{n+i_k})$.

- P. Erdős, K. Györy and Z. Papp (1980) call two arithmetic functions $f(n)$ and $g(n)$ *independent* if for every permutations (i_1, \dots, i_k) and (j_1, \dots, j_k) of $(1, \dots, k)$, there exist infinitely many integers n such that both

$$f(n + i_1) < f(n + i_2) < \dots < f(n + i_k),$$

$$g(n + j_1) < g(n + j_2) < \dots < g(n + j_k).$$

- $\varphi(n)$ and Carmichael $\lambda(n)$ are independent (N. Doyon and F. Luca (2006)).

- $\sigma(\varphi(n))$ and $\varphi(\sigma(n))$ are independent (M.O. HERNANE and F. Luca (2009)).

Open problems in F. Luca, V.J. Mejía Huguet and F. Nicolae (2009):

- Are the functions $\varphi(F_n)$ and $F_{\varphi(n)}$ independent?

- Are the functions $\varphi(F_n)$ and $\varphi(M_n)$ independent?

Entered by O. Strauch.

R. BALASUBRAMANIAN – F. LUCA: *On the sum of the first n values of the Euler function*, (2007), (Preprint).

V. BALÁŽ – P. LIARDET – O. STRAUCH: *Distribution functions of the sequence $\varphi(M)/M$, $M \in (K, K + N]$ as K, N go to infinity*, INTEGERS **10** (2010), 705–732.

H. DAVENPORT: *Über numeri abundantes*, Sitzungsber. Preuss. Acad., Phys.–Math. Kl. **27** (1933), 830–837 (Zbl 8, 197).

J.-M. DESHOUILERS – H. IWANIEC: *On the distribution modulo one of the mean values of some arithmetical functions*, Uniform Distribution Theory **3** (2008), no. 1, 111–124.

N DOYON – F. LUCA: *On the local behavior of the Carmichael λ -function*, Michigan Math. J. **54** (2006), 283–300.

P.D.T.A. ELLIOTT: *Probabilistic Number Theory I. Mean-value Theorems*, Grundlehren der mathematischen Wissenschaften, Vol. 239, Springer Verlag, New York, Heidelberg, Berlin, 1979 (MR 82h:10002a).

P. ERDŐS: *On the smoothness of the asymptotic distribution of additive arithmetical functions*, Amer. Journ. Math. **61** (1939), 722–725 (Zbl 22, 10).

P. ERDŐS: *Some remarks about additive and multiplicative functions*, Bull. Amer. Math. Soc. **52** (1946), 527–537 (MR 0016078).

P. ERDŐS – K. GYÖRY – Z. PAPP: *On some new properties of functions $\sigma(n)$, $\varphi(n)$, $d(n)$ and $\nu(n)$* , Mat. Lapok **28** (1980), 125–131.

A.S. FAÏNLEÏB: *Distribution of values of Euler's function* (Russian), Mat. Zametki **1** (1967), 645–652; English translation: Math. Notes **1** (1976), 428–432 (MR **35**#6636)

M.O. HERNANE – F. LUCA: *On the independence of ϕ and σ* , Acta Arith. **138** (2009), 337–346.

E. KOVÁČ: *On φ -convergence and φ -density*, Math. Slovaca **55** (2005), no. 3, 329–351.

- [a] F. LUCA: *On the sum divisors of the Mersenne numbers*, Math. Slovaca **53** (2003), no. 5, 457–466 (MR 2038513).
- [b] F. LUCA: *Some mean values related to average multiplicative orders of elements in finite fields*, Ramanujan J. **9** (2005), no. 1–2, 33–44 (MR 2166376).
- F. LUCA – I.E. SHPARLINSKI: *Arithmetic functions with linear recurrences*, J. Number Theory **125** (2007), 459–472.
- F. LUCA – V.J. MEJÍA HUGUET – F. NICOLAE: *On the Euler function of Fibonacci numbers*, Journal of Integer Sequences **12** (2009), A 09.6.6.
- A.G. POSTNIKOV: *Introduction to Analytic Number Theory* (Russian), Izd. Nauka, Moscow, 1971 (MR **55**#7859; for the English translation see MR 89a:11001).
- A. SCHINZEL – Y. WANG: *A note on some properties of the functions $\phi(n)$, $\sigma(n)$ and $\theta(n)$* , Bull. Acad. Polon. Sci. Cl. III **4** (1956), 207–209 (MR **18**, 17c).
- I.J. SCHOENBERG: *Über die asymptotische Verteilung reeller Zahlen mod 1*, Math. Z. **28** (1928), 171–199 (JFM 54.0212.02).
- I.J. SCHOENBERG: *On asymptotic distribution of arithmetical functions*, Trans. Amer. Math. Soc. **39** (1936), 315–330.
- I.J. SCHOENBERG: *The integrability of certain functions and related summability methods*, Amer. Math. Monthly **66** (1959), 361–375 (MR **21**#3696).
- W. SCHWARZ: *Über die Summe $\sum_{n \leq x} \varphi(f(n))$ und verwandte Probleme*, Monatsh. Math. **66** (1962), 43–54.
- O. STRAUCH: *Integral of the square of the asymptotic distribution function of $\phi(n)/n$* , Math. Institute, Slovak Acad. Sci., Bratislava, Slovak Republic, 1996, 7 pp.
- O. STRAUCH: *A new moment problem of distribution functions in the unit interval*, Math. Slovaca **44** (1994), no. 2, 171–211 (MR 95i:11082).
- M.M. TJAN: *Remainder terms in the problem of the distribution of values of two arithmetic functions*, (Russian), Dokl. Akad. Nauk SSSR **150** (1963), 998–1000 (MR **27**#4789).
- A.B. VENKOV: *On one monotone function*, (Russian), Učen. zap. Leningr. gos. un-ta. ser. matem. **16** (1949), 3–19.

1.12 van der Corput sequence in the base q

[SP, p. 2–102, 2.11]. Let $q \geq 2$ be an integer and

$$n = a_0(n) + a_1(n)q + \cdots + a_{k(n)}(n)q^{k(n)}, \quad a_j(n) \in \{0, 1, \dots, q-1\}, \quad a_{k(n)}(n) > 0,$$

be the q -adic digit expansion of integer n in the base q . Then the van der Corput sequence $\gamma_q(n)$, $n = 0, 1, 2, \dots$, in the base q defined by

$$\gamma_q(n) = \frac{a_0(n)}{q} + \frac{a_1(n)}{q^2} + \cdots + \frac{a_{k(n)}(n)}{q^{k(n)+1}}$$

is u.d.

Open problem: Find the distribution function of the sequence

$$(\gamma_q(n), \dots, \gamma_q(n+s-1)), \quad n = 0, 1, 2, \dots, \text{ in } [0, 1]^s.$$

NOTES:

(I) The $\gamma_q(n)$ is called the **radical inverse function** of the natural q -adic digit expansion of n .

(II) The Halton sequence in the bases q_1, \dots, q_s is defined by

$$\mathbf{x}_n = (\gamma_{q_1}(n), \dots, \gamma_{q_s}(n)), \quad n = 0, 1, 2, \dots$$

For the pairwise coprime bases q_1, \dots, q_s the Halton sequence is u.d., cf. [SP, p. 3–72].

(i) The Halton sequence \mathbf{x}_n is u.d. in $[0, 1]^s$ if and only if the bases q_1, \dots, q_s are coprime, see P. Hellekalek and H. Niederreiter (2011).

(III) Van der Corput sequence $\gamma_q(n)$, $n = 0, 1, \dots, N - 1$ has discrepancy

$$D_N^*(\gamma_q(n)) < \frac{1}{N} \left(\frac{q \log(qN)}{\log q} \right),$$

i.e., it is low discrepancy sequence, but for $s = 2$ we have, see O. Blažeková (2007),

$$D_N(\gamma_q(n), \gamma_q(n+1)) = \frac{1}{4} + O(D_N(\gamma_q(n))),$$

$$D_N^*((\gamma_q(n), \gamma_q(n+1))) = \max \left(\frac{1}{q} \left(1 - \frac{1}{q} \right), \frac{1}{4} \left(1 - \frac{1}{q} \right)^2 \right) + O(D_N(\gamma_q(n))).$$

Thus, van der Corput sequence is not pseudo-random.

(IV) Discrepancy bounds of the van der Corput and Halton sequences can be found in the added bibliography.

(V) **Solution for $s = 2$** is given in J. Fialová and O. Strauch (2010):

Every point $(\gamma_q(n), \gamma_q(n+1))$, $n = 0, 1, 2, \dots$, lie on the line segment

$$Y = X - 1 + \frac{1}{q^k} + \frac{1}{q^{k+1}}, \quad X \in \left[1 - \frac{1}{q^k}, 1 - \frac{1}{q^{k+1}} \right]$$

for $k = 0, 1, \dots$ and let T be their union. Because $\gamma_q(n)$ is u.d., then the sequence $(\gamma_q(n), \gamma_q(n+1))$ has a.d.f. $g(x, y)$ of the form

$$g(x, y) = |\text{Project}_x((\{0, x\} \times [0, y]) \cap T)|,$$

where Project_x is a projection of a two dimensional set to the x -axis. It is a copula and $g(x, y)$ can be computed explicitly as

$$g(x, y) = \begin{cases} 0 & \text{if } (x, y) \in A, \\ 1 - (1 - y) - (1 - x) = x + y - 1 & \text{if } (x, y) \in B, \\ y - \frac{1}{q^i} & \text{if } (x, y) \in C_i, \\ x - 1 + \frac{1}{q^{i-1}} & \text{if } (x, y) \in D_i, \end{cases}$$

$i = 1, 2, \dots$, where




Figure 1: Line segments containing $(\gamma_q(n), \gamma_q(n+1))$, $n = 1, 2, \dots$. The graph of von Neumann-Kakutani transformation.

(VI) **Formal solution.** Ch. Aisleitner and M. Hofer (2013): Let T denote von Neuman-Kakutani transformation described in Fig. 1. Define an s -dimensional curve $\{\gamma(t); t \in [0, 1)\}$, where $\gamma(t) = (t, T(t), T^2(t), \dots, T^{s-1}t)$. Then the searched a.d.f. is

$$g(x_1, x_2, \dots, x_s) = |\{t \in [0, 1]; \gamma(t) \in [0, x_1] \times [0, x_2] \times \dots \times [0, x_s]\}|,$$

where $|X|$ is the Lebesgue measure of set X . An explicit formula of a such a.d.f. for $s = 4$ is **open**.

(VI') For an arbitrary continuous $F(x_1, x_2, \dots, x_s)$ we have

$$\int_{[0,1]^s} F(x_1, x_2, \dots, x_s) dg(x_1, x_2, \dots, x_s) = \int_0^1 F(x, T(x), T^2(x), \dots, T^{s-1}(x)) dx.$$

Proof: Put $\gamma(n) = x_n$. Then

$$(\gamma_q(n), \dots, \gamma_q(n+s-1)) = (x_n, T(x_n), T^2(x_n), \dots, T^{s-1}(x_n))$$

and by Weyl's limit relation

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N F(\gamma_q(n), \dots, \gamma_q(n+s-1)) = \int_{[0,1]^s} F(x_1, x_2, \dots, x_s) dg(x_1, x_2, \dots, x_s)$$

and

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N F(x_n, T(x_n), \dots, T^{s-1}(x_n)) = \int_0^1 F(x, T(x), \dots, T^{s-1}(x)) dx.$$


(VII) **A.d.f. of** $(\gamma_q(n), \gamma_q(n+2)), n = 1, 2, \dots$. All terms of the sequence $(\gamma_q(n), \gamma_q(n+2)), n = 1, 2, \dots$, lie in the line segments

$$Y = X + \frac{2}{q}, \quad X \in \left[0, 1 - \frac{2}{q}\right), \text{ or}$$

$$Y = X + \frac{1}{q} + \frac{1}{q^{i+1}} + \frac{1}{q^{i+2}} - 1, \quad X \in \left[1 - \frac{1}{q} - \frac{1}{q^{i+1}}, 1 - \frac{1}{q} - \frac{1}{q^{i+2}}\right) \text{ or}$$

$$Y = X + \frac{1}{q} + \frac{1}{q^{i+1}} + \frac{1}{q^{i+2}} - 1, \quad X \in \left[1 - \frac{1}{q^{i+1}}, 1 - \frac{1}{q^{i+2}}\right)$$

for $i = 0, 1, \dots$. Divide $[0, 1]^2$ by the following figure



then we have the following explicit form of a.d.f. $g(x, y)$ of the sequence $(\gamma_q(n), \gamma_q(n+2))$

2)).

$$g(x, y) = \begin{cases} x & \text{if } (x, y) \in D_0, \\ y - \frac{2}{q} & \text{if } (x, y) \in C_0, \\ 0 & \text{if } (x, y) \in A_0, \\ y + x - 1 & \text{if } (x, y) \in B_0, \\ x - 1 + \frac{2}{q} & \text{if } (x, y) \in E_0, \\ y & \text{if } (x, y) \in F_0, \\ 0 & \text{if } (x, y) \in A', \\ x + y - 1 + \frac{1}{q} & \text{if } (x, y) \in B', \\ x - 1 + \frac{1}{q} + \frac{1}{q^i} & \text{if } (x, y) \in D'_i, \\ y - \frac{1}{q^{i+1}} & \text{if } (x, y) \in C'_i, \\ \frac{1}{q} & \text{if } (x, y) \in A'', \\ x + y - 1 & \text{if } (x, y) \in B'', \\ x - 1 + \frac{1}{q} + \frac{1}{q^i} & \text{if } (x, y) \in D''_i, \\ y - \frac{1}{q^{i+1}} & \text{if } (x, y) \in C''_i. \end{cases}$$

(VIII) **A.d.f. of** $(\gamma_q(n), \gamma_q(n+1), \gamma_q(n+2)), n = 1, 2, \dots$. An explicit form of $g(x, y, z)$ is given in Fialová, J., Mišík, L. and Strauch, O (2013) and it have 27 possibilities. For example, if $q \geq 3$ then

$$g(x, x, x) = \begin{cases} 0 & \text{if } x \in \left[0, \frac{2}{q}\right], \\ x - \frac{2}{q} & \text{if } x \in \left[\frac{2}{q}, 1 - \frac{1}{q}\right], \\ 3x - 2 & \text{if } x \in \left[1 - \frac{1}{q}, 1\right]. \end{cases}$$

(IX) As an applications, by the Weyl limit relation, we have

$$\begin{aligned} & \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} F(\gamma_q(n), \gamma_q(n+1), \gamma_q(n+2)) \\ &= \int_0^1 \int_0^1 \int_0^1 F(x, y, z) \, d_x \, d_y \, d_z \, g(x, y, z), \end{aligned}$$

where $F(x, y, z)$ is an arbitrary continuous function in $[0, 1]^3$.

(X) For the right-hand side of (IX) we can using

$$\begin{aligned}
& \int_0^1 \int_0^1 \int_0^1 F(x, y, z) \, d_x d_y d_z g(x, y, z) \\
&= F(1, 1, 1) - \int_0^1 g(1, 1, z) \, d_z F(1, 1, z) - \int_0^1 g(1, y, 1) \, d_y F(1, y, 1) \\
&\quad - \int_0^1 g(x, 1, 1) \, d_x F(x, 1, 1) + \int_0^1 \int_0^1 g(1, y, z) \, d_y d_z F(1, y, z) \\
&\quad + \int_0^1 \int_0^1 g(x, 1, z) \, d_x d_z F(x, 1, z) + \int_0^1 \int_0^1 g(x, y, 1) \, d_x d_y F(x, y, 1) \\
&\quad - \int_0^1 \int_0^1 \int_0^1 g(x, y, z) \, d_x d_y d_z F(x, y, z).
\end{aligned}$$

(XI) Example: Put $F(x, y, z) = \max(x, y, z)$. Then by (X)

$$\begin{aligned}
& \int_0^1 \int_0^1 \int_0^1 F(x, y, z) \, d_x d_y d_z g(x, y, z) \\
&= 1 - \int_0^1 \int_0^1 \int_0^1 g(x, y, z) \, d_x d_y d_z F(x, y, z) \\
&= 1 - \int_0^1 g(x, x, x) \, dx
\end{aligned}$$

and for $q \geq 3$ we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} \max(\gamma_q(n), \gamma_q(n+1), \gamma_q(n+2)) = \frac{1}{2} + \frac{2}{q} - \frac{3}{q^2}.$$

(XII) Example: Put $F(x, y, z) = \min(x, y, z)$. Then by (X) we have

$$\begin{aligned}
&= \int_0^1 \int_0^1 \int_0^1 F(x, y, z) \, d_x d_y d_z g(x, y, z) \\
&= 1 - 3 \cdot \frac{1}{2} + 2 \cdot \int_0^1 g(x, x, 1) \, dx + \int_0^1 g(x, 1, x) \, dx - \int_0^1 g(x, x, x) \, dx
\end{aligned}$$

which implies

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} \min(\gamma_2(n), \gamma_2(n+1), \gamma_2(n+2)) = \begin{cases} \frac{1}{2} - \frac{2}{q} + \frac{3}{q^2} & \text{if } q \geq 4, \\ \frac{1}{6} & \text{if } q = 3, \\ \frac{3}{16} & \text{if } q = 2. \end{cases}$$

(XII) Example: Put $F(x, y, z) = xyz$. By (X) we have

$$\begin{aligned}
& \int_0^1 \int_0^1 \int_0^1 F(x, y, z) \, d_x d_y d_z g(x, y, z) = 1 - 3 \cdot \frac{1}{2} + 2 \cdot \int_0^1 \int_0^1 g(x, y, 1) \, dx \, dy \\
& + \int_0^1 \int_0^1 g(x, 1, z) \, dx \, dz - \int_0^1 \int_0^1 \int_0^1 g(x, y, z) \, dx \, dy \, dz
\end{aligned}$$

and we find

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} \gamma_q(n) \cdot \gamma_q(n+1) \cdot \gamma_q(n+2) = \frac{q^4 - 3q^3 + 3q^2 + 2q + 2}{4q^4 + 4q^3 + 4q^2}$$

for $q \geq 3$.

Entered by O. Strauch.

CH. AISLEITNER – M. HOFER: *On the limit distribution of consecutive elements of the van der Corput sequence*, Uniform Distribution Theory **8** (2013), no. 1, 89–96.

O. BLAŽEKOVÁ: *Pseudo-randomness of van der Corput's sequences*, Math. Slovaca **59** (2009), no. 3, 291–298.

H. FAURE: *Discrépances de suites associées à un système de numération (en dimension un)*, Bull. Soc. Math. France **109** (1981), 143–182 (MR 82i:10069).

J. FIALOVÁ – L. MIŠÍK – O. STRAUCH: *An asymptotic distribution function of three-dimensional shifted van der Corput sequence*, pp. 27, INTEGERS: Electronic Journal of Combinatorial Number Theory, 2013 send.

J. FIALOVÁ – O. STRAUCH: *On two-dimensional sequences composed of one-dimensional uniformly distributed sequences*, Uniform Distribution Theory **6** (2011), 101–125.

J.H. HALTON: *On the efficiency of certain quasi-random sequences of points in evaluating multi-dimensional integrals*, Numer. Math. **2** (1960), 84–90 (MR 22#12688).

P. HELLEKALEK – H. NIEDERREITER: *Constructions of uniformly distributed sequences using the \mathbf{b} -adic method*, Uniform Distribution Theory **6** (2011), 185–200.

L.–K. HUA – Y. WANG: *Applications of Number Theory to Numerical Analysis*, Springer Verlag & Science Press, Berlin, Heidelberg, New York, Beijing, 1981 (83g:10034). (Chinese edition Science Press, Beijing, 1978, Zbl 451.10001).

P.D. PROINOV – V.S. GROZDANOV: *On the diaphony of the van der Corput – Halton sequence*, J. Number Theory **30** (1988), no. 1, 94–104 (MR 89k:11065).

I.M. SOBOĽ: *Evaluation of multiple integrals*, (Russian), Dokl. Akad. Nauk SSSR **139** (1961), no. 4, 821–823 (MR 25#3608).

I.M. SOBOĽ: *Multidimensional Quadrature Formulas and Haar Functions*, (Russian), Library of Applied Analysis and Computational Mathematics, Izd. "Nauka", Moscow, 1969, 288 pp. (MR 54#10952).

J.G. VAN DER CORPUT: *Verteilungsfunktionen III–VIII*, Proc. Akad. Amsterdam **39** (1936), 10–19, 19–26, 149–153, 339–344, 489–494, 579–590 (JFM 61.0204.01, 61.0204.02, 62.0206.06, 62.0207.01, 62.0207.02, 62.0207.03).

1.13 Sequences of differences

[SP, p. 2–7, 2.1.7]. The sequence

$$x_n \in [0, 1), \quad n = 1, 2, \dots,$$

is u.d. if and only if the sequence

$$|x_m - x_n|, \quad m, n = 1, 2, \dots,$$

has the a.d.f.

$$g(x) = 2x - x^2.$$

Here the double sequence $|x_m - x_n|$, for $m, n = 1, 2, \dots$, is ordered to an ordinary sequence y_n in such a way that the first N^2 terms of y_n are $|x_m - x_n|$ for $m, n = 1, 2, \dots, N$.

Open problem: Assuming u.d. of x_n , $n = 1, 2, \dots$ find a.d.f of the sequences

- $||x_m - x_n| - |x_k - x_l||$, $m, n, k, l = 1, 2, \dots$,
- $|||x_m - x_n| - |x_k - x_l|| - ||x_i - x_j| - |x_r - x_s||$, $m, n, k, l, i, j, r, s = 1, 2, \dots$,
etc.

NOTES: We can use the following method: Let us denote by $g_j(x)$ an asymptotic distribution function of the sequence of j th differences (thus $g_1(x) = 2x - x^2$). For k th moment we have

$$\int_0^1 x^k dg_{j+1}(x) = \int_0^1 \int_0^1 |x - y|^k dg_j(x) dg_j(y).$$

For $j = 1$ we have

$$\int_0^1 \int_0^1 |x - y|^k dg_1(x) dg_1(y) = \frac{8}{(k+1)(k+2)(k+4)}.$$

which implies

$$g_2(x) = \frac{8}{3}x - 2x^2 + \frac{1}{3}x^4.$$

Conjecture proposed by S.Steinerberger (2010): The density function $\frac{dg_j(x)}{dx}$ of the a.d.f. $g_j(x)$ of j -th iterated differences is of the form

$$\frac{dg_j(x)}{dx} = \frac{2^{2^j+j-1}}{2^j!} (x-1)^j p(x),$$

where $p(x)$ is a polynomial with integer coefficients. What can be said about $p(x)$?

For $j = 3$ he found

$$\frac{dg_3(x)}{dx} = \frac{8}{315} (x-1)^3 (-132 - 116x - 36x^2 + 3x^3 + x^4).$$

Proposed by O. Strauch.

O. STRAUCH: *On the L^2 discrepancy of distances of points from a finite sequence*, Math. Slovaca **40** (1990), 245–259 (MR 92c:11078).

1.14 Bernoulli numbers

Open problem. For Bernoulli numbers B_{2n} find the distribution

$$B_{2n} \bmod 1 \quad n = 1, 2, \dots$$

NOTES: (I) By von Staudt-Clausen formula $B_{2n} = A_{2n} - \sum_{(p-1)|2n} \frac{1}{p}$, where p are primes and A_{2n} are integers.

(II) F. Luca's comment: This problem was studied by P. Erdős and S.S. Wagstaff Jr. (1980). They proved that $\sum_{(p-1)|2n} \frac{1}{p}$ is everywhere dense in $[5/6, \infty)$.

Proposed by O. Strauch.

P. ERDŐS – S.S. WAGSTAFF, JR.: *The fractional parts of the Bernoulli numbers*, Illinois J. Math. **24** (1980), no. 1, 104–112 (MR550654 (MR 81c:10064)).

1.15 Ratio sequences.

[SP, p. 2–215, 2.22.2]. For an increasing sequence of positive integers x_n let $\underline{d}(x_n)$, and $\bar{d}(x_n)$ denote the lower and upper asymptotic density of x_n , resp., and $d(x_n)$ ($= \underline{d}(x_n) = \bar{d}(x_n)$) its asymptotic density if it exists. The double sequence, called the **ratio sequence of x_n** ,

$$\frac{x_m}{x_n}, \quad m, n = 1, 2, \dots,$$

is everywhere dense in $[0, \infty)$ assuming that one of the following conditions holds:

- (i) $d(x_n) > 0$,
- (ii) $\bar{d}(x_n) = 1$,
- (iii) $\underline{d}(x_n) + \bar{d}(x_n) \geq 1$,
- (iv) $\underline{d}(x_n) \geq 1/2$,
- (v) $A([0, x]; x_n) \sim \frac{cx}{\log^\alpha x}$, where $c > 0, \alpha > 0$ are constant, $A([0, x]; x_n) = \#\{n \in \mathbb{N}; x_n \in [0, x]\}$, and \sim denotes the asymptotically equivalence (i.e. the ratio of the left and the right-hand side tends to 1 as $x \rightarrow \infty$).

NOTES: (I) (i), (ii) and (v) were proved by T. Šalát (1969), for (iii) see O. Strauch and J.T. Tóth (1998) and (iv) follows from (iii).

(II) Strauch and Tóth (1998, Th. 2) proved that if the interval $(\alpha, \beta) \subset [0, 1]$ has an empty intersection with $\frac{x_m}{x_n}$ for $m, n = 1, 2, \dots$, then

$$\underline{d}(x_n) \leq \frac{\alpha}{\beta} \min(1 - \bar{d}(x_n), \bar{d}(x_n)), \quad \bar{d}(x_n) \leq 1 - (\beta - \alpha). \quad (1)$$

(III) S. Konyagin (1999, personal communication) improved the second inequality to

$$\bar{d}(x_n) \leq \frac{1 - \beta}{1 - \alpha\beta}. \quad (2)$$

Problem: Find a best possible estimation of $\bar{d}(x_n)$.

Solution: G. Grekos (2006, personal communication) notes that in (2) the equation is valid for the sequence $x_n, n = 1, 2, \dots$ defined in O. Strauch and J.T. Tóth (1998, Ex. 1) such that x_n is the sequence of all integer points lying in the intervals

$$(\gamma, \delta), (\gamma a, \delta a), (\gamma a^2, \delta a^2), \dots, (\gamma a^n, \delta a^n), \dots,$$

where γ, δ and a are positive real numbers satisfying $\gamma < \delta$ and $a > 1$. In this case the lower \underline{d} and upper \bar{d} asymptotic density of x_n can be given explicitly by

$$\underline{d}(x_n) = \frac{(\delta - \gamma)}{\gamma(a - 1)}, \quad \bar{d}(x_n) = \frac{(\delta - \gamma)a}{\delta(a - 1)},$$

and an interval (α, β) which does not contain points $\frac{x_m}{x_n}, m, n = 1, 2, \dots$ is

$$(\alpha, \beta) = \left(\frac{\delta}{\gamma a}, \frac{\gamma}{\delta} \right), \quad (3)$$

assuming $\delta/\gamma < \sqrt{a}$. All others subintervals of $[0, 1]$ with empty intersection have the form $(\alpha/a^i, \beta/a^i), i = 1, 2, \dots$. For interval (3) we have $\bar{d}(x_n) = \frac{1-\beta}{1-\alpha\beta}$ then (2) is the best possible.

(V) Strauch and Tóth (1998, Th. 6) also proved that (1) and (2) are also valid for interval (α, β) containing no accumulation points of $\frac{x_m}{x_n}, m, n = 1, 2, \dots$. If $X \subset [0, 1]$ is an union of such intervals, then

$$\bar{d}(x_n) \leq 1 - |X|,$$

where $|X|$ denotes the Lebesgue measure of X . Its improvement is **open**.

Proposed by O. Strauch.

T. ŠALÁT: *On ratio sets of sets of natural numbers*, Acta Arith. **15** (1968/69), 273–278 (MR **39**#4083).

O. STRAUCH – J.T. TÓTH: *Asymptotic density of $A \subset \mathbb{N}$ and density of the ration set $R(A)$* , Acta Arith. **87** (1998), no. 1, 67–78 (MR 99k:11020).

O. STRAUCH – J.T. TÓTH: *Distribution functions of ratio sequences*, Publ. Math. (Debrecen) **58** (2001), 751–778 (MR 2002h:11068).

1.16 Continued fractions.

[SP, p. 2–264, 2.26.8]. Let $\theta = [0; a_1, a_2, \dots]$ be an irrational number in $[0, 1]$ given by its continued fraction expansion and let $p_n(\theta)/q_n(\theta), n = 0, 1, 2, \dots$, be the corresponding sequence of its convergents. An **open problem** is to find, for the sequence

$$x_n = q_n(\theta) \pmod{2}$$

the frequency of each possible block $(\dots, 0, \dots, 1, \dots, 0, \dots)$ of length s which occurs in x_n as $(x_{n+1}, \dots, x_{n+s})$ for a special class of θ (e.g., with bounded a_i).

NOTES: R. Moeckel (1982) proved that, for almost all θ , the three possible blocks $(0, 1)$, $(1, 0)$ and $(1, 1)$ of length $s = 2$ occur in x_n with equal frequencies. The blocks of lengths $s = 3$ and $s = 4$ are investigated in V.N. Nolte (1990).

Proposed by O. Strauch.

R. MOECKEL: *Geodesic on modular surfaces and continued fractions*, Ergodic Theory Dynamical Systems **2** (1982), no. 1, 69–83 (MR 84k:58176).

V.N. NOLTE: *Some probabilistic results on the convergents of continued fractions*, Indag. Math. (N.S.) **1** (1990), no. 3, 381–389 (MR 92b:11053).

1.17 Strong uniform distribution.

Say a sequence of integers $(a_n)_{n=1}^{\infty}$ is in \mathbf{A}^* for a class of measurable functions \mathbf{A} on $[0, 1)$ if

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(\{a_n x\}) = \int_0^1 f(t) dt \text{ a.e.}$$

Here of course, for a real number y we have used $\{y\}$ to denote its fractional part. In a paper solving a well known classical problem of A. Khinchin's [Kh], J. M. Marstrand [M] also showed that if q_1, \dots, q_k is a finite list of coprime natural numbers all greater than one then the semigroup it generates, $\mathbf{m} = (m_l)_{l=1}^{\infty} = \{q_1^{i_1} \cdots q_k^{i_k} : (i_1, \dots, i_k) \in (\mathbb{Z}_0^+)^k\}$ when ordered by size is in $(L^\infty)^*$. To do this he invoked D. Birkhoff's pointwise ergodic theorem [W] and the following lemma:

For strictly increasing sequences of natural numbers $(a_n)_{n=1}^{\infty}$ and $(b_n)_{n=1}^{\infty}$ if $G(u) = \{(r, s) : a_r b_s \leq u\}$, $(u = 1, 2, \dots)$ and $f \in L^\infty$ then

$$\lim_{u \rightarrow \infty} \frac{1}{|G(u)|} \sum_{(r,s) \in G(u)} f(\{a_r b_s x\}) = \int_0^1 f(t) dt \text{ a.e.}$$

Here for a finite set A we have used $|A|$ to denote its cardinality. R.C. Baker [B] asked if \mathbf{m} is in $(L^1)^*$. This was proved by the author [Na1] using A. A. Tempelman's generalization of Birkhoff's pointwise ergodic theorem [T]. With a view to applications to sequences other than \mathbf{m} , it would be interesting to know if an L^1 version of Marstrand's lemma is true. Some partial results in this direction are known. In [Na2] we show that for strictly

increasing sequences of natural numbers $a = (a_r)_{r=1}^\infty$ and $b = (b_s)_{s=1}^\infty$, both of which are $(L^p)^*$ sequences for all $p > 1$, if there exists $C > 0$ such that

$$|\{r : a_r \leq u\}||\{s : b_s \leq u\}| \leq C|\{(r, s) : a_r b_s \leq u\}|, \quad (1)$$

for $(u = 1, 2, \dots)$ then $a \circ b = \{a_r b_s : (r, s) \in \mathbf{N}^2\}$ (the sequence of products of pairs of elements in a and b) once ordered by size is also an $(L^p)^*$ sequence. An open question is whether this result from [Na2] is true for $p = 1$. We have the following partial result. Let

$$a_1 = (a_{1,i})_{i=1}^\infty, \dots, a_k = (a_{k,i})_{i=1}^\infty,$$

denote finitely many $(L^1)^*$ sequences, and for a sequence a , let

$$G_a(u) = |\{i : a_i \leq u\}|.$$

Also let $a_1 \circ \dots \circ a_k$ denote the set

$$\{b_1 \cdots b_k : b_1 \in a_1, \dots, b_k \in a_k\},$$

counted with multiplicity and ordered by absolute value. Suppose there exists $K > 0$ such that for all $u \geq 1$

$$|G_{a_1}(u)| \cdots |G_{a_k}(u)| \leq K|G_{a_1 \circ \dots \circ a_k}(u)|.$$

Then if $\log_+ |x| = \log \max(1, |x|)$ we show that $a_1 \circ \dots \circ a_k$ is an $(L(\log_+ L)^{k-1})^*$ sequence [Na3]. To be more specific, we ask if the space $(L(\log_+ L)^{k-1})^*$ can be replaced by $(L^1)^*$. A second open question is whether a condition like (1) necessary for any of these results. As Marstrand observed the answer is no when $p = \infty$. It might be the case that if for fixed $p \in [1, \infty]$ if all three sequences $a = (a_r)_{r=1}^\infty$, $b = (b_s)_{s=1}^\infty$ and $a \circ b$ are $(L^p)^*$ then (1) automatically holds. This too is unknown.

Proposed by R. Nair.

[B] R.C. BAKER: *Riemann sums and Lebesgue integrals*, Quart. J. Math. Oxford (2) **27** (1976), 191–198.

[K] A. KHINCHIN: *Ein Satz über Kettenbrüche, mit Arithmetischen Anwendungen*, Math. Zeit. **18** (1923), 289–306.

[M] J.M. MARSTRAND: *On Khinchine's conjecture about strong uniform distribution*, Proc. Lond. Math. Soc. **21** (1977), 540–556.

[Na1] R. NAIR: *On strong uniform distribution*, Acta Arith. **54** (1990), 183–193.

[Na2] R. NAIR: *On strong uniform distribution II*, Monatsh. Math. **132** (2001), no. 4, 341–348.

[Na3] R. NAIR: *On strong uniform distribution III*, Indag. Math. (N.S.) **14** (2003), no. 2, 233–240.

[T] A.A. TEMPELMAN: *Ergodic theorems for general dynamical systems*, Soviet. Math. Dokl. **8** (1967), 1213–1216 (English Translation).

[W] P. WALTERS: *Introduction to Ergodic Theory*, Graduate Texts in Mathematics, Vol. 79, Springer-Verlag, Berlin, 1981.

1.18 Algebraic dilatations.

Let $\theta_1, \dots, \theta_k$ denote a finite set of real algebraic numbers all greater than 1 and let $(m_l)_{l=1}^\infty$ denote the semigroup generated multiplicatively by these numbers given an order consistent with the magnitude of the elements of this semigroup. Suppose $f \in L^p([0, 1])$ for some $p \in [1, \infty)$. Under what conditions on p and the sequence $(m_l)_{l=1}^\infty$ is it the case that

$$\lim_{L \rightarrow \infty} \frac{1}{L} \sum_{l=1}^L f(\{m_l x\}) = \int_0^1 f(t) dt \text{ a.e.}$$

This question is probably best understood by comparison with the cases where the $\theta_1, \dots, \theta_k$ are coprime natural numbers, in which case the result is true for $p = 1$ [Na1] and the case where $k = 1$ in which case the result is known for $p = 2$ [Bo][vPS]. In unpublished work the author has also shown that the result is true for $p > 1$ if only one of the set $\theta_1, \dots, \theta_k$ is not a rational integer. It seems likely some other arithmetic condition on the $\theta_1, \dots, \theta_k$ is necessary. For instance is the result true for instance when $p > 1$ assuming the natural logarithms of the numbers $\theta_1, \dots, \theta_k$ are linearly independent over the rationals? Are there circumstances where the assumption that k is finite can be dropped? The answer is yes for $\theta_1, \theta_2, \dots$ chosen to be some rapidly growing rational primes as shown in [L].

Proposed by R. Nair.

Bo J. BOURGAIN: *The Riesz-Raikov theorem for algebraic numbers*, in: *Festschrift in honor of I.I. Piatetski-Shapiro on the occasion of his sixtieth birthday, Part II (Ramat Aviv, 1989)*, Weizmann, Jerusalem, 1990, pp. 1–45.

[L] Y. LACROIX: *On strong uniform distribution. III*, *Monatsh. Math.* **143** (2004), no. 1, 13–19.

[vPS] A.J. VAN DER POORTEN – H.P. SCHLICKEWEI: *A Diophantine problem in harmonic analysis*, *Math. Proc. Cambridge Philos. Soc.* **108** (1990), no. 3, 417–420.

1.19 Subsequence ergodic theorems.

Suppose (X, β, μ) is a measure space. Say T a map from X to itself is measurable if $T^{-1}A := \{x : Tx \in A\} \in \beta$ for all A in the σ -algebra β . We say a measurable map T from X to itself is measure preserving if $\mu(T^{-1}A) = \mu(A)$ for all $A \in \beta$. Building on earlier work of J. Bourgain [Bo1],[Bo2],[Bo3], the author showed [Na4] [Na5] that if ϕ is a non-constant polynomial mapping the natural numbers to themselves, $(p_n)_{n=1}^\infty$ is the sequence of rational primes and if for $p > 1$ we have $f \in L^p(X, \beta, \mu)$ then $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(T^{\phi(p_n)}x)$ exists μ almost everywhere. See also [Wi].

Our first question is what happens when $p = 1$. The analogous question for the case where p_n is the n^{th} natural number is also open. For the second question suppose T_1, \dots, T_k is a finite set of commuting measure preserving maps on (X, β, μ) and that ϕ_1, \dots, ϕ_k are nonconstant polynomials mapping the natural numbers to themselves and for $p > 1$ that $f \in L^p(X, \beta, \mu)$. Then is it the case that $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(T_1^{\phi_1(p_n)} \dots T_k^{\phi_k(p_n)} x)$ μ exists μ almost everywhere? In the case where p_n is the n^{th} natural number this is a theorem, at least when $p = 2$ [Bo2].

Proposed by R. Nair.

[Bo1] J. BOURGAIN: *On the maximal ergodic theorem for certain subsets of the integers*, Israel J. Math. **61** (1988), no. 1, 39–72.

[Bo2] J. BOURGAIN: *On the pointwise ergodic theorem on L^p for arithmetic sets*, Israel J. Math. **61** (1988), no. 1, 73–84.

[Bo2] J. BOURGAIN: *Pointwise ergodic theorems for arithmetic sets. With an appendix by the author, Harry Furstenberg, Yitzhak Katznelson and Donald S. Ornstein.*, Inst. Hautes Études Sci. Publ. Math. (1989), 5–45.

[Na4] R. NAIR: *On polynomials in primes and J. Bourgain's circle method approach to ergodic theorems*, Ergod. Th. & Dynam. Sys. **11** (1991), 485–499.

[Na5] R. NAIR: *On polynomials in primes and J. Bourgain's circle method approach to ergodic theorems II*, Studia Math. **105** (3) (1993), 207–233.

[Wi] M. WIERDL: *Pointwise ergodic theorem along the prime numbers*, Israel J. Math. **64** (1988), 315–336.

1.20 Square functions for subsequence ergodic averages.

For a probability space (X, β, μ) , measure preserving $T : X \rightarrow X$, a non-constant polynomial ϕ mapping the natural numbers to themselves and a μ measurable function f defined on X and a strictly increasing sequence of integers $(N_k)_{k=1}^{\infty}$, set $A_N f(x) = \frac{1}{N} \sum_{n=1}^N f(T^{\phi(n)} x)$ ($N = 1, 2, \dots$) and set

$$S(f)(x) = \left(\sum_{k=1}^{\infty} |A_{N_{k+1}} f(x) - A_{N_k} f(x)|^2 \right)^{\frac{1}{2}}.$$

In the situation where $\phi(n) = n$ and $(N_k)_{k=1}^{\infty}$ is any strictly increasing sequence of natural numbers it is shown in [JOR] that there exists $C > 0$ such that $\mu(\{x : S(f)(x) > \lambda\}) \leq C \frac{\|f\|_1}{\lambda}$. One implication of this is that for any $p > 1$ there exists $C_p > 0$ such that $\|S(f)\|_p \leq C_p \|f\|_p$. Results of this sort provide an alternative means, to almost everywhere convergence, of measuring the stability of the averages $(A_N f)_{N=1}^{\infty}$. Ideally we would like to prove an analogue of the [JOR] inequality for general ϕ or if it were not true to find out the extent to which it was and how and when it

fails. One approach to questions of these sorts is via spectral theory and this reduces to a study of the behaviour of exponential sums of the form $a_n(\alpha) = \frac{1}{n} \sum_{l=1}^n e^{2\pi i \phi(n)\alpha}$ ($n = 1, 2, \dots$). In the case where $\phi(n) = n$ these are for each α , averages of geometric progressions and consequently have well understood distributions. For more general ϕ these are Weyl sums, are more complicated and our understanding is less complete. What is known follows from an application of the Hardy-Littlewood circle method. Using this, for $N_k = k$ ($k = 1, 2, \dots$), the author has shown that for each $p > 1$ there exists $C_p > 0$ such that we have $\|S(f)\|_p \leq C_p \|f\|_p$. The same inequality is also true in the case $p = 2$ where $1 < a \leq \frac{N_{k+1}}{N_k} < b$ for some a and b . These results are not as yet published [Na6] though see [NW] where variants of the method involved appear.

Proposed by R. Nair.

[Na6] R. NAIR: *Polynomial ergodic averages and square functions*, (unpublished manuscript).

[NW] R. NAIR – M. WEBER: *On variation functions for subsequence ergodic averages*, *Monatsh. Math.* **128** (1999), 131-150.

[JOR] R.L. JONES – I.V. OSTROVSKII – J.M. ROSENBLATT: *Square functions in ergodic theory*, *Ergodic Theory Dynam. Systems* **16** (1996), no. 2, 267–305.

1.21 Sets of integers of positive density

In this section for simplicity we confine attention to sets of integers of positive density in \mathbb{N} , though much of what we deal with can be meaningfully discussed in higher dimensions. For a set of natural numbers $S \subset \mathbb{N}$ and a countable collection $\mathbf{I} = (I_n)_{n \geq 1}$ of intervals with $|I_n|$ tending to infinity as n does, let

$$B(S, \mathbf{I}) = \limsup_{n \rightarrow \infty} \frac{|S \cap I_n|}{|I_n|}.$$

We call

$$B(S) = \sup_{\mathbf{I}} B(S, \mathbf{I}),$$

where the supremum is taken over all collections \mathbf{I} , the Banach density of S . The second definition we need is that of upper density defined as

$$d^*(S) = \limsup_{n \rightarrow \infty} \frac{|S \cap [0, n]|}{n}.$$

If the limit exists, we call $d^*(S)$ the density of S and denote it by $d(S)$. Plainly $B(S) > 0$ if $d^*(S) > 0$. A sequence of integers $\mathbf{k} = (k(n))_{n \geq 1}$ is called *intersective* if given any $S \subset \mathbb{N}$ with $B(S) > 0$ there exist a and b in

S such that $a - b$ is in \mathbf{k} . A sequence of integers $\mathbf{k} = (k(n))_{n \geq 1}$ is called a set of *Poincaré recurrence* if given any (X, β, μ) any measure preserving transformation $T : X \rightarrow X$, and any A in β with $\mu(A) > 0$ there exists k in \mathbf{k} such that

$$\mu(A \cap T^k A) > 0.$$

The terminology is motivated by the fact that as proved by H. Poincaré, \mathbf{k} is a set of Poincaré recurrence in the case $k(n) = n$. What makes ergodic theory relevant to the study of intersectivity is that as proved by A. Bertrand-Mathis [Be] and Furstenberg [F] a sequence on integers is a set of intersectivity if and only if it is a set of Poincaré recurrence. Furstenberg used this viewpoint to show that when $k(n) = n^2$ ($n = 1, 2, \dots$) then $\mathbf{k} = (k(n))_{n=1}^{\infty}$ is intersective. We say a sequence $\mathbf{k} = (k_n)_{n=1}^{\infty}$ is Hartman uniformly distributed [Ha] if for every non-integer θ we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N e^{2\pi i \theta k_n} = 0.$$

It can be shown that any \mathbf{k} that is Hartman uniformly distributed is also intersective [Bo]. E. Szemerédi [S] showed that if $B(S) > 0$ then S contains arithmetic progressions of arbitrary length. This was shown by Furstenberg to be a consequence of the fact that given any (X, β, μ) any measure preserving transformation $T : X \rightarrow X$, and any A in β with $\mu(A) > 0$ there exists a natural number n such that [F]

$$(1) \quad \mu(A \cap T^n A \cap \dots \cap T^{n(k-1)} A) > 0.$$

Motivated by (1) it would be interesting to decide whether if \mathbf{k} is Hartman uniformly distributed then given any $S \subset \mathbb{N}$ with $B(S) > 0$ there exist $R \subset \mathbb{N}$ with $d(R) > 0$ existing and $d(R) \geq B(S)$ such that for any finite subset $\{h(1), \dots, h(l)\}$ of R ,

$$B(S \cap S + k(h(1)) \cap \dots \cap S + k(h(l))) > 0.$$

The question arose in a conversation with H. Furstenberg at the Newton Institute in 2000. So far it has been possible to prove this hypothesis subject to conditions similar to but stronger than Hartman uniform distribution [Na7][NW][NZ]. A theorem of V. Bergelson and A. Leibman [BL] is that given any polynomials

$P_1(x), \dots, P_k(x)$ with integer coefficients such that $0 = P_1(0) = P_2(0) = \dots = P_k(0)$, any (X, β, μ) any measure preserving transformation $T : X \rightarrow$

X and any A in β with $\mu(A) > 0$ there exists a natural number n such that

$$\mu(A \cap T^{P_1(n)} A \cap \dots \cap T^{P_k(n)} A) > 0.$$

This implies both Szemerédi's theorem and the intersectivity of \mathbf{k} where $k(n) = n^2$. It can also be shown [Na8] that given any polynomial $Q(x)$ such that for each non-zero integer m there exists another integer $l(m)$ with $(m, Q(l(m))) = 1$ any probability space (X, β, μ) any measure preserving transformation $T : X \rightarrow X$, any A in β with $\mu(A) > 0$ there exists a prime p such that

$$\mu(A \cap T^{Q(p)} A) > 0.$$

This leads one to conjecture that given polynomials $Q_1(x), \dots, Q_k(x)$, all with the property assumed for $Q(x)$, then given any probability space (X, β, μ) , any measure preserving transformation $T : X \rightarrow X$, and any A in β with $\mu(A) > 0$, there exists a prime number p such that

$$\mu(A \cap T^{Q_1(p)} A \cap \dots \cap T^{Q_k(p)} A) > 0.$$

If proved this conjecture would imply analogues of both the Szemerédi theorem and the Furstenberg intersectivity phenomenon with the integer n chosen to be a prime. Examples of polynomials Q aren't quite as easy to construct as examples of polynomials satisfying the Bergelson-Leibman condition but they include $Q(x) = x^n + 1$ or $x^n - 1$ for any natural number n . Recently B. Green and T. Tao have proved that the set of primes contain arithmetic progressions of arbitrary length. It becomes natural to ask if, by analogy with the Bergelson-Leibman theorem, the primes contain arithmetic progressions of arbitrary length whose common difference is the value $P(a)$ say, for a given integer valued polynomial $P(x)$ with $P(0) = 0$ and some natural number a . We might also ask whether the primes contain arithmetic progressions of arbitrary length whose common difference is $Q(p)$ for a given polynomial Q such that for each non-zero integer m there exists another integer $l(m)$ with $(m, Q(l(m))) = 1$ and some prime p .

Proposed by R. Nair.

[Be] A. BERTRAND-MATHIS: *Ensembles intersectifs et recurrence de Poincaré*, Israel J. Math. **55** (1986), 184–198.

[BL] V. BERGELSON – A. LEIBMAN: *Polynomial extentions of van der Waerden's and Szemerédi's theorems*, J. Amer. Math. Soc. **9** (1996), no. 3, 725–753.

[Bo] M. BOSHERNITZAN: *Homogeneously distributed sequences and Poincaré sequences of integers of sublacunary growth*, Monatsh. Math. **96** (1983), no. 3, 173–181.

[F] H. FURSTENBERG: *Recurrence in Ergodic Theory and Combinatorial Number Theory*, Princeton University Press, 1981.

[Ha] S. HARTMAN: *Remarks on equidistribution on non-compact groups*, Compositio **16** (1964), 66–71.

- [Na7] R. NAIR: *On uniformly distributed sequences of integers and Poincaré recurrence II*, Indag. Math., N. S. **9** (1998), no. 3, 405–415.
- [Na8] R. NAIR: *On certain solutions of the diophantine equation $x - y = p(z)$* , Acta Arith. **LXII** (1992), no. 1, 61–71.
- [NW] R. NAIR – M. WEBER: *Random perturbations of intersective sets are intersective*, Indag. Math. (N.S.) **15** (2004), no. 3, 373–381.
- [NZ] R. NAIR – P. ZARIS: *On certain sets of integers and intersectivity*, Proc. Camb. Phil. Soc. **131** (2001), no. 1, 157–164.
- [S] E. SZEMERÉDI: *On sets of integers containing no k terms in arithmetic progression*, Acta Arith. **27** (1975), 199–245.

1.22 Uniform distribution of the weighted sum-of-digits function.

Let $\gamma = (\gamma_0, \gamma_1, \dots)$ be a sequence in \mathbb{R} and let $q \in \mathbb{N}$, $q \geq 2$. For $n \in \mathbb{N}_0$ with base q representation $n = n_0 + n_1q + n_2q^2 + \dots$ define the *weighted q -ary sum-of-digits function* by

$$s_{q,\gamma}(n) := \gamma_0 n_0 + \gamma_1 n_1 + \gamma_2 n_2 + \dots.$$

For $d \in \mathbb{N}$, weight-sequences $\gamma^{(j)} = (\gamma_0^{(j)}, \gamma_1^{(j)}, \dots)$ in \mathbb{R} and $q_j \in \mathbb{N}$, $q_j \geq 2$, $j \in \{1, \dots, d\}$, define

$$s_{q_1, \dots, q_d, \gamma}(n) := (s_{q_1, \gamma^{(1)}}(n), \dots, s_{q_d, \gamma^{(d)}}(n)),$$

where $\gamma = (\gamma_0, \gamma_1, \dots)$ and $\gamma_k = (\gamma_k^{(1)}, \dots, \gamma_k^{(d)})$ for $k \in \mathbb{N}_0$.

Open question: Let $q_1, \dots, q_d \geq 2$ be pairwise coprime integers. Under which conditions on the weight-sequences $\gamma^{(j)} = (\gamma_0^{(j)}, \gamma_1^{(j)}, \dots)$ in \mathbb{R} , $j \in \{1, \dots, d\}$, is the sequence

$$s_{q_1, \dots, q_d, \gamma}(n) \bmod 1, \quad n = 0, 1, 2, \dots \tag{1}$$

u.d. mod 1?

Proposed by F. Pillichshammer.

NOTES:

(I) For example if $\gamma_k^{(j)} = q_j^{-k-1}$ for all $j \in \{1, \dots, d\}$ and all $k \in \mathbb{N}_0$, then we obtain the d -dimensional van der Corput-Halton sequence which is well known to be uniformly distributed modulo one.

(II) If $\gamma_k^{(j)} = q_j^k \alpha_j$ for all $j \in \{1, \dots, d\}$ and all $k \in \mathbb{N}_0$, then the sequence (1) is the sequence $(\{n(\alpha_1, \dots, \alpha_d)\})_{n \geq 0}$ which is well known to be uniformly distributed modulo one if and only if $1, \alpha_1, \dots, \alpha_d$ are linearly independent over \mathbb{Q} .

(III) If $\gamma_k^{(j)} = \alpha_j \in \mathbb{R}$ for all $j \in \{1, \dots, d\}$ and all $k \in \mathbb{N}_0$, then it was shown by M. Drmota and G. Larcher (2001) that the sequence (1) is u.d. mod 1 if and only

if $\alpha_1, \dots, \alpha_d \in \mathbb{R} \setminus \mathbb{Q}$.

(IV) For $q_1 = \dots = q_d = q$ it was shown by F. Pillichshammer (2007) that the sequence (1) is u.d. mod 1 if and only if for every $\mathbf{h} \in \mathbb{Z}^d \setminus \{\mathbf{0}\}$ one of the following properties holds: Either

$$\sum_{\substack{k=0 \\ \langle \mathbf{h}, \boldsymbol{\gamma}_k \rangle q \notin \mathbb{Z}}}^{\infty} \|\langle \mathbf{h}, \boldsymbol{\gamma}_k \rangle\|^2 = \infty$$

or there exists a $k \in \mathbb{N}_0$ such that $\langle \mathbf{h}, \boldsymbol{\gamma}_k \rangle \notin \mathbb{Z}$ and $\langle \mathbf{h}, \boldsymbol{\gamma}_k \rangle q \in \mathbb{Z}$. Here $\|\cdot\|$ denotes the distance to the nearest integer, i.e., for $x \in \mathbb{R}$, $\|x\| = \min_{k \in \mathbb{Z}} |x - k|$ and $\langle \cdot, \cdot \rangle$ is the usual inner product.

(V) The generalization can be found in R. Hofer, G. Larcher and F. Pillichshammer (2007), where a similar result was proved with the weighted sum-of-digits function replaced by a generalized weighted digit-block-counting function.

(VI) R. Hofer (2007) proved: Let $q_1, \dots, q_d \geq 2$ be pairwise coprime integers and $\gamma^{(1)}, \dots, \gamma^{(d)}$ be given weight sequences in \mathbb{R} . If for each dimension $j \in \{1, \dots, d\}$ the following sum

$$\sum_{i=0}^{\infty} \left\| h(\gamma_{2^{i+1}}^{(j)} - q_j \gamma_{2^i}^{(j)}) \right\|^2$$

is divergent for every nonzero integer h , then the sequence (1) is u.d. in $[0, 1]^d$.

M. DRMOTA – G. LARCHER: *The sum-of-digits function and uniform distribution modulo 1*, J. Number Theory **89** (2001), 65–96.

R. HOFER: *Note on the joint distribution of the weighted sum-of-digits function modulo one in case of pairwise coprime bases*, Uniform Distribution Theory **2** (2007), no. 1, 1–10.

R. HOFER – G. LARCHER – F. PILLICHSHAMMER: *Average growth-behavior and distribution properties of generalized weighted digit-block-counting functions*, (2007), (Submitted).

F. PILLICHSHAMMER: *Uniform distribution of sequences connected with the weighted sum-of-digits function*, Uniform Distribution Theory **2** (2007), no. 1, 1–10.

1.23 Moment problem.

1.23.1 Truncated Hausdorff moment problem

Recovered a d.f. $g(x)$, given its moments

$$s_n = \int_0^1 x^n dg(x), \quad n = 1, 2, \dots, N. \quad (1)$$

The set of all points (s_1, s_2, \dots, s_N) in $[0, 1]^N$ for which there exists d.f. $g(x)$ satisfying (1) is called the N -th moment space $\boldsymbol{\Omega}_N$. It can be shown:

(i) the point (s_1, s_2, \dots, s_N) belongs to the moment space $\boldsymbol{\Omega}_N$ if and only if $\sum_{i=0}^k (-1)^i \binom{k}{i} s_{i+j} \geq 0$ for all $j, k = 0, 1, 2, \dots, N$.

(ii) $\boldsymbol{\Omega}_N$ is a simply connected, convex, and closed subset of $[0, 1]^N$.

(iii) If the point (s_1, s_2, \dots, s_N) belongs to the interior of the moment space $\mathbf{\Omega}_N$ the truncated moment problem (1) has infinitely many solutions.

(iv) If (s_1, s_2, \dots, s_N) belongs to the boundary of the $\mathbf{\Omega}_N$, the (1) has a unique solution $g(x)$.

(v) If the sequence $x_n \in [0, 1)$, $n = 1, 2, \dots$, satisfies $\lim \frac{1}{N} \sum_{n=1}^N x_n = s_1$, $\lim \frac{1}{N} \sum_{n=1}^N x_n^2 = s_2, \dots$, $\lim \frac{1}{N} \sum_{n=1}^N x_n^N = s_N$, where (s_1, s_2, \dots, s_N) belongs to the boundary of the $\mathbf{\Omega}_N$, then x_n has an a.d.f. $g(x)$.

Exact characterization of the moment space $\mathbf{\Omega}_N$ can be found in S. Karlin and L.S. Shapley (1953), also see G.A. Athanassoulis and P.N. Gavriliadis (2002).

1.23.2 L^2 moment problem

Given a triple of numbers $(X_1, X_2, X_3) \in [0, 1]^3$ O. Strauch (1994) gave a complete solution to the moment problem

$$(X_1, X_2, X_3) = \left(\int_0^1 g(x) dx, \int_0^1 xg(x) dx, \int_0^1 g^2(x) dx \right)$$

in d.f. $g(x) : [0, 1] \rightarrow [0, 1]$. He expresses the boundary of the body

$$\Omega = \left\{ \left(\int_0^1 g(x) dx, \int_0^1 xg(x) dx, \int_0^1 g^2(x) dx \right); g \text{ is d.f.} \right\}$$

as Π_1, \dots, Π_6 surfaces and the curve Π_7 such that for $(X_1, X_2, X_3) \in \cup_{i=1}^6 \Pi_i$ the moment problem has unique solution, for $(X_1, X_2, X_3) \in \Pi_7$ exactly two solutions, and in the interior of Ω has infinitely many solutions (see [SP, 2–20, 2.2.21] for exact results). Now, if a sequence x_n , $n = 1, 2, \dots$, in $[0, 1]$ has limits

$$X_1 = 1 - \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N x_n,$$

$$X_2 = \frac{1}{2} - \frac{1}{2} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N x_n^2 \text{ and}$$

$$X_3 = 1 - \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N x_n - \frac{1}{2} \lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{m,n=1}^N |x_m - x_n|$$

and if $(X_1, X_2, X_3) \in \bigcup_{1 \leq i \leq 7} \Pi_i$, then the sequence x_n possess an asymptotic distribution function $g(x)$.

Open problem is to solve the moment problem

$$(X_1, X_2, X_3, X_4) = \left(\int_0^1 g(x) dx, \int_0^1 xg(x) dx, \int_0^1 x^2g(x) dx, \int_0^1 g^2(x) dx \right).$$

E.g. for $g(x) = 2x - x^2$ it has the unique solution.

Proposed by O. Strauch.

G.A. ATHANASSOULIS – P.N. GAVRILIADIS: *The truncated Hausdorff moment problem solved by using kernel density functions*, Probabilistic Engineering Mechanics **17** (2002), 273–291.

S. KARLIN – L.S. SHAPLEY: *Geometry of moment spaces*, AMS Memoires, Vol. 12, Providence RI, 1953.

O. STRAUCH: *A new moment problem of distribution functions in the unit interval*, Math. Slovaca **44** (1994), no. 2, 171–211 (MR 95i:11082).

1.24 Scalar product.

Let $\mathbf{x}_n = (x_{n,1}, \dots, x_{n,s})$ and $\mathbf{y}_n = (y_{n,1}, \dots, y_{n,s})$ be infinite sequences in $[0, 1]^s$. Assume that the sequence $(\mathbf{x}_n, \mathbf{y}_n)$, $n = 1, 2, \dots$, is u.d. in $[0, 1]^{2s}$. Then the sequence of the inner (i.e. scalar) products

$$x_n = \mathbf{x}_n \cdot \mathbf{y}_n = \sum_{i=1}^s x_{n,i} y_{n,i}, \quad n = 1, 2, \dots$$

has the a.d.f. $g_s(x) = |\{(\mathbf{x}, \mathbf{y}) \in [0, 1]^{2s}; \mathbf{x} \cdot \mathbf{y} < x\}|$ on the interval $[0, s]$, and $g_s(x) = (-1)^s \int_{\substack{x_1 + \dots + x_s < x \\ x_1 \in [0, 1], \dots, x_s \in [0, 1]}} 1 \cdot \log x_1 \dots \log x_s \, dx_1 \dots dx_s$.

For $x \in [0, 1]$ we have

$$g_1(x) = x - \log x,$$

$$g_2(x) = \frac{x^2}{2} \left((\log x)^2 - 3 \log x + \frac{7}{2} - \frac{1}{6} \pi^2 \right),$$

$$g_3(x) = \frac{x^3}{27} \left(-\frac{9}{2} (\log x)^3 + \frac{99}{4} (\log x)^2 + \left(-\frac{255}{4} + \frac{9}{4} \pi^2 \right) \log x + \frac{575}{8} - \frac{33}{8} \pi^2 - 9\zeta(3) \right),$$

$$g_s(x) = (-1)^s x^s \sum_{j=0}^s \binom{s}{j} (\log x)^{s-j} \frac{1}{(s-j)!} \cdot$$

$$\int_{[0, 1]^j} \prod_{i=1}^j (\log x_1 + \dots + \log x_{j-1} + \log(1 - x_j)) x_1^{s-1} \dots x_j^{s-j} \, dx_1 \dots dx_j.$$

Open is the explicit formula of $g_s(x)$ for $x \in [1, s]$.

NOTES:

(I) O. Strauch (2003). The formula for $g_s(x)$ with $x \in [0, 1]$ was proved by L. Habsieger (Bordeaux) (personal communication). A motivation is an application of $g_s(x)$ to one-time pad cipher, see O. Strauch (2004).

(II) E. Hlawka (1982) investigated the question of the distribution of the scalar product of two vectors on an s -dimensional sphere and also the problem of the associated discrepancies.

Proposed by O. Strauch.

E. HLAJKA: *Gleichverteilung auf Produkten von Sphären*, J. Reine Angew. Math. **330** (1982), 1–43 (MR 83i:10066).

O. STRAUCH: *On distribution functions of sequences generated by scalar and mixed product*, Math. Slovaca **53** (2003), no. 5, 467–478 (MR 2038514).

O. STRAUCH: *Some modification of one-time pad cipher*, Tatra Mt. Math. Publ. **29** (2004), 157–171.

1.25 Determinant.

Let $\mathbf{x}_n^{(i)} = (x_{n,1}^{(i)}, \dots, x_{n,s}^{(i)})$, $i = 1, \dots, s$, be infinite sequences in the s -dimensional ball $B(r)$ with the center at $(0, \dots, 0)$ and radius r . Assume that these sequences are u.d. and statistically independent in $B(r)$, i.e., $(\mathbf{x}_n^{(1)}, \dots, \mathbf{x}_n^{(s)})$ is u.d. in $B(r)^s$. Then the sequence

$$x_n = |\det(\mathbf{x}_n^{(1)}, \dots, \mathbf{x}_n^{(s)})|$$

has the a.d.f. $g_s(r, x)$ defined on the interval $[0, r^s]$ by

$$g_s(r, x) = \frac{|\{(\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(s)}) \in B(r)^s; |\det(\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(s)})| < x\}|}{|B(r)|^s},$$

and for $\lambda = \frac{x}{r^s}$ there exists $\tilde{g}_s(\lambda)$ such that $g_s(r, x) = \tilde{g}_s(\lambda)$ if $\lambda \in [0, 1]$.

Here we have

$$\tilde{g}_1(\lambda) = \lambda,$$

$$\tilde{g}_2(\lambda) = \frac{2}{\pi}(1 + 2\lambda^2) \arcsin \lambda + \frac{6}{\pi} \lambda \sqrt{1 - \lambda^2} - 2\lambda^2,$$

$$\tilde{g}_3(\lambda) = 1 + \frac{9}{4} \lambda \int_{\lambda}^1 \frac{\arccos x}{x} dx - \frac{3}{4} \lambda^3 \arccos \lambda - \sqrt{1 - \lambda^2} + \frac{7}{4} \lambda^2 \sqrt{1 - \lambda^2}.$$

Open is the explicit form of $\tilde{g}_s(\lambda)$ for $s > 3$. A further open question is the explicit form of the a.d.f. of the above sequence with $[0, 1]^s$ instead of $B(r)$.

NOTES: (I) O. Strauch (2003).

(II) Note that the integral in $\tilde{g}_3(\lambda)$ cannot be expressed as a finite combination of elementary functions, cf. I.M. Ryshik and I.S. Gradshteyn (1951, p. 122).

(III) The d.f.'s $\tilde{g}_s(\lambda)$ and $g_s(x)$ from 1.24 form the basis of a new one-time pad cryptosystem introduced in Strauch (2002).

Proposed by O. Strauch.

I.M. RYSHIK – I.S. GRADSTEIN: *Tables of Series, Products, and Integrals*, (German and English dual language edition), VEB Deutscher Verlag der Wissenschaften, Berlin, 1957; translation from the Russian original, Gos. Izd. Tech.–Teor. Lit., Moscow, 1951 (MR 22#3120).

O. STRAUCH: *On distribution functions of sequences generated by scalar and mixed product*, Math. Slovaca **53** (2003), no. 5, 467–478 (MR 2038514).

O. STRAUCH: *Some modification of one-time pad cipher*, Tatra Mt. Math. Publ. **29** (2004), 157–171.

1.26 Ramanujan function.

Let $\tau(n)$ be the Ramanujan function given by

$$q \prod_{n \geq 1} (1 - q^n)^{24} = \sum_{n \geq 1} \tau(n) q^n.$$

Is $\{\tau(n+1)/\tau(n)\}_{n \in \mathbb{N}}$ dense in \mathbb{R} ?

Proposed by F. Luca.

1.27 Apéry sequence.

Let $(A_n)_{n \geq 0}$ be the Apéry sequence given by

$$A_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2.$$

Let $\mathcal{P} = \{p \text{ prime} : p \mid A_n \text{ for some } n\}$.

(i) Is it true that \mathcal{P} misses infinitely many primes?

(ii) For a positive real number x let $\mathcal{P}(x) = \mathcal{P} \cap [1, x]$. Find lower bounds for $\#\mathcal{P}(x)$.

Regarding (ii), it follows from the results of Luca (2007) that $\#\mathcal{P}(x) \gg \log \log x$.

Proposed by F. Luca.

F. LUCA: *Prime divisors of binary holonomic sequences*, Adv. in Appl. Math. **40** (2008), no. 2, 168–179 (MR2388609(2009e:11004)).

1.28 Urban's conjecture.

Let $k \in \mathbb{N}$ be fixed, and let λ_i, μ_i , for $1 \leq i \leq k$ be real algebraic numbers with absolute values greater than 1. Assume that, for $i = 1, 2, \dots, k$, the pairs λ_i, μ_i are multiplicatively independent (i.e. they are not integers m, n such that $\lambda_i^m = \mu_i^n$), and $(\lambda_i, \mu_i) \neq (\lambda_j, \mu_j)$ for $i \neq j$. Then for any real numbers $\theta_1, \dots, \theta_k$ with at least one $\theta_i \notin \mathbb{Q}(\cup_{i=1}^k \{\lambda_i, \mu_i\})$ the double sequence

$$\sum_{i=1}^k \lambda_i^m \mu_i^n \theta_i \pmod{1}, \quad m, n = 1, 2, \dots$$

is dense in $[0, 1]$.

NOTES:

(I) R. Urban (2007). In a first step he proved (Theorem 1.6):

Let λ_1, μ_1 and λ_2, μ_2 be two distinct pairs of multiplicatively independent real algebraic integers of degree 2, with absolute values greater than 1, such that the absolute values of their conjugates $\tilde{\lambda}_1, \tilde{\mu}_1, \tilde{\lambda}_2, \tilde{\mu}_2$ are also greater than 1. Let $\mu_1 = g_1(\lambda_1)$ for some $g_1 \in \mathbb{Z}[x]$ and $\mu_2 = g_2(\lambda_2)$ for some $g_2 \in \mathbb{Z}[x]$. Assume that at least one element in each pair λ_i, μ_i has all positive powers irrational. Assume further that there exist $k, lk', l' \in \mathbb{N}$ such that

(a) $\min(|\lambda_2|^k |\mu_2|^{l'}, |\tilde{\lambda}_2|^k |\tilde{\mu}_2|^{l'}) > \max(|\lambda_1|^k |\mu_1|^l, |\tilde{\lambda}_1|^k |\tilde{\mu}_1|^l)$ and

$$(b) \min(|\lambda_1|^{k'}|\mu_1|^{l'}, |\tilde{\lambda}_1|^{k'}|\tilde{\mu}_1|^{l'}) > \max(|\lambda_2|^{k'}|\mu_2|^{l'}, |\tilde{\lambda}_2|^{k'}|\tilde{\mu}_2|^{l'}).$$

Then for any real numbers θ_1, θ_2 with at least one $\theta_i \neq 0$ the sequence

$$\lambda_1^m \mu_1^n \theta_1 + \lambda_2^m \mu_2^n \theta_2 \pmod{1}, \quad m, n = 1, 2, \dots$$

is dense in $[0, 1]$. For illustration

$$(\sqrt{23} + 1)^m (\sqrt{23} + 2)^n \theta_1 + (\sqrt{61} + 1)^m (\sqrt{61} - 6)^n \theta_2 \pmod{1}, \quad m, n = 1, 2, \dots$$

is dense in $[0, 1]$, assuming $(\theta_1, \theta_2) \neq (0, 0)$.

R. Urban note that (a) and (b) hold, when

$$|\lambda_2| > |\tilde{\lambda}_2| > |\lambda_1| > |\tilde{\lambda}_1| > 1 \text{ and } |\mu_1| > |\tilde{\mu}_1| > |\mu_2| > |\tilde{\mu}_2| > 1.$$

He also note that Theorem 1.6 can be extended in the case when not all of λ_i, μ_i are of degree 2, but if λ_i, μ_i are rationals, then θ_i must be irrational. As example, for every $\theta_2 \neq 0$, the sequence

$$(3 + \sqrt{3})^m 2^n + 5^m 7^n \theta_2 \sqrt{2} \pmod{1}, \quad m, n = 1, 2, \dots$$

is dense in $[0, 1]$.

(II) The Conjecture is motivated by H. Furstenberg's (1967) result: If $p, q > 1$ are multiplicatively independent integers, i.e., they are not both integer powers of some integer, then for every irrational θ the double sequence

$$p^n q^m \theta \pmod{1}, \quad m, n = 1, 2, \dots$$

is everywhere dense in $[0, 1]$.

(III) Further generalization was given by B. Kra (1999): For positive integers $1 < p_i < q_i, i = 1, 2, \dots, k$, assume that all pairs p_i, q_i are multiplicatively independent and pairs $(p_i, q_i) \neq (p_j, q_j)$ for $i \neq j$. Then for distinct $\theta_1, \dots, \theta_k$ with at least one irrational θ_i the sequence

$$\sum_{i=1}^K p_i^n q_i^m \theta_i \pmod{1}, \quad m, n = 1, 2, \dots \quad (1)$$

is dense in $[0, 1]$.

(IV) Berend in MR1487320 (99j:11079) reformulated Kra's result:

Let p_i, q_i integers and θ_i real, $i = 1, 2, \dots, k$. If p_1 and q_1 are multiplicatively independent, θ_1 is irrational, and pairs $(p_i, q_i) \neq (p_1, q_1)$ for $i \geq 2$, then the sequence (1) is dense in $[0, 1]$.

(V) Precisely, Furstenberg (1967) proved: Let S be a non-lacunary semigroup of rational integers. Then $S\alpha \pmod{1}$ is dense in $[0, 1]$ for any irrational α .

D. Berend (1987) extends it:

Let K be a real algebraic number field and S a subsemigroup of the multiplicative group of K such that

- (i) $S \subset (-\infty, -1) \cup (1, \infty)$,
- (ii) there exist multiplicatively independent $\lambda, \mu \in S$ (i.e., there exist no integers m and n , not both of which are 0, with $\lambda^m = \mu^n$),
- (iii) $\mathbb{Q}(S) = K$.

Then for every $\alpha \notin K$ the set $S\alpha \bmod 1$ is dense in $[0, 1]$. If, moreover

- (iv) $S \not\subset PS(K)$,

then $S\alpha \bmod 1$ is dense in $[0, 1]$ for every $\alpha \neq 0$.

Here, if $[K : \mathbb{Q}] = m$ denotes by $PS(K)$ the semigroup of all Pisot or Salem number of degree m over \mathbb{Q} .

Furthermore, if $S\alpha \bmod 1$ is dense in $[0, 1]$ for every $\alpha \notin K$ or for all $\alpha \neq 0$, then S has a subsemigroup having the same property generated by two elements.

(VI) D. Berend (1987a): Let p, q , and c be non-zero integers with p and q multiplicatively independent, ξ an irrational and β arbitrary. Then the set

$$\{p^m q^n \xi + c^{m+n} \beta : m, n \in \mathbb{N}\}$$

is dense modulo 1.

(VII) R. Urban (2009): Let $a_1 > a_2 > 1$ and $b_1 > b_2 > 1$ be two pairs of multiplicatively independent integers, and let c be a positive real number. Suppose that $a_1 < b_1$ and $a_2 > b_2$. Then, for any real numbers ξ_1, ξ_2 with at least one ξ_i irrational, there exists $q \in \mathbb{N}$ such that for any real number β , the set

$$\{a_1^m a_2^n q \xi_1 + b_1^m b_2^n q \xi_2 + c^{m+n} \beta : m, n \in \mathbb{N}\}$$

is dense modulo 1.

Entered by O. Strauch.

[a] D. BEREND: *Actions of sets of integers on irrationals*, Acta Arith. **48** (1987), 275–306.
 D. BEREND: *Dense (mod 1) semigroups of algebraic numbers*, J. Number Theory. **26** (1987), no. 3, 246–256 (MR901238 (88e:11102)).
 H. FURSTENBERG: *Disjointness in ergodic theory, minimal sets, and a problem in Diophantine approximation*, Math. System Theory. **1** (1967), 1–49 (Zbl 0146.28502).
 B. KRA: *A generalization of Furstenberg’s diophantine theorem*, Proc. Amer. Math. Soc. **127** (1999), no. 7, 1951–1956 (MR1487320(99j:11079)).
 R. URBAN: *On density modulo 1 of some expressions containing algebraic integers*, Acta Arith. **127** (2007), no. 3, 217–229.
 R. URBAN: *Multiplicatively independent integers and dense modulo 1 sets of sums*, Uniform Distribution Theory **4** (2009), no. 1, 27–33.

1.29 Extreme values of $\int_0^1 \int_0^1 F(x, y) d_x d_y g(x, y)$ for copulas $g(x, y)$

Let $F(x, y)$ be a Riemann integrable function defined on $[0, 1]^2$ and x_n, y_n , $n = 1, 2, \dots$, be two u.d. sequences in $[0, 1]$. A problem is to find limit

points of the sequence

$$\frac{1}{N} \sum_{n=1}^N F(x_n, y_n), \quad N = 1, 2, \dots \quad (1)$$

Applying Helly theorems we obtain limit points of (1) form the set

$$\left\{ \int_0^1 \int_0^1 F(x, y) d_x d_y g(x, y); g(x, y) \in G((x_n, y_n)) \right\}, \quad (2)$$

where $G(x_n, y_n)$ is the set of all d.f.'s of the two-dimensional sequence (x_n, y_n) , $n = 1, 2, \dots$. In this case, two-dimensional sequence (x_n, y_n) need not be u.d. but every d.f. $g(x, y) \in G((x_n, y_n))$ satisfies

(i) $g(x, 1) = x$ for $x \in [0, 1]$ and

(ii) $g(1, y) = y$ for $y \in [0, 1]$.

The d.f. $g(x, y)$ which satisfies (i) and (ii) is called *copula* and a basic theory of copulas can be found in R.B. Nelsen (1999), see [OP, 2.3, Deterministic analysis of sequences].

Open problem: Find extreme values of $\int_0^1 \int_0^1 F(x, y) d_x d_y g(x, y)$, where $g(x, y)$ is a copula.

NOTES:

(I) Firstly, for $F(x, y) = |x - y|$, this problem was formulated by F. Pillichshammer and S. Steinerberger (2009). They proved: *Let x_n and y_n be two uniformly distributed sequences in $[0, 1]$. Then $\limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} |x_n - y_n| \leq \frac{1}{2}$ and in particular, $\limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} |x_{n+1} - x_n| \leq \frac{1}{2}$ and this result is best possible.* They also found $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} |x_{n+1} - x_n| = \frac{2(b-1)}{b^2}$ for van der Corput sequence x_n in the base b and $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} |x_{n+1} - x_n| = 2\{\alpha\}(1 - \{\alpha\})$ for $x_n = n\alpha \bmod 1$, where α is irrational.

(II) Secondly, S. Steinerberger (2009) study (1) for $F(x, y) = f_1(x)f_2(x)$ and for u.d. sequences $x_n = \Phi(z_n)$ and $y_n = \Psi(z_n)$, where $\Phi(x)$ and $\Psi(x)$ are uniformly distributed preserving (u.d.p.) functions and z_n is a u.d. sequence. For u.d.p. see this [OP, 2.1 Uniform distribution theories]. He proved: Let $f : [0, 1] \rightarrow \mathbb{R}$ be a Lebesgue measurable function, we see it as random variable, and $g(x) = |f^{-1}([0, x])|$ be its d.f. and put $f^*(x) = g^{-1}(x)$. If d.f. $g(x)$ does not have inverse function we put $f^*(x) = \inf\{t \in \mathbb{R}; g(t) \geq x\}$.

Let f_1, f_2 be Riemann integrable functions on $[0, 1]$. Let Φ, Ψ be arbitrary u.d.p. transformations. Then

$$\int_0^1 f_1^*(x) f_2^*(1-x) dx \leq \int_0^1 f_1(\Phi(x)) f_2(\Psi(x)) dx \leq \int_0^1 f_1^*(x) f_2^*(x) dx$$

and these bounds are best possible. Also, every number within the bounds is attained by some u.d.p. Φ, Ψ . In his proof Steinerberger used the Hardy-Littlewood

inequality [Hardy, Littlewood and Pólya (1934), Th. 378] $\int_0^1 f_1(x)f_2(x) \, dx \leq \int_0^1 f_1^*(x)f_2^*(x) \, dx$.

(III) J. Fialová and O. Strauch (2010) proved: *Let $F(x, y)$ be a Riemann integrable function defined on $[0, 1]^2$. For differential of $F(x, y)$ let us assume that $d_x d_y F(x, y) > 0$ for every $(x, y) \in [0, 1]^2$. Then*

$$\begin{aligned} \max_{g(x,y)\text{-copula}} \int_0^1 \int_0^1 F(x, y) \, d_x d_y g(x, y) &= \int_0^1 F(x, x) \, dx, \\ \min_{g(x,y)\text{-copula}} \int_0^1 \int_0^1 F(x, y) \, d_x d_y g(x, y) &= \int_0^1 F(x, 1 - x) \, dx, \end{aligned}$$

where, precisely, max is attained in $g(x, y) = \min(x, y)$ and min in $g(x, y) = \max(x + y - 1, 0)$, uniquely. In proof they used expression

$$\begin{aligned} \int_0^1 \int_0^1 F(x, y) \, d_x d_y g(x, y) &= F(1, 1) - \int_0^1 g(1, y) \, d_y F(1, y) \\ &\quad - \int_0^1 g(x, 1) \, d_x F(x, 1) + \int_0^1 \int_0^1 g(x, y) \, d_x d_y F(x, y) \end{aligned}$$

which holds for every Riemann integrable function $F(x, y)$ and d.f. $g(x, y)$ which have no any common discontinuity points. And then they used Fréchet-Hoeffding bounds [Nelsen (1999), p. 9] $\max(x + y - 1, 0) \leq g(x, y) \leq \min(x, y)$ which holds for every $(x, y) \in [0, 1]^2$ and for every copula $g(x, y)$.

(IIIa) Using Sklar theorem that every d.f. $g(x, y)$ can be express as $g(x, y) = c(g(x, 1), g(1, y))$ for related copula $c(x, y)$ Fialová and Strauch extend: *Let us assume that $F(x, y)$ is a continuous function such that $d_x d_y F(x, y) > 0$ for every $(x, y) \in (0, 1)^2$. Then for the extremes of integral $\int_0^1 \int_0^1 F(x, y) \, d_x d_y g(x, y)$ for $g(x, y)$ for which $g(x, 1) = g_1(x)$ and $g(1, y) = g_2(y)$ we have*

$$\begin{aligned} \max_{g(x,y)} \int_0^1 \int_0^1 F(x, y) \, d_x d_y g(x, y) &= \int_0^1 F(g_1^{-1}(x), g_2^{-1}(x)) \, dx, \\ \min_{g(x,y)} \int_0^1 \int_0^1 F(x, y) \, d_x d_y g(x, y) &= \int_0^1 F(g_1^{-1}(x), g_2^{-1}(1 - x)) \, dx, \end{aligned}$$

where the maximum is attained in $g(x, y) = \min(g_1(x), g_2(y))$ and the minimum in $g(x, y) = \max(g_1(x) + g_2(y) - 1, 0)$, uniquely.

(IIIb) Fialová and Strauch (2011) criterion: *Assume that a copula $g(x, y)$ maximize $\int_0^1 \int_0^1 F(x, y) \, d_x d_y g(x, y)$ and let $[X_1, X_2] \times [Y_1, Y_2]$ be an interval in $[0, 1]^2$ such that the differential*

$$g(X_2, Y_2) + g(X_1, Y_1) - g(X_2, Y_1) - g(X_1, Y_2) > 0.$$

If for every interior point (x, y) of $[X_1, X_2] \times [Y_1, Y_2]$ the differential $d_x d_y F(x, y)$ has a constant sign, then

(i) if $d_x d_y F(x, y) > 0$ then

$$g(x, y) = \min(g(x, Y_2) + g(X_1, y) - g(X_1, Y_2), g(x, Y_1) + g(X_2, y) - g(X_2, Y_1))$$

(ii) if $d_x d_y F(x, y) < 0$ then

$$g(x, y) = \max(g(x, Y_2) + g(X_2, y) - g(X_2, Y_2), g(x, Y_1) + g(X_1, y) - g(X_1, Y_1))$$


for every $(x, y) \in [X_1, X_2] \times [Y_1, Y_2]$.

(IV) Fialová and Strauch (2010) also consider $F(x, y)$ in the form $F(x, y) = f(x)y$ and study the limit points of $\frac{1}{N} \sum_{n=1}^N f(x_n)y_n$, where x_n is u.d. sequence and u.d. sequence y_n is given by $y_n = \Phi(x_n)$, where $\Phi(x)$ is a u.d.p. This problem is equivalent to find

$$\max_{\Phi(x) \text{--u.d.p.}} \int_0^1 f(x)\Phi(x) dx, \quad \min_{\Phi(x) \text{--u.d.p.}} \int_0^1 f(x)\Phi(x) dx.$$

Fialová (2010) solve this problem for piecewise linear $f(x)$. Here the function $f : [0, 1] \rightarrow [0, 1]$ is *piecewise linear* (p.l.) if there exists a system of ordinate intervals $J_j, j = 1, 2, \dots, u$ which are disjoint and fulfilled the whole interval $[0, 1]$, and a corresponding system of abscissa intervals $I_{j,i}, i = 1, 2, \dots, l_j$, such that $f(x)/I_{j,i}$ is the increasing or decreasing diagonal of $I_{j,i} \times J_j$. Fialová (2010) proved: *Let $f : [0, 1] \rightarrow [0, 1]$ be a p.l. function with ordinate decomposition $J_j, j = 1, 2, \dots, n$, and abscissa decomposition $I_{j,i}, i = 1, 2, \dots, l_j$. Define a p.l. function $\Psi(x)$ in the same abscissa decomposition $I_{j,i}$, but in a new ordinate decomposition J'_j , with the lengths $|J'_j| = \sum_{i=1}^{l_j} |I_{j,i}|, j = 1, 2, \dots, n$, ordered similarly as $J_j, j = 1, 2, \dots, u$. Put the graph of $\Psi(x)/I_{j,i}$ on $I_{j,i} \times J'_j$ as the diagonal \nearrow or \searrow if and only if $f(x)/I_{j,i}$ is the \nearrow or \searrow diagonal. Note that if $f(x)$ is a constant on the interval $I_{j,i}$, then J_i is a point, and the graph $\Psi(x)/I_{j,i}$ can be defined arbitrary, either increasing or decreasing in $I_{j,i} \times J'_j$. Then $\Psi(x)$ is the best u.d.p. approximation of $f(x)$.*

For example



(IVa) The result in (IV) correspond (II) since we have $\Psi(x) = g_f(f(x))$. But Fialová used $\int_0^1 (f(x) - \Psi(x))^2 \, dx = \int_0^1 f^2(x) \, dx - 2 \int_0^1 f(x)\Psi(x) \, dx + \int_0^1 \Psi^2(x) \, dx$, $\int_0^1 \Psi^2(x) \, dx = \frac{1}{3}$ which gives

$$\max \int_0^1 f(x)\Psi(x) \, dx = \min \int_0^1 (f(x) - \Psi(x))^2 \, dx.$$

(V) Steinerberger (2009) generalized **open problem** to give bounds for

$$\int_0^1 f_1(\Phi_1(x))f_2(\Phi_2(x)) \dots f_s(\Phi_s(x)) \, dx$$

of Riemann integrable f_1, \dots, f_s and u.d.p. maps Φ_1, \dots, Φ_s . He proved the following partial results:

a) $\max_{\Phi_1, \dots, \Phi_s} \int_0^1 f_1(\Phi_1(x))f_2(\Phi_2(x)) \dots f_s(\Phi_s(x)) \, dx$

$$\leq \left(\prod_{i=1}^n \int_0^1 |f_i(x)|^s \, dx \right)^{\frac{1}{s}}.$$

b) $\min_{\Phi_1, \dots, \Phi_n} \int_0^1 \Phi_1(x)\Phi_2(x) \dots \Phi_s(x) \, dx \geq \frac{1}{e^s}.$

c) $\min_{\Phi_1, \dots, \Phi_s} \int_0^1 \Phi_1(x)\Phi_2(x) \dots \Phi_s(x) \, dx \leq e^{\frac{1}{6s} \frac{s}{s-2} \frac{4}{\pi} \frac{1}{e^s}}.$

Comments: Let $x_n, n = 1, 2, \dots$, be a u.d. sequence in $[0, 1)$ and $g(t_1, \dots, t_s)$ be an a.d.f. of the s -dimensional sequence $(\Phi_1(x_n), \dots, \Phi_s(x_n)), n = 1, 2, \dots$

We have

$$g(t_1, \dots, t_s) = |\Phi_1^{-1}([0, t_1]) \cap \dots \cap \Phi_s^{-1}([0, t_s])|,$$

$g(1 \dots, t_i, 1 \dots, 1) = t_i$ for $i = 1, \dots, s$, i.e., it is a copula and

$$\int_0^1 f_1(\Phi_1(x)) \dots f_s(\Phi_s(x)) \, dx = \int_{[0,1]^s} f_1(t_1) \dots f_s(t_s) \, dg(t_1, \dots, t_s).$$

(VI) Thus we arrive at the **open problem**: Find extreme values of $\int_{[0,1]^s} F(\mathbf{x}) \, dg(\mathbf{x})$, where $g(\mathbf{x})$ is an s -dimensional copula.

(VII) Steinerberger (2010) generalized (I) to give bounds for the asymptotic behavior of

$$\frac{1}{N} \sum_{n=1}^N \|\mathbf{x}_n - \mathbf{y}_n\|,$$

where $\mathbf{x}_n, \mathbf{y}_n$ are u.d. sequences in a bounded Jordan measurable domain Ω . E.g., for s -dimensional ball he found the sharp inequality $\frac{1}{N} \sum_{n=1}^N \|\mathbf{x}_n - \mathbf{y}_n\| \leq \frac{2s}{s+1}$.

(VIII) **Open problem:** Transform the theory of d.f.'s to the multidimensional unit sphere \mathbb{S} and find extremes of the energy integral

$$\int_{\mathbb{S}} \|\mathbf{x} - \mathbf{y}\|^s dg(\mathbf{x}) dg(\mathbf{y}).$$

The expository paper on Riesz energy can be found in J. Brauchart (2011).
(IX) See also problem 1.37.

Entered by O. Strauch.

J.S. BRAUCHART: *Optimal discrete Riesz energy and discrepancy*, Uniform Distribution Theory **6** (2011), no. 2, 207–220.

G.H. HARDY – E. LITTLEWOOD – G. PÓLYA: *Inequalities*, Cambridge University Press, London, 1934.

R.B. NELSEN: *An Introduction to Copulas. Properties and Applications*, Lecture Notes in Statistics **139**, Springer-Verlag, New York, 1999.

F. PILLICHSHAMMER – S. STEINERBERGER: *Average distance between consecutive points of uniformly distributed sequences*, Uniform Distribution Theory **4** (2009), no. 1, 51–67.

S. STEINERBERGER: *Uniform distribution preserving mappings and variational problems*, Uniform Distribution Theory **4** (2009), no. 1, 117–145.

S. STEINERBERGER: *Extremal uniform distribution and random chords*, Acta Math. Hungar. **130** (2011), no. 4, 321–339 (MR2771097 (2012a:11106)).

J. FIALOVÁ – O. STRAUCH: *On two-dimensional sequences composed of one-dimensional uniformly distributed sequences*, Uniform Distribution Theory **6** (2011), no. 1, 101–125.

1.30 Niederreiter-Halton (NH) sequence

Directly from R. Hofer and G. Larcher (2010): Niederreiter-Halton (NH) sequence is a combination of different digital (\mathbf{T}_i, w_i) -sequences in different prime bases q_1, \dots, q_r with $w_1 + \dots + w_r = s$ into a single sequence in $[0, 1)^s$.

Finite row NH sequence is a NH sequence if all generating matrices of the component digital (\mathbf{T}_i, w_i) -sequences have each row containing only finitely many entries different from zero.

Infinite row NH sequence is a (NH) sequence which is not finite row.

Digital (\mathbf{T}, s) -sequence over \mathbb{F}_q .

- Let s be a dimension;
- q be a prime;
- Represent $n = n_0 + n_1q + n_2q^2 + \dots$ in base q ;
- Let C_1, \dots, C_s be $\mathbb{N} \times \mathbb{N}$ -matrices in the finite field \mathbb{F}_q ;
- $C_i \cdot (n_0, n_1, \dots)^T = (y_0^{(i)}, y_1^{(i)}, \dots)^T \in \mathbb{F}_q^{\mathbb{N}}$;
- $x_n^{(i)} := \frac{y_0^{(i)}}{q} + \frac{y_1^{(i)}}{q^2} + \dots$;

- The sequence $\mathbf{x}_n = (x_n^{(1)}, \dots, x_n^{(s)})$ is said to be (\mathbf{T}, s) -sequence if for every $m \in \mathbb{N}$ there exists $\mathbf{T}(m)$ such that $0 \leq \mathbf{T}(m) \leq m$ and for all $d_1 + \dots + d_s = m - \mathbf{T}(m)$ and the $(m - \mathbf{T}(m)) \times m$ -matrix consisting of the upper left $d_1 \times m$ -submatrix of C_1
the upper left $d_2 \times m$ -submatrix of C_2
...
the upper left $d_s \times m$ -submatrix of C_s
has rank $m - \mathbf{T}(m)$.

If \mathbf{T} is minimal we speak **strict digital (\mathbf{T}, s) -sequence**.

Open Problem: Determine whether the following two-dimensional NH sequences in base 3 and respectively 2 are low-discrepancy sequences (i.e. $D_N^* = O((\log N)^s)/N$) or not:

1. $C^{(1)}$ is the unit matrix in \mathbb{F}_3 and

$$C^{(2)} = \begin{pmatrix} 1 & \underbrace{00\dots 0}_{l_1} & 1 & & 0 & 0 & \dots \\ 0 & 1 & \underbrace{00\dots\dots 0}_{l_2} & & 1 & 0 & 0 & \dots \\ 0 & 0 & 1 & & \underbrace{00\dots\dots\dots 0}_{l_3} & 1 & 0 & 0 & \dots \\ \dots & \dots \\ \dots & \dots \end{pmatrix}$$

in \mathbb{F}_2 with l_1, l_2, l_3, \dots arbitrary but $\lim_{n \rightarrow \infty} l_n = \infty$.

2. $C^{(1)}$ is the unit matrix in \mathbb{F}_3 and

$$C^{(2)} = \begin{pmatrix} 1 & \underbrace{00\dots 0}_{l_1} & 1 & \underbrace{00\dots 0}_{l_2} & 1 & 0 & \dots \\ 0 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 1 & 0 & 0 & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \end{pmatrix}$$

in \mathbb{F}_2 and the first row contains infinitely many 1's but with density 0.

3.

$$C^{(1)} = C^{(2)} = \begin{pmatrix} 1 & 1 & 1 & 1 & \dots \\ 0 & 1 & 0 & 0 & \dots \\ 0 & 0 & 1 & 0 & \dots \\ 0 & 0 & 1 & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix}$$

but $C^{(1)}$ in \mathbb{F}_3 and $C^{(2)}$ in \mathbb{F}_2 .

NOTES:

(I) A basic example is the Halton sequence which is a combination of s digital $(0, 1)$ -sequences in different prime bases q_1, \dots, q_s generated by the unit matrices in \mathbb{F}_{q_i} for each i .

(II) General NH sequences were first investigated by R. Hofer, P. Kritzer, G. Larcher and F. Pillichshammer (2009) and R. Hofer (2009) and she proved: NH sequence is u.d. if and only if each (\mathbf{T}_i, w_i) is u.d.

(III) A strictly digital (\mathbf{T}, s) -sequence is u.d. if and only if $\lim_{m \rightarrow \infty} (m - \mathbf{T}(m)) = \infty$. If $\mathbf{T}(m) \leq t$ for all m , then (\mathbf{T}, s) -sequence is (t, s) -sequence.

(IV) R. Hofer and G. Larcher (2010) give concrete examples of digital $(0, s)$ -sequences generated by matrices with finite rows.

R. HOFER: *On the distribution properties of Niederreiter-Halton sequences*, J. Number Theory **129** (2009), 451–463.

R. HOFER – P. KRITZER – G. LARCHER – F. PILLICHSHAMMER: *Distribution properties of generalized van der Corput-Halton sequences and their subsequences*, Int. J. Number Theory **5** (2009), 719–746.

R. HOFER – G. LARCHER: *On existence and discrepancy of certain digital Niederreiter-Halton sequences*, Acta Arith. **141** (2010), no. 4, 369–394.

1.31 Gauss-Kuzmin theorem and $g(x) = g_f(x)$

Denote

$$\begin{aligned} f(x) &= 1/x \bmod 1, \\ g_f(x) &= \int_{f^{-1}([0,x])} 1. \, dg(x), \\ (g^{f^n})_f(x) &= g_{f^{n+1}}(x), \\ g_0(x) &= \frac{\log(1+x)}{\log 2}. \end{aligned}$$

The problem is to find all solutions $g(x)$ of the functional equation $g(x) = g_f(x)$ for $x \in [0, 1]$. It is equivalent to

$$g(x) = \sum_{n=1}^{\infty} g\left(\frac{1}{n}\right) - g\left(\frac{1}{n+x}\right) \text{ for d.f. } g(x), x \in [0, 1]. \quad (1)$$

The following is known:

- (I) $g_0(x)$ satisfies (1).
- (II) Gauss-Kuzmin theorem: If $g(x) = x$, then $g_{f^n}(x) \rightarrow g_0(x)$ and the rate of convergence is $O(q^{\sqrt{n}})$, $0 < q < 1$.
- (III) Theorem in (II) was proved by R. Kuzmin (1928) assuming for a starting function $g(x)$
 - (i) $0 < g'(x) < M$ and
 - (ii) $|g''(x)| < \mu$.

Thus, if $g(x)$ satisfies (i), (ii), and (1), then $g(x) = g_0(x)$.

- (IV) Theorem (II) was inspired by Gauss. He conjectured $m_n(x) \rightarrow g_0(x)$, where $m_n(x) = |\{\alpha \in [0, 1]; 1/r_n(\alpha) < x\}|$ and for continued fraction expansion $\alpha = [a_0(\alpha); a_1(\alpha), a_2(\alpha), \dots]$, $r_n(\alpha) = [a_{n+1}(\alpha); a_{n+2}(\alpha), \dots]$. In this case $m_n(x) = g_{f^n}(x)$ for $g(x) = x$, since $f(1/r_n(\alpha)) = 1/r_{n+1}(\alpha)$.
- (V) For starting point $x_0 \in [0, 1]$ we define the iterate sequence x_n as

$$x_1 = f(x_0), x_2 = f(f(x_0)), x_3 = f(f(f(x_0))), \dots$$

Then a.d.f. $g(x)$ solves (1). For example, the sequence $x_1 = 1/r_1, x_2 = 1/r_2, \dots$ for $\frac{\sqrt{5}-1}{2} = [0; 1, 1, 1, \dots]$ produces solution $g(x) = c_{\frac{\sqrt{5}-1}{2}}(x)$.

- (VI) **Chain of solutions.** If d.f. $g_1(x)$ solve the equation $g_f(x) = g(x)$ and $(g_2)_f(x) = g_1(x)$, then $g_2(x)$ solve $g_f(x) = g(x)$, again. The $g_2(x)$ can be found as solution

$$g_1(x) = \sum_{n=1}^{\infty} g_2\left(\frac{1}{n}\right) - g_2\left(\frac{1}{n+x}\right).$$

From it

$$g_2\left(\frac{1}{x+1}\right) = 1 - g_1(x) + \sum_{n=2}^{\infty} g_2\left(\frac{1}{n}\right) - g_2\left(\frac{1}{n+x}\right)$$

and thus it is suffice to define a non-increasing $g_2(x)$ on $[0, 1/2)$, such that

- (i) $g_2(0) = 0$,
- (ii) $\sum_{n=2}^{\infty} g_2\left(\frac{1}{n}\right) - g_2\left(\frac{1}{n+x}\right) \leq \sum_{n=1}^{\infty} g_1\left(\frac{1}{n}\right) - g_1\left(\frac{1}{n+x}\right)$,
- (iii) $\sum_{n=2}^{\infty} g_2'\left(\frac{1}{n}\right) \frac{1}{(x+n)^2} \leq \sum_{n=1}^{\infty} g_1'\left(\frac{1}{n}\right) \frac{1}{(x+n)^2}$.

- (VII) If d.f. $g_1(x)$ and $g_2(x)$ satisfy (1) and $g_1(x) = g_2(x)$ for $x \in [0, 1/2)$, then $g_1(x)$ and $g_2(x)$ coincide on the whole interval $[0, 1]$. Other sets of uniqueness are $[0, \frac{1}{n+1}) \cup (\frac{1}{n}, 1]$ for arbitrary positive integer n .
- (VIII) If $x_n \in [0, 1)$ has a.d.f. $g_1(x)$, then $\{\frac{1}{x_n}\}$ has a.d.f.

$$g_2(x) = \sum_{n=1}^{\infty} g_1\left(\frac{1}{n}\right) - g_1\left(\frac{1}{n+x}\right).$$

Entered by O. Strauch.

R.O. KUZMIN: *A problem of Gauss*, Dokl. Akad. Nauk, Ser. A, 375–380, 1928.

1.32 Benford law

Let $b \geq 2$ be an integer considered as a base for the development of positive real number $x > 0$ and $M_b(x)$ be a mantissa of x defined by $x = M_b(x) \times b^{n(x)}$ such that $1 \leq M_b(x) < b$ holds, where $n(x)$ is a uniquely determined integer. Let $K = k_1 k_2 \dots k_r$ be a positive integer expressed in the base b , that is $K = k_1 \times b^{r-1} + k_2 \times b^{r-2} + \dots + k_{r-1} \times b + k_r$, where $k_1 \neq 0$ and at the same time $K = k_1 k_2 \dots k_r$ is considered as an r -consecutive block of integers in the base b . We have

$$\begin{aligned} K \leq M_b(x) \times b^{r-1} < K + 1 &\iff \\ \frac{K}{b^{r-1}} \leq M_b(x) < \frac{K + 1}{b^{r-1}} &\iff \\ \log_b \left(\frac{K}{b^{r-1}} \right) \leq \log_b(M_b(x)) < \log_b \left(\frac{K + 1}{b^{r-1}} \right) &\iff \\ \log_b \left(\frac{K}{b^{r-1}} \right) \leq \log_b x \bmod 1 < \log_b \left(\frac{K + 1}{b^{r-1}} \right). &\quad (1) \end{aligned}$$

• Definition: A sequence x_n , $n = 1, 2, \dots$, of positive real numbers satisfies Benford law (abbreviated to B.L.) of order r if for every r -digits number $K = k_1 k_2 \dots k_r$ we have

$$\lim_{N \rightarrow \infty} \frac{\#\{n \leq N; \text{ first } r \text{ digits of } M_b(x_n) \text{ are equal to } K\}}{N} = \log_b(K+1) - \log_b K.$$

Here "the first r digits of $M_b(x_n) = K$ is the same as "the first r digits (starting a non-zero digit) of $x_n = K$.

• Definition: If a sequence x_n , $n = 1, 2, \dots$, satisfies B.L. of order r , for every $r = 1, 2, \dots$, then it is called that x_n satisfies strong B.L. or extended or generalized B.L. In the following we will described it as B.L.

From (1) directly follows:

(I) Theorem: A sequence x_n , $x_n > 0$, $n = 1, 2, \dots$, satisfies B.L. if and only if the sequence $\log_b x_n \bmod 1$ is u.d. in $[0, 1)$.

(II) Theorem: For every K and r there exists infinitely many n such that the first r digits (starting a non-zero digit) of $x_n = K$ if and only if $\log_b x_n \bmod 1$ is dense in $[0, 1)$.

Characterization u.d. of $\log_b x_n \bmod 1$ using d.f.'s in $G(x_n \bmod 1)$

In V. Baláž, K. Nagasaka and O. Strauch (2010) is proved:

(III) Theorem: Let x_n , $n = 1, 2, \dots$, be a sequence in $(0, 1)$ and $G(x_n)$ be the set of all d.f.s of x_n . Assume that every d.f. $g(x) \in G(x_n)$ is continuous at $x = 0$. Then the sequence x_n satisfies B.L. in the base b if and only if for every $g(x) \in G(x_n)$ we have

$$x = \sum_{i=0}^{\infty} \left(g\left(\frac{1}{b^i}\right) - g\left(\frac{1}{b^{i+x}}\right) \right) \text{ for } x \in [0, 1]. \quad (2)$$

Find all solutions of (2). Some examples are:

$$g(x) = \begin{cases} x & \text{if } x \in [0, \frac{1}{b}], \\ 1 + \log_b x + (1-x)\frac{1}{b-1} & \text{if } x \in [\frac{1}{b}, 1]. \end{cases}$$

$$\tilde{g}(x) = \begin{cases} 0 & \text{if } x \in [0, \frac{1}{b}], \\ 1 + \frac{\log x}{\log b} & \text{if } x \in [\frac{1}{b}, 1], \end{cases}$$

$$g^*(x) = \begin{cases} 0 & \text{if } x \in [0, \frac{1}{b^2}], \\ 2 + \frac{\log x}{\log b} & \text{if } x \in [\frac{1}{b^2}, \frac{1}{b}], \\ 1 & \text{if } x \in [\frac{1}{b}, 1] \end{cases}$$

$$g^{**}(x) = \begin{cases} 0 & \text{if } x \in [0, \frac{1}{b^3}], \\ 3 + \frac{\log x}{\log b} & \text{if } x \in [\frac{1}{b^3}, \frac{1}{b^2}], \\ 1 & \text{if } x \in [\frac{1}{b^2}, 1]. \end{cases}$$

(IV) Simple results:

(i) Fibonacci numbers F_n , $n!$, n^n , n^{n^2} , satisfy B.L.

(ii) The positive sequences x_n and $1/x_n$, $n = 1, 2, \dots$ satisfy B.L. in the base b simultaneously.

(iii) The positive sequences x_n and nx_n , $n = 1, 2, \dots$ satisfy B.L. in the base b simultaneously.

(iv) For a sequence $x_n > 0$, $n = 1, 2, \dots$, assume that

- $\lim_{n \rightarrow \infty} x_n = \infty$ monotonically,
- $\lim_{n \rightarrow \infty} \log \frac{x_{n+1}}{x_n} = 0$ monotonically.

Then the sequence x_n satisfies B.L. in every base b if and only if

$$\lim_{n \rightarrow \infty} n \log \frac{x_{n+1}}{x_n} = \infty.$$

(v) Assume $x_n > 0$, $n = 1, 2, \dots$. If for every $k = 1, 2, \dots$ the ratio sequence x_{n+k}/x_n , $n = 1, 2, \dots$, satisfies B.L. in the base b , then the original sequence x_n , $n = 1, 2, \dots$ also satisfies B.L. in the base b , see A.I. Pavlov (1981).

(IX) J.L. Brown, Jr. and R.L. Duncan (1970): Let x_n be a sequence generated by the recursion relation

$$x_{n+k} = a_{k-1}x_{n+k-1} + \dots + a_1x_{n+1} + a_0x_n, \quad n = 1, 2, \dots,$$

where a_0, a_1, \dots, a_{k-1} are non-negative rationals with $a_0 \neq 0$, k is a fixed integer, and x_1, x_2, \dots, x_k are starting points. Assume that the characteristic polynomial

$$x^k - a_{k-1}x^{k-1} - \dots - a_1x - a_0$$

has k distinct roots $\beta_1, \beta_2, \dots, \beta_k$ satisfying

$$0 < |\beta_1| < \dots < |\beta_k|$$

and such that none of the roots has magnitude equal to 1.

Then $\log x_n \pmod 1$ is u.d.

Furthermore, the general solution of the recurrence is $x_n = \sum_{j=1}^k \alpha_j \beta_j^n$ and if j_0 is the largest value of j for which $\alpha_j \neq 0$ and if $\log_b \beta_{j_0}$ is irrational, then also

$\log_b x_n \pmod 1$ is u.d.,

i.e. x_n satisfies B.L. in the base b . This implies that Fibonacci and Lucas numbers obey B.L. what rediscovered L.C. Washington (1981).

Entered by O. Strauch.

V. BALÁŽ – K. NAGASAKA – O. STRAUCH: *Benford's law and distribution functions of sequences in (0, 1)* (Russian), *Mat. Zametki* **88** (2010), no. 4, 485–501; (English translation) *Math. Notes* **88** (2010), no. 4, 449–463.

J.L. BROWN, JR. – R.L. DUNCAN: *Modulo one uniform distribution of the sequence of logarithms of certain recursive sequences*, *Fibonacci Quart.* **8** (1970), 482–486.

F. LUCA – P. STANICA: *On the first digits of the Fibonacci numbers and their Euler function*, *Uniform Distribution Theory*, accepted 2013.

A.I. PAVLOV: *On distribution of fractional parts and Benford's law*, (Russian), *Izv. Aka. Nauk SSSR Ser. Mat.* **45** (1981), no. 4, 760–774.

L.C. WASHINGTON: *Benford's law for Fibonacci and Lucas numbers*, *Fibonacci Quart.* **19** (1981), 175–177.

1.33 The integral $\int_{[0,1]^s} F(\mathbf{x}, \mathbf{y}) dg(\mathbf{x}) dg(\mathbf{y})$ for d.f.'s $g(\mathbf{x})$

• Let $F(x, y)$ be a real continuous symmetric function defined on $[0, 1]^2$ and let $G(F)$ be a set of all d.f.s $g(x)$ satisfying

$$\int_0^1 \int_0^1 F(x, y) dg(x) dg(y) = 0. \quad (1)$$

The study of $G(F)$ is motivated by the fact that for every sequence $x_n \in [0, 1)$ we have

$$G(x_n) \subset G(F) \iff \lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{m,n=1}^N F(x_m, x_n) = 0,$$

where $G(x_n)$ is the set of all d.f.s of the sequence $x_n, n = 1, 2, \dots$ ¹

This immediately follows from Riemann-Stieltjes integral

$$\int_0^1 \int_0^1 F(x, y) dF_N(x) dF_N(y) = \frac{1}{N^2} \sum_{m,n=1}^N F(x_m, x_n),$$

where $F_N(x) = \frac{1}{N} \# \{n \leq N; x_n < x\}$. Assuming $\lim_{k \rightarrow \infty} F_{N_k}(x) = g(x)$ for all continuity points x of g , then Helly-Bray lemma implies

$$\lim_{k \rightarrow \infty} \int_0^1 \int_0^1 F(x, y) dF_{N_k}(x) dF_{N_k}(y) = \int_0^1 \int_0^1 F(x, y) dg(x) dg(y).$$

Open problem is to solve $\int_0^1 \int_0^1 F(x, y) dg(x) dg(y) = 0$ in d.f.s $g(x)$. The multi-dimensional case is mentioned in Problem 2.2 (II).

Partial results:

(I) Let us denote

$$F_{\tilde{g}}(x, y) = \int_0^1 \tilde{g}^2(t) dt - \int_x^1 \tilde{g}(t) dt - \int_y^1 \tilde{g}(t) dt + 1 - \max(x, y).$$

From the relation

$$\int_0^1 (g(x) - \tilde{g}(x))^2 dx = \int_0^1 \int_0^1 F_{\tilde{g}}(x, y) dg(x) dg(y),$$

¹Define $G(F = A)$ is the set of all d.f.s $g(x)$ for which $\int_0^1 \int_0^1 F(x, y) dg(x) dg(y) = A$, and $G(A \leq F \leq B)$ is the set of all d.f.s $g(x)$ for which $A \leq \int_0^1 \int_0^1 F(x, y) dg(x) dg(y) \leq B$. Then again $G(x_n) \subset G(F = A)$ is equivalent to $\lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{m,n=1}^N F(x_m, x_n) = A$, and $G(x_n) \subset G(A \leq F \leq B)$ is equivalent to $A \leq \liminf_{N \rightarrow \infty} \frac{1}{N^2} \sum_{m,n=1}^N F(x_m, x_n)$ and $\limsup_{N \rightarrow \infty} \frac{1}{N^2} \sum_{m,n=1}^N F(x_m, x_n) \leq B$.

we see that the moment problem (1) with $F(x, y) = F_{\tilde{g}}(x, y)$ has the unique solution $g(x) = \tilde{g}(x)$.

(II) Let $F : [0, 1]^2 \rightarrow \mathbb{R}$ be a continuous and symmetric function. For every distribution functions $g(x), \tilde{g}(x)$ we have

$$\begin{aligned} \int_0^1 \int_0^1 F(x, y) dg(x) dg(y) = 0 &\iff \int_0^1 \int_0^1 F(x, y) d\tilde{g}(x) d\tilde{g}(y) \\ &= \int_0^1 (g(x) - \tilde{g}(x)) \left(2 d_x F(x, 1) - \int_0^1 (g(y) + \tilde{g}(y)) d_y d_x F(x, y) \right). \end{aligned}$$

Especially, putting $\tilde{g}(x) = c_0(x)$, we have

$$\begin{aligned} \int_0^1 \int_0^1 F(x, y) dg(x) dg(y) = 0 &\iff \\ F(0, 0) = \int_0^1 (g(x) - 1) \left(2 d_x F(x, 1) - \int_0^1 (g(y) + 1) d_y d_x F(x, y) \right). \end{aligned}$$

• A symmetric continuous $F(x, y)$ defined on $[0, 1]^2$ is called *copositive* if

$$\int_0^1 \int_0^1 F(x, y) dg(x) dg(y) \geq 0$$

for all distribution functions $g : [0, 1] \rightarrow [0, 1]$.

(III) Let $F(x, y)$ be a copositive function having continuous $F'_x(x, 1)$ a.e., and let $g_1(x)$ be a strictly increasing solution of the moment problem (1). Then for every strictly increasing d.f. $g(x)$ we have

$$\int_0^1 \int_0^1 F(x, y) dg(x) dg(y) = 0 \iff F'_x(x, 1) = \int_0^1 g(y) d_y F'_x(x, y) \text{ a.e. on } [0, 1],$$

Proposed by O. Strauch.

O. STRAUCH: *On set of distribution functions of a sequence*, in: *Proc. Conf. Analytic and Elementary Number Theory*, In Honor of E. Hlawka's 80th Birthday, Vienna, July 18-20, 1996, Universität Wien and Universität für Bodenkultur, (W. G. Nowak and J. Schoißengeier, eds.) Vienna, 1997, 214–229 (Zbl 886.11044).

O. STRAUCH: *Moment problem of the type $\int_0^1 \int_0^1 F(x, y) dg(x) dg(y) = 0$* , in: *Proceedings of the International Conference on Algebraic Number Theory and Diophantine Analysis held in Graz*, August 30 to September 5, 1998, Eds. F. Halter-Koch, R.F. Tichy, Walter de Gruyter, Berlin, New York, 2000, 423–443 (MR 2001d:11079).

1.34 Comparison of random sequences using the game theory

A finite two-person zero-sum matrix game with the payoff matrix \mathbf{A} .

$$\mathbf{A} = \begin{pmatrix} a_{1,1} & a_{1,2} & \dots & a_{1,m} \\ a_{1,1} & a_{1,2} & \dots & a_{1,m} \\ \dots & \dots & \dots & \dots \\ a_{m,1} & a_{m,2} & \dots & a_{m,m} \end{pmatrix}$$

In this form, Player I chooses a row, Player II chooses a column, and II pays I the entry in the chosen row and column. Note that the entries of the matrix are the winnings of the row chooser and losses of the column chooser. This pure strategy for Player I of choosing row i may be represented as the $\mathbf{e}_i = (0, \dots, 0, 1, 0, \dots, 0)$, the unit vector with a 1 in the i -th position and 0's elsewhere. Similarly, the pure strategy for II of choosing the j -th column may be represented by $\mathbf{e}_j = (0, \dots, 0, 1, 0, \dots, 0)$ and the payoff to I is

$$\mathbf{e}_i \mathbf{A} \mathbf{e}_j^T = a_{i,j}.$$

Now, let Player I use a sequence $\mathbf{e}_n^{(I)}$, $n = 1, 2, \dots$, of pure strategy and Player II a sequence $\mathbf{e}_n^{(II)}$, $n = 1, 2, \dots$, of pure strategy. Then the mean-value of the payoff of I after N games is

$$\frac{1}{N} \sum_{n=1}^N \mathbf{e}_n^{(I)} \mathbf{A} (\mathbf{e}_n^{(II)})^T.$$

Assume that there exist densities

$$\lim_{N \rightarrow \infty} \frac{1}{N} \#\{n \leq N; \mathbf{e}_n^{(I)} = \mathbf{e}_i\} = p_i, \quad i = 1, 2, \dots, m$$

$$\lim_{N \rightarrow \infty} \frac{1}{N} \#\{n \leq N; \mathbf{e}_n^{(II)} = \mathbf{e}_i\} = q_i, \quad i = 1, 2, \dots, m$$

The vector $\mathbf{p} = (p_1, p_2, \dots, p_m)$ is called a mixed strategy for Player I. Similarly $\mathbf{q} = (q_1, q_2, \dots, q_m)$ is a mixed strategy for Player II. If the sequences $\mathbf{e}_n^{(I)}$ and $\mathbf{e}_n^{(II)}$ are statistically independent, then we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mathbf{e}_n^{(I)} \mathbf{A} (\mathbf{e}_n^{(II)})^T = \mathbf{p} \mathbf{A} \mathbf{q}^T = \sum_{i,j=1}^m p_i a_{i,j} q_j. \tag{1}$$

The mixed strategies \mathbf{p} and \mathbf{q} can be computed optimally such that $\mathbf{p} \mathbf{A} \mathbf{q}^T = 0$, but independence of $\mathbf{e}_n^{(I)}$ and $\mathbf{e}_n^{(II)}$ is a **problem**. Player with better sequence can be found payoff positive.

Now, we transform the above matrix game to continuous case: Put

$$I_{i,j} = [p_1+p_2+\cdots+p_{i-1}, p_1+p_2+\cdots+p_i) \times [q_1+q_2+\cdots+q_{j-1}, q_1+q_2+\cdots+q_j).$$

Define $F(x, y)$ on $[0, 1]^2$ such that

$$F(x, y) = a_{i,j} \text{ if } (x, y) \in I_{i,j}, i, j, = 1, 2, \dots, m. \quad (1)$$

Let Player I use u.d. sequence x_n and Player II u.d. sequence y_n , $n = 1, 2, \dots$. If $(x_n, y_n) \in I_{i,j}$ the Player I chooses pure strategy \mathbf{e}_i and Player II pure strategy \mathbf{e}_j . Then mean value of the payoff of the Player I is

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N F(x_n, y_n) = \int_0^1 \int_0^1 F(x, y) dg(x, y), \quad (2)$$

where $g(x, y)$ is a d.f. of the sequence (x_n, y_n) , $n = 1, 2, \dots$.

Example: Odd or Even. Players I and II simultaneously call out one of the numbers one or two. Player I wins if the sum of the numbers is odd. Player II wins if the sum of the numbers is even. The payoff matrix \mathbf{A} is

$$\mathbf{A} = \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix}$$

and the function $F(x, y)$ corresponding to (1) is

		$F(x, y)$	
		1	-1
1	-1	1	-1
-1	1	-1	1

Let x_n and y_n , $n = 1, 2, \dots$ be u.d. sequences such that (x_n, y_n) has a.d.f. $g(x, y)$.

(i) If $g(x, y) = xy$ then $\int_0^1 \int_0^1 F(x, y) dx dy = 0$.

(ii) If $g(x, y) = \min(x, y)$ then $\int_0^1 \int_0^1 F(x, y) d\min(x, y) = \int_0^1 F(x, x) dx = -1$.

(iii) If $g(x, y) = \max(x + y - 1, 0)$ then $\int_0^1 \int_0^1 F(x, y) d\max(x + y - 1, 0) = \int_0^1 F(x, 1 - x) dx = 1$.

(iv) If $(x_n, y_n) = (\gamma_q(n), \gamma_q(n + 1))$, where $\gamma_q(n)$ is the van der Corput sequence in the base q , then

$\int_0^1 \int_0^1 F(x, y) dg(x, y) = \frac{4}{q} - 1$. Thus for $q = 2$ Player I wins prize 1 and in the case $q > 5$ he loses.

Proposed by O. Strauch.

1.35 Oscillating sums

Directly from J. Arias de Reyna and J. van de Lune (2008):

- $S_\alpha(n) = \sum_{j=1}^n (-1)^{[j\alpha]}$ where α is any real number.
- Denoted by $t_0 = 0, t_1, t_2, \dots$ the sequence of those n for which $S_\alpha(n)$ assumes a value for the first time, i.e., is larger/smaller than ever before.
- Let $\alpha = [a_0; a_1, a_2, \dots]$ and $\beta = \alpha/2 = [b_0; b_1, b_2, \dots]$ be simple continued fraction expansions.

Open Problem: Determine whether the t_k is recurrent and the sequence $\text{sign}(S(t_k))$ is to be purely periodic.

NOTES:

(I) H.D. Ruderman (1977) proposed and D. Borwein (1978) solved (among other) that the series $\sum_{n=1}^{\infty} (-1)^{[n\sqrt{2}]} / n$ converges.

(II) P. Bundschuh (1977) proved that the series $\sum_{n=1}^{\infty} (-1)^{[n\alpha]} / n$ converges for numbers α with bounded b_i of $\beta = \alpha/2 = [b_0; b_1, b_2, \dots]$.

(III) J. Schoisengeier (2007) proved that the series

$$\sum_{n=1}^{\infty} (-1)^{[n\alpha]} / n \text{ and } \sum_{k=0, 2 \nmid q_k}^{\infty} (-1)^k (\log b_{k+1}) / q_k$$

converge simultaneously. Here $\frac{p_k}{q_k}$ are convergents of $\beta = \alpha/2 = [b_0; b_1, b_2, \dots]$.

(IV) A.E. Brouwer and J. van de Lune (1976) have shown that $S_\alpha(n) \geq 0$ for all n if and only if the partial quotients a_{2i} of $\alpha = [a_0; a_1, a_2, \dots]$ are even for all $i \geq 0$.

(V) J. Arias de Reyna and J. van de Lune (2008) proved that $S_\alpha(n)$ is not bounded, so that the corresponding sequence t_k actually is an infinite sequence. They also prove that for every $j \geq 1$ there is an index k such that $t_j - t_{j-1} = Q_k$, where P_k/Q_k is a certain convergent of $\alpha = [a_0; a_1, a_2, \dots]$. They also give a fast algorithm for the computation of $S_\alpha(n)$ for any irrational α and for very large n in terms of $\beta = \alpha/2 = [b_0; b_1, b_2, \dots]$, e.g., $S_{\sqrt{2}}(10^{1000}) = -10$, $S_{\sqrt{2}}(10^{10000}) = 166$, $S_\pi(10^{10000}) = 11726$.

J. ARIAS DE REYNA–J. VAN DE LUNE: *On some oscillating sums*, Uniform Distribution Theory **3** (2008), no. 1, 35–72.

D. BORWEIN: *Solution to problem no. 6105*, Amer. Math. Monthly **85** (1978), 207–208.

H.D. RUDERMAN: *Problem 6105** Amer. Math. Monthly **83** (1977), 573.

P. BUNDSCHUH: *Konvergenz unendlicher Reihen und Gleichverteilung mod 1*, Arch. Math. **29** (1977), 518–523 (Zbl 0365.10025).

J. SCHOISSENGEIER: *The integral mean of discrepancy of the sequence $(n\alpha)$* , Monatsh. Math. **131** (2000), 227–234 (Zbl 1153.11033).

A.E. BROUWER – J. VAN DE LUNE: *A note on certain oscillating sums*, Math. Centrum, Amsterdam, Afd. zuivere Wisk. ZW 90/76, 16 p. 1976 (Zbl 0359.10029).

1.36 Discrepancy system in the unit cube


Let S^{n-1} be the unit sphere of the n -dimensional euclidean space \mathbb{R}^n and a cap is a portion of the sphere cut of by hyperplane. P. Gruber (2009) discuss the problem whether the family of all caps of given size is a discrepancy system. A. Volčič (2011) in the planar case proved that the family of all arcs of S^1 of a constant length l is a discrepancy system if $\frac{l}{2\pi}$ is irrational. For rational $\frac{l}{2\pi}$ there exists a non-uniformly distributed sequence x_m , $m = 1, 2, \dots$, in S^1 such that $\frac{\#\{m \leq N; x_m \in C\}}{N} \rightarrow \frac{l}{2\pi}$ for every arc $C \subset S^1$ of the length l . In the case $n - 1 \geq 2$, Volčič (2011) proved that if s is a zero of a $d + 2$ dimensional Legendre polynomial of even degree, then x_m need not be uniformly distributed even if $\frac{\#\{n \leq N; x_m \in C_s\}}{N} \rightarrow P(C_s)$ for any spherical cap $C_s(u) = \{v \in S^{n-1}; u \cdot v \geq s\}$. Here P is the the normalized Hausdorff measure on the sphere and $u \cdot v$ is the usual scalar product in \mathbb{R}^n . In his proof he used Ungar result (1954) that $\int_{C_s} f dP = 0$ for all spherical caps C_s need not imply $f = 0$.

(I) **Open Problem:** Discuss a similar problem in $[0, 1]^s$. For example:

(II) Let x_n , $n = 1, 2, \dots$, be a sequence in $[0, 1)$ such that $\frac{\#\{n \leq N; x_n \in I\}}{N} \rightarrow |I|$ for all intervals $I \subset [0, 1]$ of the fixed length $|I| = a$. Then x_n need not be u.d. Proof: All $g(x) \in G(x_n)$ must satisfy

$$g(x + a) = g(x) + a \text{ for } x \in [0, 1 - a]. \quad (1)$$

The following d.f. $g(x)$ satisfies (1) but $g(x) \neq x$.



P. UNGAR: *Freak theorem about functions on a sphere*, J. London Math. Soc. **29** (1954), 100–103.
 A. VOLČIČ: *On Grubner's problem concerning uniform distribution on the sphere*, Arch. Math. **97** (2011), no. 4, 385–390.

1.37 Extremes of $\int_0^1 \int_0^1 F(x, y) d_x d_y g(x, y)$ attained at shuffles of M

(I) R.B. Nelsen (1999, p. 59, 3.2.3.): Let $I_i, i = 1, 2, \dots, n$ be a decomposition of the unit interval $[0, 1]$, let π be a permutation of $(1, 2, \dots, n)$, and let $T : [0, 1] \rightarrow [0, 1]$ be an one-to-one map whose graph T is formed by diagonals or anti-diagonals of squares $I_i \times I_{\pi(i)}, i = 1, 2, \dots, n$. Then the copula $C(x, y)$ defined by

$$C(x, y) = |\text{Project}_x(([0, x] \times [0, y]) \cap T)|$$

is called the shuffle of M .

(II) M. Hofer and M.R. Iacò (2013) proved: Let $(a_{i,j}), i, j = 1, 2, \dots, n$ be a real-valued $n \times n$ matrix. Let $I_{i,j} = [\frac{i-1}{n}, \frac{i}{n}] \times [\frac{j-1}{n}, \frac{j}{n}], i, j = 1, 2, \dots, n$ and let the piecewise constant function $F(x, y)$ be defined as

$$F(x, y) = a_{i,j} \text{ if } (x, y) \in I_{i,j}, i, j = 1, 2, \dots, n.$$

Then

$$\max_{g(x,y)\text{-copula}} \int_0^1 \int_0^1 F(x, y) d_x d_y g(x, y) = \frac{1}{n} \sum_{i=1}^n a_{i,\pi^*(i)}.$$

Here $\pi^*(i)$ maximizes $\sum_{i=1}^n a_{i,\pi(i)}$, where π is a permutation of $(1, 2, \dots, n)$. The maximum is attained at $g(x, y) = C(x, y)$, where $C(x, y)$ is the shuffle of M whose graph T is formed by diagonals or anti-diagonals in $I_{i,\pi^*(i)}, i = 1, 2, \dots, n$.

(III) Applying (II) Hofer and Iacò approximate extremes of

$$\int_0^1 \int_0^1 F(x, y) dg(x, y)$$

which respect to copulas $g(x, y)$ by the following: For continuous $F(x, y)$ on $[0, 1]^2$ define piecewise constant functions $F_1(x, y), F_2(x, y)$ as

$$F_1(x, y) = \min_{(u,v) \in I_{i,j}} F(u, v) \text{ if } (x, y) \in I_{i,j}, i, j = 1, 2, \dots, n,$$

$$F_2(x, y) = \max_{(u,v) \in I_{i,j}} F(u, v) \text{ if } (x, y) \in I_{i,j}, i, j = 1, 2, \dots, n, \text{ where}$$

$$I_{i,j} = \left[\frac{i-1}{n}, \frac{i}{n} \right] \times \left[\frac{j-1}{n}, \frac{j}{n} \right].$$

Let $C_0(x, y), C_1(x, y), C_2(x, y)$ be copulas such that

$$\begin{aligned} C_1(x, y) &\text{ maximizes } \int_0^1 \int_0^1 F_1(x, y) \, dg(x, y), \\ C_2(x, y) &\text{ maximizes } \int_0^1 \int_0^1 F_2(x, y) \, dg(x, y) \text{ and} \\ C_0(x, y) &\text{ maximizes } \int_0^1 \int_0^1 F(x, y) \, dg(x, y). \end{aligned}$$

over all copulas $g(x, y)$. Then

$$\begin{aligned} &\int_0^1 \int_0^1 F(x, y) \, dC_0(x, y) \\ &= \lim_{n \rightarrow \infty} \int_0^1 \int_0^1 F_1(x, y) \, dC_1(x, y) = \lim_{n \rightarrow \infty} \int_0^1 \int_0^1 F_2(x, y) \, dC_2(x, y). \end{aligned}$$

(IV) **Open problem:** Using (III) and a numerical experiment the authors conjecture that the extreme of $\int_0^1 \int_0^1 \sin(\pi(x+y)) \, d_x d_y g(x, y)$ is

$$\max_{g(x,y)\text{-copula}} \int_0^1 \int_0^1 \sin(\pi(x+y)) \, d_x d_y g(x, y) = \frac{3}{4\sqrt{2}} - \frac{1}{2\pi}$$

and it is attained at shuffle of M formed by anti-diagonal of $[0, \frac{3}{4}] \times [0, \frac{3}{4}]$ and by diagonal $[\frac{3}{4}, 1] \times [\frac{3}{4}, 1]$. Compare with Problem 1.29.

(V) The copula in (IV) satisfies (IIIb) in 1.29 which is a necessary condition for a copula maximizing a related integral.

(VI) Note that if $x_n, n = 1, 2, \dots$, is a u.d. sequence, then two-dimensional sequence $(x_n, T(x_n))$ has a.d.f $C(x, y)$ and thus for every continuous $F(x, y)$ we have

$$\int_0^1 \int_0^1 F(x, y) \, dC(x, y) = \int_0^1 F(x, T(x)) \, dx.$$

Solution of (IV). S. Thonhauser noted that the problem of optimizing the integral $\int_0^1 \int_0^1 F(x, y) dC(x, y)$ over copulas $C(x, y)$ (cf. Problem 1.29) belongs to the well-known mass transportation problems, or the Monge-Kantorovich transportation problem, see e.g. L. Ambrosio and N. Gigli (2013).

(i) He found that L. Uckelmann (1997) proved: Let

$$F(x, y) = \Phi(x+y) \text{ for } (x, y) \in [0, 1]^2;$$

For $0 < k_1 < k_2 < 2$ let $\Phi(x)$ be a twice differentiable function such that $\Phi(x)$ is strictly convex on $[0, k_1] \cup [k_2, 2]$ and concave on $[k_1, k_2]$, i.e.

$\Phi''(x) > 0$ for $x \in [0, k_1) \cup (k_2, 2]$,


$\Phi''(x) > 0$ for $x \in [0, k_1) \cup (k_2, 2]$.

If α and β are the solutions of

$$\Phi(2\alpha) - \Phi(\alpha + \beta) + (\beta - \alpha)\Phi'(\alpha + \beta) = 0,$$

$$\Phi(2\beta) - \Phi(\alpha + \beta) + (\alpha - \beta)\Phi'(\alpha + \beta) = 0$$

such that $0 < \alpha < \beta < 1$, then the optimal copula $C(x, y)$ is the shuffle of M




with the support

$$\Gamma(x) = \begin{cases} x & \text{for } x \in [0, \alpha] \cup [\beta, 1], \\ \alpha + \beta - x & \text{for } x \in (\alpha, \beta). \end{cases}$$

Then

$$\max \int_0^1 \int_0^1 F(x, y) dC(x, y) = \int_0^\alpha \Phi(2x) dx + (\beta - \alpha)\Phi(\alpha + \beta) + \int_\beta^1 \Phi(2x) dx.$$

(ii) S. Thonhauser adapts (i) for $\Phi(x) = \sin \pi x$. In this case $\Phi(x)$ is convex-concave and he proved that the copula $C(x, y)$ which maximizes $\int_0^1 \int_0^1 \sin \pi(x + y) dC(x, y)$ is the shuffle of M



with the support

$$\Gamma(x) = \begin{cases} \beta - x & \text{for } x \in [0, \beta), \\ x & \text{for } x \in [\beta, 1], \end{cases}$$

where $\beta = 0.7541996008265638$ solves

$$\sin(2\pi\beta) - \sin(\pi\beta) = \pi\beta \cos(\pi\beta).$$

For not colliding with the corresponding literature the function c plays the role of the cost function F from above in the following.

The proof of the $\sin(\pi(x+y))$ result, or originally of Uckelmann's statement, is based on a particular optimality criterion based on c -convex functions. A function $f : X \rightarrow \mathbb{R}$ is called c -convex if it has a representation

$$f(x) = \sup_y \{c(x, y) + a(y)\},$$

for some function a . The associated c -subdifferential of f at x is given by

$$\partial_c f(x) = \{y \mid f(z) - f(x) \geq c(z, y) - c(x, y) \ \forall z \in X\}$$

and $\partial_c f = \{(x, y) \in X \times Y \mid y \in \partial_c f(x)\}$.

Now the criterion is as follows: A pair (X_1, X_2) of random variables with $X_1 \stackrel{d}{\sim} \mu$, $X_2 \stackrel{d}{\sim} \nu$ is an optimal c -coupling between μ and ν if and only if

$$(X_1, X_2) \in \partial_c f \text{ a.s.}$$

for some c -convex function f , equivalently, $X_2 \in \delta_c f(X_1)$ a.s. Additionally one has that $y \in \partial_c f(x)$ if and only if $\exists a(= a(y)) \in \mathbb{R}$ such that

$$\psi_{y,a}(x) = c(x, y) + a(y) = f(x) \quad \text{and} \quad \psi_{y,a}(\xi) = c(\xi, y) + a(y) \leq f(\xi) \quad \forall \xi \in X. \quad (1.0.1)$$

For the case $\Phi(x+y) = \sin(\pi(x+y))$, with one change from convexity to concavity, one can set β as the solution to

$$\phi(2\beta) - \phi(\beta) = \beta\phi'(\beta).$$

Furthermore define:

$$\begin{aligned} f_1(x) &= x\phi'(\beta), & f_2(x) &= \frac{1}{2}(\phi(2x) - \phi(2\beta)) + \beta\phi'(\beta), \\ \psi^1(\xi) &= \phi(\beta - x + \xi) + x\phi'(\beta) - \phi(\beta), \\ \psi^2(\xi) &= \phi(x + \xi) - \frac{1}{2}\phi(2x) - \frac{1}{2}\phi(2\beta) + \beta\phi'(\beta). \end{aligned}$$

For building up a candidate c -convex function define

$$f(x) = f_1(x)I_{[0,\beta)}(x) + f_2(x)I_{[\beta,1]}(x),$$

and put for $\xi \in [0, 1]$:

$$\psi_{\Gamma(x)}(\xi) = \begin{cases} \psi^1(\xi), & x \in [0, \beta), \\ \psi^2(\xi), & x \in [\beta, 1]. \end{cases}$$

Here $\psi_{\Gamma(x)}(\xi)$ plays the role of $\psi_{y,a}(\xi) = c(\xi, y) + a(y)$ with $y = \Gamma(x)$ in (1.0.1). Some tedious calculations show that $y = \Gamma(x)$ is in the c -subdifferential of $f(x)$ for all $x \in [0, 1]$ which implies optimality of this particular coupling and optimality of the distribution induced by $(U, \Gamma(U))$.

(iii) L. Rüschendorf and L. Uckelmann (2000): Let $F(x, y) = \Phi(x + y)$ where $\Phi(x)$ changes its convexity behaviour more than two times. They assume e.g. that $\Phi(x)$ is convex-concave-convex-concave-convex and let $\alpha < \beta$ solve equations in (i) and $0 < \alpha < \beta < \delta < \varepsilon < \gamma$ solve additionally

$$\Phi(2\delta) - \Phi(\delta + \varepsilon) + (\varepsilon - \delta)\Phi'(\delta + \varepsilon) = 0,$$


$$\Phi(2\varepsilon) - \Phi(\delta + \varepsilon) + (\delta - \varepsilon)\Phi'(\delta + \varepsilon) = 0.$$

Then they conjectured that an optimal copula $C(x, y)$ is the shuffle of M with the support

$$\Gamma(x) = \begin{cases} x & \text{for } x \in [0, \alpha] \cup [\beta, \delta] \cup [\varepsilon, 1], \\ \alpha + \beta - x, & \text{for } x \in (\alpha, \beta), \\ \delta + \varepsilon - x & \text{for } x \in (\delta, \varepsilon). \end{cases}$$

The authors do not have a formal proof of this statement but one can see however analytically that $\Gamma(x)$ is optimal in the class of all copulas of this special form.

Example. Consider $\Phi(x) = 0.092x^6 - 0.498x^5 + x^4 + 0.910x^3 + 0.406x^2$, then $\Phi(x)$ has four convexity changes and one obtains $\alpha = 0.1$, $\beta = 0.3$, $\delta = 0.6$, $\varepsilon = 0.8$. The shuffle of M is



- L. AMBROSIO– N. GIGLI: *A User's Guide to Optimal Transport Modelling and Optimisation of Flows an Networks*, Lecture Notes in Mathematics, Springer, 2013.
- M. HOFER – M.R. IACÒ: *Optimal bounds for integrals with respect to copulas and applications*, Journal of Optimization Theory and Applications (2013), accepted.
- R.B. NELSEN: *An Introduction to Copulas. Properties and Applications*, Lecture Notes in Statistics **139** Springer-Verlag, New York, 1999.
- L. RÜSCHENDORF – L. UCKELMANN: *Numerical and analytical results for the transportation problem of Monge-Kantorovich*, *Metrika*, 51 (2000), 25–258.
- L. UCKELMANN: *Optimal couplings between one-dimensional distributions*, Distribution with given Marginals and Moment Problems, V. Beneš and J. Štěpán (eds.) (Prague, 1996), 275–281, Kluwer, 1997.

1.38 Two-dimensional Benford's law

Let $x_n > 0$, $y_n > 0$, $n = 1, 2, \dots$ and $b > 1$ be an integer base, K_1, K_2 be positive integers, and

$$K_1 = k_1^{(1)} k_2^{(1)} \dots k_{r_1}^{(1)} \text{ in base } b,$$

$$K_2 = k_1^{(2)} k_2^{(2)} \dots k_{r_2}^{(2)} \text{ in base } b,$$

$$u_1 = \log_b \left(\frac{K_1}{b^{r_1-1}} \right),$$

$$u_2 = \log_b \left(\frac{K_1+1}{b^{r_1-1}} \right),$$

$$v_1 = \log_b \left(\frac{K_2}{b^{r_2-1}} \right),$$

$$v_2 = \log_b \left(\frac{K_2+1}{b^{r_2-1}} \right).$$

(I) As in Problem 1.32 we have

first r_1 digits (starting a non-zero digit) of $x_n = K_1 \iff \{\log_b x_n\} \in [u_1, u_2)$,

first r_2 digits (starting a non-zero digit) of $y_n = K_2 \iff \{\log_b y_n\} \in [v_1, v_2)$.

Denote ²

$$F_N(x, y) = \frac{\#\{n \leq N; \{\log_b x_n\} < x \text{ and } \{\log_b y_n\} < y\}}{N}.$$

(II) From definition of d.f.'s the following holds:

Let $g(x, y) \in G(\{\log_b x_n\}, \{\log_b y_n\})$ and $\lim_{k \rightarrow \infty} F_{N_k}(x, y) = g(x, y)$ for $(x, y) \in [0, 1]^2$. Then

$$\begin{aligned} & \lim_{k \rightarrow \infty} \frac{\#\{n \leq N_k; \text{ first } r_1 \text{ digits of } x_n = K_1 \text{ and first } r_2 \text{ digits of } y_n = K_2\}}{N_k} \\ &= g(u_2, v_2) + g(u_1, v_1) - g(u_2, v_1) - g(u_1, v_2). \end{aligned}$$

²In the following the sentence "starting a non-zero digit" we will not mention.

(III) As example we give:

$$G(\{\log_b n\}, \{\log_b(n+1)\}) \\ = \left\{ g_u(x, y) = \frac{b^{\min(x,y)} - 1}{b-1} \cdot \frac{1}{b^u} + \frac{b^{\min(x,y,u)} - 1}{b^u}; u \in [0, 1] \right\}.$$

By Sklar theorem

$$g_u(x, y) = \min(g_u(x), g_u(y)), \text{ where} \\ g_u(x) = \frac{b^x - 1}{b-1} \cdot \frac{1}{b^u} + \frac{b^{\min(x,u)} - 1}{b^u}.$$

Put $x_n = \log_b n \bmod 1$ and $y_n = \log_b(n+1) \bmod 1$. Then by (II)

$$\lim_{k \rightarrow \infty} \frac{\#\{n \leq N_k; \text{ first } r_1 \text{ digits of } x_n = K_1 \text{ and first } r_2 \text{ digits of } y_n = K_2\}}{N_k} \\ = g_u(u_2, v_2) + g_u(u_1, v_1) - g_u(u_2, v_1) - g_u(u_1, v_2).$$

If $K_1 = K_2$ then $= g_u(u_2) - g_u(u_1)$. It can be found directly.

In the following examples we use statistical independent sequences:

Let $x_n \in [0, 1)$, $n = 1, 2, \dots$, be an u.d. sequence. Then

(IV) x_n and $\log_b n \bmod 1$ are statistically independent (Rauzy (1973) see [SP, p. 2–27, 2.3.6.].

(V) x_n and $\log_b(n \log n) \bmod 1$ are statistically independent (Y. Ohkubo (2011)).

(VI) x_n and $\log_b p_n \bmod 1$ are statistically independent (Y. Ohkubo (2011)).

(VII) The sequences $\log_b n \bmod 1$, $\log_b p_n \bmod 1$ and $\log_b \log n \bmod 1$ have the same set of d.f.'s. (Y. Ohkubo (2011)).

(VIII) From (IV) it follows: Let $x_n \in [0, 1)$, $n = 1, 2, \dots$, be u.d. sequence. Then

$$G(x_n, \{\log_b n\}) = \{g_u(x, y) = x \cdot g_u(y); u \in [0, 1]\},$$

where

$$g_u(x) = \frac{b^x - 1}{b-1} \cdot \frac{1}{b^u} + \frac{b^{\min(x,u)} - 1}{b^u}$$

and $F_{N_k}(x, y) \rightarrow g_u(x, y)$ if $\{\log_b N_k\} \rightarrow u$.

(IX) Let $x_n \in [0, 1)$, $n = 1, 2, \dots$, be u.d. sequence. Then

$$G(x_n, \{\log_b p_n\}) = \{g_u(x, y) = x \cdot g_u(y); u \in [0, 1]\},$$

where

$$g_u(x) = \frac{b^x - 1}{b - 1} \cdot \frac{1}{b^u} + \frac{b^{\min(x,u)} - 1}{b^u}$$

and $F_{N_k}(x, y) \rightarrow g_u(x, y)$ if $\{\log_b N_k\} \rightarrow u$.

(X) We have

$$G(\{\log_b F_n\}, \{\log_b p_n\}) = \{x.g_u(y); u \in [0, 1]\}$$

and let $\{\log_b N_k\} \rightarrow u$. Then

$$\begin{aligned} & \lim_{k \rightarrow \infty} \frac{\#\{n \leq N_k; \text{ first } r_1 \text{ digits of } F_n = K_1 \text{ and first } r_2 \text{ digits of } p_n = K_2\}}{N_k} \\ &= u_2 g_u(v_2) + u_1 g_u(v_1) - u_2 g_u(v_1) - u_1 g_u(v_2), \end{aligned}$$

where F_n is the sequence of Fibonacci numbers and p_n is the increasing sequence of all primes and

$$\begin{aligned} u_1 &= \log_b \left(\frac{K_1}{b^{r_1-1}} \right), u_2 = \log_b \left(\frac{K_1 + 1}{b^{r_1-1}} \right), \\ v_1 &= \log_b \left(\frac{K_2}{b^{r_2-1}} \right), v_2 = \log_b \left(\frac{K_2 + 1}{b^{r_2-1}} \right), \\ g_u(x) &= \frac{b^x - 1}{b - 1} \cdot \frac{1}{b^u} + \frac{b^{\min(x,u)} - 1}{b^u}. \end{aligned}$$

(XI) Problem 1.38 is inspired by the result of F. Luca and P. Stanica (2013): There exists infinite many n such that Fibonacci number F_n starts with digits K_1 and $\phi(F_n)$ starts with digits K_2 in the base b representation. Here K_1 and K_2 are arbitrary and $\varphi(x)$ is the Euler function.

We see that (XI) is equivalent to the sequence

$$(\log_b F_n, \log_b \varphi(F_n)) \bmod 1, \quad n = 1, 2, \dots,$$

is everywhere dense in $[0, 1]^2$, but the authors use the following method:

(i) By the first author $\varphi(F_n)/F_n$ is dense in $[0, 1]$. Thus, for an interval I with arbitrary small length which containing K_2/K_1 , there exists $\varphi(F_a)/F_a \in I$.

(ii) Then $\varphi(F_{ap})/F_{ap} \in I$ for all sufficiently large primes p .

(iii) There exists infinitely many primes p such that F_{ap} starts with K_1 .

(iv) Finally, multiplying I by F_{ap} they find $\varphi(F_{ap})$ which starts with K_2 .

(XII) Definition: The sequence (x_n, y_n) , $x_n > 0$, $y_n > 0$, $n = 1, 2, \dots$, satisfies 2-dimensional B.L. in base b , if for every K_1, K_2 we have

$$\lim_{N \rightarrow \infty} \frac{\#\{n \leq N; \text{ the first } r \text{ digits of } x_n = K_1 \text{ and the first } l \text{ digits of } y_n = K_2\}}{N} \\ = \log_b \left(1 + \frac{1}{K_1}\right) \cdot \log_b \left(1 + \frac{1}{K_2}\right).$$

(XIII) From definition follows: The sequence (x_n, y_n) satisfies 2-dimensional Benford law (B.L.) if and only if $(\log_b x_n, \log_b y_n) \bmod 1$ is u.d. in $[0, 1]^2$.

(XVI) **Open problem:** Prove that the sequence (n^{n^2}, n^n) satisfies 2-dimensional B.L. in any base b . Motivation is that by [SP, 3.13.4.] the sequence $(n^2 \log n, n \log n) \bmod 1$ is u.d. in $[0, 1]^2$.

(XV) **Open problem:** Prove that the sequence $(\log_b n, \log_b \log n) \bmod 1$ is dense in $[0, 1]^2$. If this is true, then there exists infinite many n such that n starts with digits K_1 and $\log n$ starts with digits K_2 in the base b representation, where K_1 and K_2 are arbitrary positive integers. By [SP, 3.13.5.] the sequence $(\log n, \log \log n) \bmod 1$ is dense in $[0, 1]^2$ but not u.d.

Proposed by O. Strauch.

F. LUCA – P. STANICA: *On the first digits of the Fibonacci numbers and their Euler function*, Uniform Distribution Theore accepted.

Y. OHKUBO: *On sequences involving primes* Uniform Distribution Theory, Vol. 6 (2011), no. 2, 221–238.

M. SKLAR: *Fonctions de répartition à n dimensions et leurs marges*, Publ. Inst. Statist. Univ. Paris **8** (1959), 229–231 (MR **23**# A2899).

G. RAUZY: *Propriétés Statistiques de Suites Arithmétiques*, *Le Mathématicien* Vol **15**, Collection SUP, Presses Universitaires de France, Paris, 1976, 133 pp. (MR **53**#13152).

O. STRAUCH – Š. PORUBSKÝ: *Distribution of Sequences: A Sampler*, Peter Lang, Frankfurt am Main, 2005.

1.39 Problem of N.M. Korobov

M.B. Levin(1999):

(I) A number α is normal in the base q if and only if the sequence $x_n = \alpha q^n \bmod 1$, $n = 1, 2, \dots$ is u.d. in $[0, 1)$. See also 2.4. Exponential sequences (40)–(50).

(II) N.M. Korobov (1955) proposed to found normal α with minimal discrepancy $D_N(x_n)$. He found α with $D_N = O(N^{-1/2})$.

(III) J. Schiffer (1986) was proved: Let $p(n)$ be a polynomial in n with rational coefficients, $p(x)$ not constant and let $d_n, n = 1, 2, \dots$, be a bounded

sequence of rational numbers, such that $p(n) + d_n$ is a positive integer for all $n \geq 1$. Then $D_N = O(\log^{-1})N$ for $\alpha = 0.(p(1) + d_1)(p(2) + d_2)\dots$. Moreover, if $p(x) \geq 1$ be a linear polynomial with rational coefficients, then $D_N \geq K/\log N$ for $K > 0$ and $N = 1, 2, \dots$ for $\alpha = 0.[p(1)][p(2)]\dots$, i.e. for Champernowne normal α .

(IV) N.M. Korobov (1966) for q -prime and M.B. Levin for arbitrary integer q found $D_N = O(N^{-2/3} \log^{4/3} N)$.

(V) M.B. Levin (1999) construct normal α such that $D_N = O(N^{-1} \log^2 N)$.

N.M. KOROBOV: *Numbers with bounded quotient and their applications to questions of Diophantine approximation*, Izv. Akad. Nauk SSSR Ser. Mat. **19** (1955), 361–380.

N.M. KOROBOV: *Distribution of fractional parts of exponential function*, Vestnik Moskov. Univ. Ser. I Mat. Mekh. 21 (1966), no. 4, 42–46.

M.B. LEVIN: *The distribution of fractional parts of the exponential function*, Soviet Math. (Izv. VUZ) 21 (1977), no. 11, 41–47.

M.B. LEVIN: *On the discrepancy estimate of normal numbers*, Acta Arith. LXXXVIII (1999), no. 2, 99–111.

J. SCHIFFER: *Discrepancy of normal numbers*, Acta Arith. 47 (1986), 175–186.

2. Open theories

2.1 Uniform distribution theories.

There are some different ways for generalizing of the classical u.d. theory, see [KN, Chap. 3 and 4], [H, Chap. 2], [DT, Chap. 2] and [SP, p. 1–5, 1.5]. For example:

- Points of investigated sequences x_n are elements from a general space.
- For basic sets $A_x = \{n \in \mathbb{N}; x_n \in [0, x)\}$ in the definition of u.d. as $d(A_x) = x$, the asymptotic density d is exchanged by another types of densities.
- The asymptotic density d is preserved but in A_x the relation $x_n \in [0, x)$ is exchanged by more complicated relations (cf. O. Strauch (1998)).

Here we start with a main theorem of u.d. theory due to H. Weyl (cf. [KN, p. 2, Th. 1.1], [SP, p. 1–4, Th. 1.4.0.1]):

Weyl's limit relation. *The sequence $u(n)$, $n = 1, 2, \dots$ from the unit interval $[0, 1]$ is u.d. if and only if for every continuous $f : [0, 1] \rightarrow \mathbb{R}$ we have*

$$\int_0^1 f(x) dx = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(u(n)).$$

This relation can be used as a definition of u.d. of $u(n)$ and also for definition of u.d. in abstract spaces, see [KN, p. 171, Def. 1.1]: Let \mathbf{X} be a compact Hausdorff space and $C(\mathbf{X})$ consists of all real-valued continuous functions on \mathbf{X} . Let dX be a nonnegative regular normed Borel measure in \mathbf{X} . The sequence $u(n) \in \mathbf{X}$, $n = 1, 2, \dots$ is called u.d. in \mathbf{X} with respect to dX if

$$\forall (f \in C(\mathbf{X})) \int_{\mathbf{X}} f(X) dX = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(u(n)).$$

The basic application of Weyl's limit relation is a possibility computing the Riemann integral $\int_0^1 f(x) dx$ on $[0, 1]$ of a continuous function $f(x)$ as the limit $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(u(n))$ of arithmetic means of $f(x)$ (quasi-Monte Carlo method) and vice-versa, the limit of arithmetic means by integral. Looking at $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(u(n))$ as an integral defined on \mathbb{Z}^+ , then the classical u.d. theory is the theory of coherence between two types of

integrals. Thus the concept of u.d. theory can be generalized (see O. Strauch (1999, Chap. 4)) to a theory of the integral equation

$$\int_{\mathbf{X}} f(X) dX = \int_{\mathbf{Y}} f(u(Y)) dY, \quad (1)$$

of two types of integrals, where \mathbf{X}, \mathbf{Y} are arbitrary spaces equipped with integrals or measures dX and dY , respectively, or more generally, equipped with functionals which in the following we also call integrals. Here $f : \mathbf{X} \rightarrow \mathbb{R}$ and $u : \mathbf{Y} \rightarrow \mathbf{X}$. The main problem is to compute integral of the first type on the left-hand (1) by the integral of the second type on the right-hand side. This is our aim in these new u.d. theories. Here are a few selected spaces with theories of integration.

- $\mathbf{X}_1 = [0, 1]$, equipped with the integral $\int_0^1 f(x) dx$;
- $\mathbf{X}_2 = \{1, 2, \dots\}$, equipped with the integral $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(n)$;
- $\mathbf{X}_3 = [0, +\infty)$, equipped with the integral $\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(x) dx$;
- $\mathbf{X}_4 = [0, 1]$, equipped with the integral $\max_{x \in [0, 1]} f(x)$;
- $\mathbf{X}_5 = \{1, 2, \dots\}$, equipped with the integral $\limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(n)$;
- $\mathbf{X}_6 = [0, +\infty)$, equipped with the integral $\int_0^{+\infty} f(x) dx$;
- $\mathbf{X}_7 = \{1, 2, \dots\}$, equipped with the integral $\lim_{N \rightarrow \infty} \frac{1}{N^\alpha} \sum_{n=1}^N f(n)$;
- $\mathbf{X}_8 = [0, 1]^s$, equipped with the integral $\int_{[0, 1]^s} f(\mathbf{x}) d\mathbf{x}$;
- $\mathbf{X}_9 = \{1, 2, \dots\} \times [0, +\infty)$, equipped with the integral $\lim_{N, T \rightarrow \infty} \frac{1}{NT} \sum_{n=1}^N \int_0^T f(n, x) dx$.

Varying couples $(\mathbf{X}_i, \mathbf{X}_j)$ we find the following known u.d. theories

- $\int_0^1 f(x) dx = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(u(n))$ - classical u.d. theory;
- $\int_0^1 f(x) dg(x) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(u(n))$ - theory of g -distributed sequences, see [KN, pp. 54–57] and [SP, p. 1–11, 1.8.1];
- $\int_0^1 f(x) dx = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(u(x)) dx$ - c.u.d. theory, see [KN, pp. 78–87] and [DT, pp. 277–300];
- $\int_0^1 f(x) dx = \int_0^1 f(u(x)) dx$ - theory of u.d. preserving functions, it was introduced by Š. Porubský, T. Šalát and O. Strauch (1998), see [SP, p. 2–45, 2.5.1];
- $\max_{x \in [0, 1]} f(x) = \limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(u(n))$ - theory of maldistributed sequences, it was introduced by G. Myerson (1993), see [SP, p. 1–19, 1.8.10];
- $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(n) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(u(n))$ - u.d. theory in \mathbb{Z} , see [KN, pp. 305–319].

These examples lead to the following common definition.

Definition. Let \mathcal{C} be a set of functions $f : \mathbf{X} \rightarrow \mathbb{R}$. The u.d. theory $(\mathcal{C}, \mathbf{X}, \mathbf{Y})$ -u.d. is a theory of functions $u : \mathbf{Y} \rightarrow \mathbf{X}$ satisfying

$$\forall(f \in \mathcal{C}) \int_{\mathbf{X}} f(X) dX = \int_{\mathbf{Y}} f(u(Y)) dY. \quad (2)$$

These functions u are called u.d. in $(\mathcal{C}, \mathbf{X}, \mathbf{Y})$ -u.d. theory.

- The (\mathbf{X}, \mathbf{Y}) -u.d. theory is a theory of the integral equation (1), in which the set \mathcal{C} is not strictly specified.
- The (\mathbf{Y}, \mathbf{X}) -u.d. is the inverse theory to the (\mathbf{X}, \mathbf{Y}) -u.d.
- The (\mathbf{X}, \mathbf{Y}) -u.d. theory is empty if there does not exist any u for some class f such that (1) is valid.
- The (\mathbf{X}, \mathbf{X}) -u.d. theory of $\int_{\mathbf{X}} f(X) dX = \int_{\mathbf{X}} f(u(X)) dX$ is a theory of integration, where the inside part $u(X)$ can be omitted. We shall call it u.d. preserving theory (abbreviating u.d.p. theory), because the equation

$$\int_{\mathbf{X}} f(X) dX = \int_{\mathbf{X}} f(u(X)) dX = \int_{\mathbf{Z}} f(u(v(Z))) dZ \quad (3)$$

gives

Theorem. Let u be u.d. in $(\mathcal{C}, \mathbf{X}, \mathbf{X})$ -u.d. theory and $\mathcal{C} \circ u = \mathcal{C}$, where $\mathcal{C} \circ u = \{f(u(X)); f \in \mathcal{C}\}$. Then

$$v \text{ is u.d. in } (\mathcal{C}, \mathbf{X}, \mathbf{Z}) \iff u \circ v \text{ is u.d. in } (\mathcal{C}, \mathbf{X}, \mathbf{Z}).$$

- Using the equation

$$\int_{\mathbf{X}} f(X) dX = \int_{\mathbf{Y}} f(u(Y)) dY = \int_{\mathbf{Z}} f(u(v(Z))) dZ \quad (4)$$

a new (\mathbf{Y}, \mathbf{Z}) -u.d. theory can be found by means of known (\mathbf{X}, \mathbf{Y}) -u.d. and (\mathbf{X}, \mathbf{Z}) -u.d. theory.

In the following we list some **new u.d theories**:

- (I) $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(n) = \int_0^1 f(u(x)) dx$ is an inverse theory to the classical u.d. one.
- (II) $\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(x) dx = \int_0^1 f(u(x)) dx$ is an inverse theory to the c.u.d. theory.
- (III) $\int_0^{+\infty} f(x) dx = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(u(n))$ is an empty u.d. theory.
- (IV) In $\int_0^1 \int_0^1 f(x, y) dx dy = \int_0^1 f(u(x), v(x)) dx$, the curve $(u(x), v(x))$ must be Peano. The equation (4) in the form $\int_0^1 \int_0^1 f(x, y) dx dy =$

$\int_0^1 f(u(x), v(x)) dx = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(u(w(n)), v(w(n)))$ gives that $(u(x), v(x))$ is u.d. in this theory if and only if for every classical u.d. sequence $w(n), n = 1, 2, \dots$, in $[0, 1]$, the sequence $(u(w(n)), v(w(n)))$ is u.d. in $[0, 1]^2$.

- (V) In $\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(x) dx = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(u(n))$, assuming that \mathcal{C} contains only bounded continuous functions f which have bounded variation on every interval $[0, n]$ such that $V(f/[0, n]) = \mathcal{O}(n)$, then we can construct u.d. sequence $u(n)$ directly: $u(i), i = 1, 2, \dots, n^2$ is composed with n parts that are contained in the intervals $[0, 1), [1, 2), \dots, [n-1, n)$, all are congruent mod 1 and having discrepancy $D_n^* \rightarrow 0$. By applying Koksma inequality we have

$$\left| \frac{1}{n} \int_0^n f(x) dx - \frac{1}{n^2} \sum_{i=1}^{n^2} f(u(i)) \right| \leq D_n^* \frac{V(f/[0, n])}{n}.$$

- (VI) In $\int_0^1 f(x) dx = \lim_{N, T \rightarrow \infty} \frac{1}{NT} \sum_{n=1}^N \int_0^T f(u(n, x)) dx$ we can use the following L^2 discrepancy

$$D_{N, T}^{(2)}(u) = \frac{1}{3} + \frac{1}{NT} \sum_{n=1}^N \int_0^T (u(n, x))^2 dx - \frac{1}{NT} \sum_{n=1}^N \int_0^T u(n, x) dx - \frac{1}{2(NT)^2} \sum_{m, n=1}^N \int_0^T \int_0^T |u(m, x_1) - u(n, x_2)| dx_1 dx_2,$$

which characterizes u.d. of $u(n, x)$.

- (VII) The $\int_0^1 f(x) dx = \int_0^1 f(u(x)) dx$ is known u.d.p. theory, where the (3) has the form

$$\int_0^1 f(x) dx = \int_0^1 f(u(x)) dx = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(u(v(n)))$$

- (*) $u(x)$ is u.d.p. $\iff u(v(n))$ is u.d. in $[0, 1]$.

In [SP, p. 2–45, 2.5.1] the result (*) was used as definition: The map $u : [0, 1] \rightarrow [0, 1]$ is called **uniform distribution preserving** (abbreviated u.d.p.) if for any u.d. sequence $x_n, n = 1, 2, \dots$, in $[0, 1]$ the sequence $u(x_n)$ is also u.d. In this u.d.p. theory we register the following progress:

A Riemann integrable function $u : [0, 1] \rightarrow [0, 1]$ is a u.d.p. transformation if and only if one of the following conditions is satisfied:

- (i) $\int_0^1 h(x) dx = \int_0^1 h(u(x)) dx$ for every continuous $h : [0, 1] \rightarrow \mathbb{R}$.
- (ii) $\int_0^1 (u(x))^k dx = \frac{1}{k+1}$ for every $k = 1, 2, \dots$.
- (iii) $\int_0^1 e^{2\pi i k u(x)} dx = 0$ for every $k = \pm 1, \pm 2, \dots$.
- (iv) There exists an increasing sequence of positive integers N_k and an N_k -almost u.d. sequence x_n for which the sequence $u(x_n)$ is also N_k -almost u.d.
- (v) There exists an almost u.d. sequence x_n in $[0, 1)$ such that the sequence $u(x_n) - x_n$ converges to a finite limit.
- (vi) There exists at least one $x \in [0, 1]$ of which orbit $x, u(x), u(u(x)), \dots$ is almost u.d.

- (vii) u is measurable in the Jordan sense and $|u^{-1}(I)| = |I|$ for every subinterval $I \subset [0, 1]$.
- (viii) $\int_0^1 u(x) dx = \int_0^1 x dx = \frac{1}{2}$,
 $\int_0^1 (u(x))^2 dx = \int_0^1 x^2 dx = \frac{1}{3}$,
 $\int_0^1 \int_0^1 |u(x) - u(y)| dx dy = \int_0^1 \int_0^1 |x - y| dx dy = \frac{1}{3}$.

From the other properties of u.d.p. transformations let us mention:

- (ix) Let u_1, u_2 be u.d.p. transformations and α a real number. Then $u_1(u_2(x))$, $1 - u_1(x)$ and $u_1(x) + \alpha \bmod 1$ are again u.d.p. transformations.
- (x) Let u_n be a sequence of u.d.p. transformations uniformly converging to u . Then u is u.d.p.
- (xi) Let $u : [0, 1] \rightarrow [0, 1]$ be piecewise differentiable. Then u is u.d.p. if and only if $\sum_{x \in u^{-1}(y)} \frac{1}{|u'(x)|} = 1$ for all but a finite number of points $y \in [0, 1]$.
- (xii) A piecewise linear transformation $u : [0, 1] \rightarrow [0, 1]$ is u.d.p. if and only if $|J_j| = |I_{j,1}| + \dots + |J_{j,n_j}|$ for every $J_j = (y_{j-1}, y_j)$, where $0 = y_0 < y_1 < \dots < y_m = 1$ is the sequence of ordinates of the ends of line segment components of the graph of f and $u^{-1}(J_j) = I_{j,1} \cup \dots \cup J_{j,n_j}$.
- (xiii) $u(x)$ is u.d.p. if and only if

$$\int_0^1 \int_0^1 F(u(x), u(y)) dx dy = 0,$$

where $F(x, y) = (1/2)(|x - u(y)| + |y - u(x)| - |x - y| - |u(x) - u(y)|)$.

NOTES: The problem to find all continuous u.d.p. is formulated in Ja.-I. Rivkid (1973). The results (i)-(vii), (ix)-(xii) are proved in Š. Porubský, T. Šalát and O. Strauch (1988). The criterion (viii) and (xiii) are given in O. Strauch (1999, p. 116, 67). Some parts of these results are also proved independently in W. Bosch (1988). R.F. Tichy and R. Winkler (1991) gave a generalization for compact metric spaces. Some related results can be found in: M. Paštéka (1987), Y. Sun (1993, 1995), P. Schatte (1993), S.H. Molnár (1994) and J. Schmeling and R. Winkler (1995).

Multidimensional u.d.p. map $\Phi : [0, 1]^s \rightarrow [0, 1]^s$ is called uniformly distribution preserving (u.d.p.) map if for every uniformly distributed (u.d.) sequence \mathbf{x}_n , $n = 1, 2, \dots$, the image $\Phi(\mathbf{x}_n)$ is again u.d. The main criterion of u.d.p. map is

Theorem 2.1.0.1. *A map $\Phi(\mathbf{x})$ is u.d.p. if and only if for every continuous $f : [0, 1]^s \rightarrow \mathbb{R}$ we have*

$$\int_{[0,1]^s} f(\Phi(\mathbf{x})) d\mathbf{x} = \int_{[0,1]^s} f(\mathbf{x}) d\mathbf{x}.$$

The multi-dimensional u.d.p. functions are:

- (i) $\Phi(\mathbf{x}) = \mathbf{x} \oplus \boldsymbol{\sigma}$, where
 $x \oplus \sigma = \frac{x_0 + \sigma_0 \pmod{b}}{b} + \frac{x_1 + \sigma_1 \pmod{b}}{b^2} + \dots$ and
 $\mathbf{x} \oplus \boldsymbol{\sigma} = (x_1 \oplus \sigma_1, x_2 \oplus \sigma_2, \dots, x_s \oplus \sigma_s)$;
- (ii) $\Phi(\mathbf{x}) = (\Phi_1(x_1), \dots, \Phi_s(x_s))$, where $\Phi_n(x)$ are one-dimensional u.d.p. maps, especially
- (iii) $\Phi(\mathbf{x}) = \mathbf{b}^\alpha \mathbf{x} \pmod{1} = (b_1^{\alpha_1} x_1, \dots, b_s^{\alpha_s} x_s) \pmod{1}$;
- (iv) $\Phi(\mathbf{x}) = \mathbf{x} + \boldsymbol{\sigma} \pmod{1} = (x_1 + \sigma_1, \dots, x_s + \sigma_s) \pmod{1}$;
- (v) $\Phi(\mathbf{x}) = (A\mathbf{x})^T \pmod{1}$, where A is an $s \times s$ nonsingular integer matrix, cf. S. Steinerberger (Th.2, 2009);
- (vi) $\Phi(\mathbf{x}) = \pi(\mathbf{x})$, where $\pi(\mathbf{x}) = (x_{\pi(1)}, \dots, x_{\pi(n)})$ is a permutation.

Open question: Find another multidimensional u.d.p.

Proposed by O. Strauch.

W. BOSCH: *Functions that preserve uniform distribution*, Trans. Amer. Math. Soc. **307** (1988), no. 1, 143–152 (MR 89h:11046).

S.H. MOLNÁR: *Sequences and their transforms with identical asymptotic distribution function modulo 1*, Studia Sci. Math. Hungarica **29** (1994), no. 3–4, 315–322 (MR 95j:11071).

M. PAŠTÉKA: *On distribution functions of sequences*, Acta Math. Univ. Comenian. **50–51** (1987), 227–235 (MR 90c:11115).

Š. PORUBSKÝ – T. ŠALÁT – O. STRAUCH: *Transformations that preserve uniform distribution*, Acta Arith. **49** (1988), 459–479 (MR 89m:11072; Zbl 656.10047).

JA.I. RIVKIND: *Problems in Mathematical Analysis*, (Russian), 2nd edition ed., Izd. Vyšejšaja škola, Minsk, 1973.

P. SCHATTE: *On transformations of distribution functions on the unit interval – a generalization of the Gauss – Kuzmin – Lévy theorem*, Zeitschrift für Analysis und ihre Anwendungen **12** (1993), 273–283 (MR 95d:11098).

J. SCHMELING – R. WINKLER: *Typical dimension of the graph of certain functions*, Monatsh. Math. **119** (1995), 303–320 (MR 96c:28005).

S. STEINERBERGER: *Uniform distribution preserving mappings and variational problems*, Uniform Distribution Theory **4** (2009), no. 1, 117–145.

O. STRAUCH: *Distribution of Sequences*, (in Slovak), Mathematical Institute of the Slovak Academy of Sciences, (DSc Thesis), Bratislava, Slovakia, 1999.

Y. SUN: *Some properties of uniform distributed sequences*, J. Number Theory **44** (1993), no. 3, 273–280 (MR 94h:11068).

Y. SUN: *Isomorphisms for convergence structures*, Adv. Math. **116** (1995), no. 2, 322–355 (MR 97c:28031).

R.F. TICHY – R. WINKLER: *Uniform distribution preserving mappings*, Acta Arith. **60** (1991), no. 2, 177–189 (MR 93c:11054).

2.2 Distribution functions of sequences.

For a multi-dimensional sequence \mathbf{x}_n , $n = 1, 2, \dots$, in $[0, 1]^s$, the theory of the set $G(\mathbf{x}_n)$ of all d.f.s of \mathbf{x}_n , $n = 1, 2, \dots$, is **open**. A motivation to study of $G(\mathbf{x}_n)$ is the deterministic analysis of sequences in 2.3.¹ The set

¹We shall identify the notion of the **distribution** of a sequence $\mathbf{x}_n \pmod{1}$, $n = 1, 2, \dots$, with the set $G(\mathbf{x}_n \pmod{1})$, i.e., the distribution of $\mathbf{x}_n \pmod{1}$ is known if we know the set $G(\mathbf{x}_n \pmod{1})$.

$G(\mathbf{x}_n)$ has the following fundamental properties for every sequence \mathbf{x}_n in $[0, 1]^s$:

(I) $G(\mathbf{x}_n)$ is non-empty, and it is either a singleton or has infinitely many elements. Precisely, $G(\mathbf{x}_n)$ is non-empty, closed and connected set in the weak topology, and these properties are characteristic for $G(\mathbf{x}_n)$, i.e., given a non-empty set H of distribution functions, there exists a sequence \mathbf{x}_n in $[0, 1]^s$ such that $G(\mathbf{x}_n) = H$ if and only if H is closed and connected.

(II) There are no general methods for computing $G(\mathbf{x}_n)$, without the following one: Let $F(\mathbf{x}, \mathbf{y})$ be a continuous function defined on $[0, 1]^s \times [0, 1]^s$ and let $G(F)$ be the set of all d.f.'s $g(\mathbf{x})$ satisfying

$$\int_{[0,1]^s \times [0,1]^s} F(\mathbf{x}, \mathbf{y}) dg(\mathbf{x}) dg(\mathbf{y}) = 0. \quad (1)$$

If the sequence \mathbf{x}_n , $n = 1, 2, \dots$, satisfies

$$\lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{n,m=1}^N F(\mathbf{x}_m, \mathbf{x}_n) = 0,$$

then $G(\mathbf{x}_n) \subset G(F)$.

Open problem 1. Find a method for solving the moment problem (1).

(III) The definition of d.f.s of sequences in the multi-dimensional case is different as in the one-dimensional one.

- If $\mathbf{x} = (x_1, \dots, x_s) \in \mathbb{R}^s$ is given, then $\mathbf{x} \bmod 1$ denotes the sequence $(\{x_1\}, \dots, \{x_s\})$. If $\mathbf{x}_n = (x_{n,1}, \dots, x_{n,s})$ is the sequence of points in \mathbb{R}^s then we define

- the s -dimensional counting function by

$$A([u_1, v_1] \times \dots \times [u_s, v_s]; N; \mathbf{x}_n \bmod 1) = \#\{n \leq N; \{x_{n,1}\} \in [u_1, v_1), \dots, \{x_{n,s}\} \in [u_s, v_s)\}.$$

- the s -dimensional step d.f. also called the empirical distribution by

(i) $F_N(\mathbf{x}) = \frac{1}{N} A([0, x_1] \times \dots \times [0, x_s]; N; \mathbf{x}_n \bmod 1)$ if $\mathbf{x} \in [0, 1]^s$,

(ii) $F_N(\mathbf{x}) = 0$ for every \mathbf{x} having a vanishing coordinate,

(iii) $F_N(\mathbf{1}) = 1$,

(iv) $F_N(1, \dots, 1, x_{i_1}, 1, \dots, 1, x_{i_2}, 1, \dots, 1, x_{i_l}, 1, \dots, 1) = F_N(x_{i_1}, x_{i_2}, \dots, x_{i_l})$ for every restricted l -dimensional face sequence $(x_{n,i_1}, x_{n,i_2}, \dots, x_{n,i_l})$ of \mathbf{x}_n for $l = 1, 2, \dots, s$.

Then

- If $f : [0, 1]^s \rightarrow \mathbb{R}$ is continuous, again

$$\frac{1}{N} \sum_{n=1}^N f(\mathbf{x}_n \bmod 1) = \int_{[0,1]^s} f(\mathbf{x}) dF_N(\mathbf{x}).$$

- A function $g : [0, 1]^s \rightarrow [0, 1]$ is called a **d.f.** if

- (i) $g(\mathbf{1}) = 1$,
- (ii) $g(\mathbf{0}) = 0$, and moreover $g(\mathbf{x}) = 0$ for any \mathbf{x} with a vanishing coordinate,
- (iii) $g(\mathbf{x})$ is non-decreasing, i.e. $\Delta_{h_s}^{(s)}(\dots(\Delta_{h_1}^{(1)}g(x_1, \dots, x_s))) \geq 0$ for any $h_i \geq 0, x_i + h_i \leq 1$, where $\Delta_{h_i}^{(i)}g(x_1, \dots, x_s) = g(x_1, \dots, x_i + h_i, \dots, x_s) - g(x_1, \dots, x_i, \dots, x_s)$.

- If g is such d.f. then $\int_{[0,1]^2} dg(\mathbf{x}) = 1$.

- If $dg(\mathbf{x}) = \Delta_{dx_s}^{(s)} \dots \Delta_{dx_1}^{(1)}g(x_1, \dots, x_s)$ is the differential of $g(\mathbf{x})$ at the point $\mathbf{x} = (x_1, \dots, x_s)$, then also $dg(\mathbf{x}) = \Delta(g, J)$, where $J = [x_1, x_1 + dx_1] \times \dots \times [x_s, x_s + dx_s]$, and $\Delta(g, J)$ is an alternating sum of the values of g at the vertices of J (function values at the adjacent vertices have opposite signs), i.e.,

$$\Delta(g, J) = \sum_{\varepsilon_1=1}^2 \dots \sum_{\varepsilon_k=1}^2 (-1)^{\varepsilon_1 + \dots + \varepsilon_k} g(x_{\varepsilon_1}^{(1)}, \dots, x_{\varepsilon_k}^{(k)})$$

for an interval $J = [x_1^{(1)}, x_2^{(1)}] \times [x_1^{(2)}, x_2^{(2)}] \times \dots \times [x_1^{(k)}, x_2^{(k)}] \subset [0, 1]^k$. Moreover, $g(\mathbf{x})$ is non-decreasing if and only if $dg(\mathbf{x}) \geq 0$ for every $\mathbf{x} \in [0, 1]^s$ and $d\mathbf{x} \geq \mathbf{0}$.

- The d.f. $g(1, \dots, 1, x_{i_1}, 1, \dots, 1, x_{i_2}, 1, \dots, 1, x_{i_l}, 1, \dots, 1)$ is called an **l -dimensional face d.f. of g in variables $(x_{i_1}, x_{i_2}, \dots, x_{i_l}) \in (0, 1)^l$** , $0 \leq l \leq s$.

- We shall identify two d.f.'s $g(\mathbf{x})$ and $\tilde{g}(\mathbf{x})$ if:

- (i) $g(\mathbf{x}) = \tilde{g}(\mathbf{x})$ at every common point $\mathbf{x} \in (0, 1)^s$ of continuity, and
- (ii) $g(1, \dots, 1, x_{i_1}, 1, \dots, 1, x_{i_2}, 1, \dots, 1, x_{i_l}, 1, \dots, 1) = \tilde{g}(1, \dots, 1, x_{i_1}, 1, \dots, 1, x_{i_2}, 1, \dots, 1, x_{i_l}, 1, \dots, 1)$ at every common point $(x_{i_1}, x_{i_2}, \dots, x_{i_l}) \in (0, 1)^l$ of continuity in every l -dimensional face d.f. of g and \tilde{g} , $l = 1, 2, \dots, s$.

- The s -dimensional d.f. $g(\mathbf{x})$ is a d.f. of the sequence $\mathbf{x}_n \bmod 1$ if

(i) $g(\mathbf{x}) = \lim_{k \rightarrow \infty} F_{N_k}(\mathbf{x})$ for all continuity points $\mathbf{x} \in (0, 1)^s$ of g (so-called the **weak limit**) and,

(ii) $g(1, \dots, 1, x_{i_1}, 1, \dots, 1, x_{i_2}, 1, \dots, 1, x_{i_l}, 1, \dots, 1) =$
 $= \lim_{k \rightarrow \infty} F_{N_k}(x_{i_1}, x_{i_2}, \dots, x_{i_l})$
 weakly over $(0, 1)^l$ and every l -dimensional face sequence of \mathbf{x}_n for $l = 1, 2, \dots, s$, and for a suitable sequence of indices $N_1 < N_2 < \dots$.

• The Second Helly theorem shows that the weak limit² $F_{N_k}(\mathbf{x}) \rightarrow g(\mathbf{x})$ implies

$$\int_{[0,1]^s} f(\mathbf{x}) dF_{N_k}(\mathbf{x}) \rightarrow \int_{[0,1]^s} f(\mathbf{x}) dg(\mathbf{x})$$

for every continuous $f : [0, 1]^s \rightarrow \mathbb{R}$.

• $G(\mathbf{x}_n \bmod 1)$ denotes the set of all d.f.'s of $\mathbf{x}_n \bmod 1$.

(IV) **For one-dimensional case** $s = 1$ we have:

(1) The continuity of all d.f.s in $G(x_n \bmod 1)$ follows from the limit

$$\lim_{K \rightarrow \infty} \frac{1}{K} \sum_{k=1}^K \beta_k = 0, \text{ where } \beta_k = \limsup_{N \rightarrow \infty} \left| \frac{1}{N} \sum_{n=1}^N e^{2\pi i k x_n} \right|^2.$$

(2) The lower and upper d.f. \underline{g}, \bar{g} of x_n belong to $G(x_n \bmod 1)$ if and only if $\int_0^1 (\bar{g}(x) - \underline{g}(x)) dx = \limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \{x_n\} - \liminf_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \{x_n\}$.

(3) Let H be non-empty, closed, and connected set of d.f.'s. Denote $\underline{g}_H(x) = \inf_{g \in H} g(x)$ and $\bar{g}_H(x) = \sup_{g \in H} g(x)$. Further, if $g \in H$ let $\text{Graph}(g)$ be the continuous curve formed by all the points $(x, g(x))$ for $x \in [0, 1]$, and the all line segments connecting the points of discontinuity $(x, \liminf_{x' \rightarrow x} g(x'))$ and $(x, \limsup_{x' \rightarrow x} g(x'))$. Assume that for every $g \in H$ there exists a point $(x, y) \in \text{Graph}(g)$ such that $(x, y) \notin \text{Graph}(\tilde{g})$ for any $\tilde{g} \in H$ with $\tilde{g} \neq g$. If moreover $\underline{g} = \underline{g}_H$ and $\bar{g} = \bar{g}_H$ for the lower d.f. \underline{g} and the upper d.f. \bar{g} of the sequence $x_n \in [0, 1)$ and $G(x_n) \subset H$, then $G(x_n) = H$.

(4) For given two different d.f.s $g_1(x)$, and $g_2(x)$, we define

$$F_{g_2}(x, y) = \int_0^x g_2(t) dt + \int_0^y g_2(t) dt - \max(x, y) + \int_0^1 (1 - g_2(t))^2 dt,$$

$$F_{g_1, g_2}(x) = \frac{\int_0^x (g_2(t) - g_1(t)) dt - \int_0^1 (1 - g_2(t))(g_2(t) - g_1(t)) dt}{\int_0^1 (g_2(t) - g_1(t))^2 dt},$$

$$F_{g_1, g_2}(x, y) = F_{g_2}(x, y) - F_{g_1, g_2}(x) F_{g_1, g_2}(y) \int_0^1 (g_2(t) - g_1(t))^2 dt.$$

Let $g_1(x) \neq g_2(x)$ be two d.f.'s. Then the set of d.f.s $G(x_n)$ of x_n in $[0, 1)$ satisfies

$$G(x_n) = \{t g_1(x) + (1 - t) g_2(x); t \in [0, 1]\}$$

if and only if

²that is (i), and (ii) above are fulfilled

$$(i) \quad \lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{m,n=1}^N F_{g_1, g_2}(x_m, x_n) = 0,$$

$$(ii) \quad \liminf_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N F_{g_1, g_2}(x_n) = 0,$$

$$(iii) \quad \limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N F_{g_1, g_2}(x_n) = 1.$$

(5) A symmetric continuous $F(x, y)$ defined on $[0, 1]^2$ is called **copositive** if $\int_0^1 \int_0^1 F(x, y) dg(x) dg(y) \geq 0$ for all distribution functions $g : [0, 1] \rightarrow [0, 1]$. Let $F(x, y)$ be a copositive function having continuous $F'_x(x, 1)$ a.e. and let d.f. $g_1(x)$ be a strictly increasing solution of the moment problem (1). Then for every strictly increasing d.f. $g(x)$ we have

$$\int_0^1 \int_0^1 F(x, y) dg(x) dg(y) = 0 \iff F'_x(x, 1) = \int_0^1 g(y) d_y F'_x(x, y) \text{ a.e. on } [0, 1].$$

(6) Let $F(x, y)$ be a continuous, symmetric, copositive and $F''_{xy} = 0$ a.e. such that the set $H(F)$ of jumps of $F'_x(x, y)$ is covered by

$$H(F) \subset \bigcup_{i,j=1}^M \{(x_i(t), x_j(t)); t \in [\alpha, \beta]\} \text{ with pairwise disjoint sets } \{x_1(t); t \in [\alpha, \beta]\}, \dots, \{x_M(t); t \in [\alpha, \beta]\}.$$

Assume that the derivatives $x'_i(t)$, $i = 1, 2, \dots, k$, are continuous and let $\mathbf{A}(t)$ denote the associated matrix defined by

$$\mathbf{A}(t) = \frac{1}{2} \left(d_y F'_x(x_i(t), x_j(t)) |x'_i(t)| + d_y F'_x(x_j(t), x_i(t)) |x'_j(t)| \right) \text{ and}$$

$\mathbf{g}(t) = (g(x_1(t)), g(x_2(t)), \dots, g(x_M(t)))$ is the vector associated with $g : [0, 1] \rightarrow [0, 1]$. Finally, let g_1 be a strictly increasing solution of the moment problem (1). Then we have

$$\int_0^1 \int_0^1 F(x, y) dg(x) dg(y) = \int_\alpha^\beta (\mathbf{g}(t) - \mathbf{g}_1(t)) \mathbf{A}(t) (\mathbf{g}(t) - \mathbf{g}_1(t))^T dt$$

for all distribution functions $g : [0, 1] \rightarrow [0, 1]$.

(7) Directly by definition $G(x_n)$ we showed: Assume

- $f(x)$ be a real-valued function defined for $x \geq 1$ such that $f(x)$ is strictly increasing with its inverse function $f^{-1}(x)$.

- $\lim_{k \rightarrow \infty} \frac{f^{-1}(k+x) - f^{-1}(k)}{f^{-1}(k+1) - f^{-1}(k)} = \tilde{g}(x)$ for each $x \in [0, 1]$, point of continuity of $\tilde{g}(x)$;

- $\lim_{k \rightarrow \infty} \frac{f^{-1}(k+u)}{f^{-1}(k)} = \psi(u)$ for each $u \in [0, 1]$, point of continuity of $\psi(u)$, or $\psi(u) = \infty$ for $u > 0$;

- $\lim_{k \rightarrow \infty} f^{-1}(k+1) - f^{-1}(k) = \infty$. Then we have:

If $1 < \psi(1) < \infty$ and $f'(x) \rightarrow 0$ as $x \rightarrow \infty$, then

$$G(f(n) \bmod 1) = \left\{ g_u(x) = \frac{\min(\psi(x), \psi(u)) - 1}{\psi(u)} + \frac{1}{\psi(u)} \tilde{g}(x); u \in [0, 1] \right\},$$

where $\tilde{g}(x) = \frac{\psi(x) - 1}{\psi(1) - 1}$ and $F_{N_i}(x) \rightarrow g_u(x)$ as $i \rightarrow \infty$ if and only if $f(N_i) \bmod 1 \rightarrow u$. The lower d.f. $\underline{g}(x)$ and the upper d.f. $\bar{g}(x)$ of

$f(n) \bmod 1$ are $\underline{g}(x) = \tilde{g}(x)$, $\bar{g}(x) = 1 - \frac{1}{\psi(x)}(1 - \tilde{g}(x))$. Furthermore $\underline{g}(x) = g_0(x) = g_1(x)$ belongs to $G(f(n) \bmod 1)$ but $\bar{g}(x) = g_x(x)$ does not.

(8) Let x_n and y_n be two sequences in $[0, 1)$ and $G((x_n, y_n))$ denote the set of all d.f.s of the two-dimensional sequence (x_n, y_n) . If $z_n = x_n + y_n \bmod 1$, then the set $G(z_n)$ of all d.f.s of z_n has the form

$$G(z_n) = \left\{ g(t) = \int_{0 \leq x+y < t} 1. \, dg(x, y) + \int_{1 \leq x+y < 1+t} 1. \, dg(x, y); \right. \\ \left. g(x, y) \in G((x_n, y_n)) \right\}$$

assuming that all the used Riemann-Stieltjes integrals exist.

NOTES:

- (I) A purely topological characterization of $G(\mathbf{x}_n)$ with a short history can be found in R. Winkler (1997).
 (II) O. Strauch (1994).
 (III) For definitions, cf. [SP, 1.11, p. 1–60].
 (IV) (1) is a generalization of the Wiener – Schoenberg theorem given by P. Kostyrko, M. Mačaj, T. Šalát and O. Strauch (2001). (2), (3) and (4) are from O. Strauch (1997). (5) and (6) are proved in O. Strauch (2000), (7) in R. Giuliano Antonini and O. Strauch (2008) and (8) in O. Strauch and O. Blažeková (2006).

Proposed by O. Strauch.

R. GIULIANO ANTONINI – O. STRAUCH: *On weighted distribution functions of sequences*, Uniform Distribution Theory **3** (2008), no. 1, 1–18.

P. KOSTYRKO – M. MAČAJ – T. ŠALÁT – O. STRAUCH: *On statistical limit points*, Proc. Amer. Math. Soc. **129** (2001), no. 9, 2647–2654 (MR 2002b:40003).(electronic).

O. STRAUCH – O. BLAŽEKOVÁ: *Distribution of the sequence $p_n/n \bmod 1$* , Uniform Distribution Theory **1** (2006), 45–63.

O. STRAUCH: *L^2 discrepancy*, Math. Slovaca **44** (1994), 601–632 (MR 96c:11085).

O. STRAUCH: *On the set of distribution functions of a sequence*, in: *Proceedings of the Conference on Analytic and Elementary Number Theory in Honor of Edmund Hlawka's 80th Birthday, Vienna, July 18–20, 1996*, (W.–G. Nowak – J. Schöißengeier Eds.), Universität Wien & Universität für Bodenkultur, Vienna, 1997, pp. 214–229 (Zbl 886.11044).

O. STRAUCH: *Moment problem of the type $\int_0^1 \int_0^1 F(x, y) \, dx \, dy = 0$* , in: *Proceedings of the International Conference on Algebraic Number Theory and Diophantine Analysis held in Graz, August 30 to September 5, 1998*, (F. Halter-Koch – R.F. Tichy Eds.), Walter de Gruyter, Berlin, New York, 2000, pp. 423–443 (MR 2001d:11079).

J.G. VAN DER CORPUT: *Verteilungsfunktionen I–II*, Proc. Akad. Amsterdam **38** (1935), 813–821, 1058–1066 (JFM 61.0202.08, 61.0203.01; Zbl 12, 347; 13, 57).

J.G. VAN DER CORPUT: *Verteilungsfunktionen III–VIII*, Proc. Akad. Amsterdam **39** (1936), 10–19, 19–26, 149–153, 339–344, 489–494, 579–590 (JFM 61.0204.01, 61.0204.02, 62.0206.06, 62.0207.01, 62.0207.02, 62.0207.03).

R. WINKLER: *On the distribution behaviour of sequences*, Math. Nachr. **186** (1997), 303–312 (MR 99a:28012).

2.3 Deterministic analysis of sequences.

Assume that the s -dimensional sequence

$$\mathbf{x}_n = (x_{n,1}, \dots, x_{n,s}) \in [0, 1]^s, \quad n = 1, 2, \dots, N$$

is a result of an N often repeated measurement of s physical variables X_1, \dots, X_s . If $g(x_1, \dots, x_s)$ is an a.d.f. of \mathbf{x}_n , $n = 1, 2, \dots$ (assuming in the moment that $N \rightarrow \infty$), then $g(x_1, \dots, x_s)$ contains some informations about relations between variables X_1, \dots, X_s . For example

(i) X_1, \dots, X_s are independent if and only if every d.f. $g(\mathbf{x}) \in G(\mathbf{x}_n)$ can be written as a product $g(\mathbf{x}) = g_1(x_1) \dots g_s(x_s)$ of one-dimensional d.f.s. Here g_i , $i = 1, \dots, s$ depend on $g \in G(\mathbf{x}_n)$.

(ii) If X_s depends on X_1, \dots, X_{s-1} , and I_1, \dots, I_s are subintervals in $[0, 1)$, then the implication

$$(X_1 \in I_1 \wedge \dots \wedge X_{s-1} \in I_{s-1}) \implies (X_s \in I_s),$$

can be evaluated by $\int_{I_1 \times \dots \times I_s} h(\mathbf{x}) \, d\mathbf{x}$, where $h(\mathbf{x})$ is the density of $g(\mathbf{x})$ (if it exists).

The studying of \mathbf{x}_n , $n = 1, 2, \dots$ via $G(\mathbf{x}_n)$ we shall call **deterministic analysis** of the sequence \mathbf{x}_n , since for approximate computation of $g(\mathbf{x}) \in G(\mathbf{x}_n)$ can be used discrepancies of \mathbf{x}_n , $n = 1, 2, \dots, N$. We do not use probabilities and statistical methods.

For approximate computation of $G(\mathbf{x}_n)$ we need there solve the following problem.

Open problem. For a big dimension s there exists no real employing N for a good approximation of a d.f. $g(x_1, \dots, x_s)$ of \mathbf{x}_n by the step d.f.

$$F_N(x_1, \dots, x_s) = \frac{\#\{n \leq N; x_{n,1} < x_1, \dots, x_{n,s} < x_s\}}{N}.$$

But using partial sequences $(x_{n,i_1}, \dots, x_{n,i_k})$, $n = 1, 2, \dots, N$ with small dimension k it can be found N such that the corresponding step d.f. well approximates the marginal d.f. $g(1, \dots, 1, x_{i_1}, 1, \dots, 1, x_{i_2}, 1, \dots)$. Problem is to reconstruct $g(x_1, \dots, x_s)$ by using marginals

$$g(1, \dots, 1, x_{i_1}, 1, \dots, 1, x_{i_2}, 1, \dots)$$

with small dimensions.

In the following we shall formulate above problem more elementary.

Open problem 1. Let $\mathbf{x}_n = (x_{n,1}, x_{n,2}, \dots, x_{n,s}), n = 1, 2, \dots$, be an infinite s -dimensional sequence in the unit cube $[0, 1]^s$. Assume that, for fixed $k < s$, all k -dimensional marginal sequences $(x_{n,i_1}, \dots, x_{n,i_k})$ are u.d.

(I) Find all possible d.f.s of \mathbf{x}_n .

(II) Find some (possible "minimal") criteria which imply u.d. of the original sequence $\mathbf{x}_n, n = 1, 2, \dots$

• In connection with (I) we denote by $G_{s,k}$ the set of all d.f.s $g(\mathbf{x})$ on $[0, 1]^s$ for which all k -dimensional marginals (i.e. faces) of d.f.'s satisfy

$$g(1, \dots, 1, x_{i_1}, 1, \dots, 1, x_{i_2}, 1, \dots, 1, x_{i_k}, 1, \dots, 1) = x_{i_1} x_{i_2} \dots x_{i_k}.$$

• For $k = 1$, these d.f.'s are called *copulas*, which were introduced by M. Sklar (1959). All basic properties of copulas can be found in the monograph R.B. Nelsen (1999).

• Thus, by definition $G_{s,k}$, the $G_{2,1}$ is the set of all two-dimensional d.f.s $g(x, y)$ defined on $[0, 1]^2$ such that their marginals d.f.'s satisfy $g(x, 1) = x$ and $g(1, y) = y$. $G_{2,1}$ contains:

- $g_1(x, y) = xy$,

- $g_2(x, y) = \min(x, y)$,

- $g_3(x, y) = \max(x + y - 1, 0)$,

- $g_\theta(x, y) = (\min(x, y))^\theta (xy)^{1-\theta}$, where $\theta \in [0, 1]$ (Cuadras-Augé family, cf. Nelsen (1999, p. 12, Ex. 2.5),

- $g_4(x, y) = \frac{xy}{x+y-xy}$ (see Nelsen (1999, p. 19, 2.3.4),

- $\tilde{g}(x, y) = x + y - 1 + g(1 - x, 1 - y)$ for every $g(x, y) \in G_{2,1}$ (Survival copula, see Nelsen (1999, p. 28, 2.6.1),

- $g_5(x, y) = \min(ya(x), x(b(y)))$, where $a(0) = b(0) = 0, a(1) = b(1) = 1$ and $a(x)/x, b(y)/y$ are both decreasing on $(0, 1]$ (Marshall copula, cf. Nelsen (1999, p. 51, Exerc. 3.3).

Here are some new copulas:

- $g_6(x, y) = \frac{1}{z_0} \min(xy, xz_0, yz_0)$ for fixed $z_0, 0 < z_0 \leq 1$.

- $g_7(x, y) = \frac{1}{z_0 u_0} \min(xyz_0, xyu_0, xz_0 u_0, yz_0 u_0)$ for fixed $z_0, u_0 \in [0, 1]^2$.

- Shuffle of M is a copula defined in R.B. Nelsen (1999, p. 59, 3.2.3.), cf. the Problem 1.37.

- Generalized shuffle of M : Let $f : [0, 1] \rightarrow [0, 1]$ be an arbitrary uniform distribution preserving function (called u.d.p., see Problem 2.1 (VII)) and graph $f = \{(x, f(x)); x \in [0, 1]\}$. Then the generalized shuffle of M is the copula

$$g(x, y) = |\text{Project}_x(\text{graph } f \cap [0, x] \times [0, y])|.$$

There are some basic properties of $G_{2,1}$:

- $G_{2,1}$ is closed under point-wise limit and convex linear combinations.
- For every $g(x, y) \in G_{2,1}$ and every $(x_1, y_1), (x_2, y_2) \in [0, 1]^2$ we have $|g(x_2, y_2) - g(x_1, y_1)| \leq |x_2 - x_1| + |y_2 - y_1|$.
- For every $g(x, y) \in G_{2,1}$ we have $g_3(x, y) = \max(x + y - 1, 0) \leq g(x, y) \leq \min(x, y) = g_2(x, y)$ (Fréchet-Hoeffding bounds, see Nelsen (1999, p. 9)).
- M. Sklar (1959) proved that for every d.f. $g(x, y)$ on $[0, 1]^2$ there exists $\tilde{g}(x, y) \in G_{2,1}$ such that $g(x, y) = \tilde{g}(g(x, 1), g(1, y))$ for every $(x, y) \in [0, 1]^2$. If $g(x, 1)$ and $g(1, y)$ are continuous, then $\tilde{g}(x, y)$ is unique (cf. Nelsen (1999, p. 15, Th. 2.3.3)).
- Let $(x_n, y_n), n = 1, 2, \dots$, be a sequence in $[0, 1]^2$ such that both coordinate sequences $x_n, n = 1, 2, \dots$, and $y_n, n = 1, 2, \dots$ are u.d. Then the set $G((x_n, y_n))$ of all d.f. of $(x_n, y_n), n = 1, 2, \dots$ satisfies
 - $G((x_n, y_n)) \subset G_{2,1}$,
 - $G((x_n, y_n))$ is nonempty, closed and connected, and vice-versa
 - for every nonempty, closed and connected $H \subset G_{2,1}$, there exists a sequence $(x_n, y_n) \in [0, 1]^2$ such that $G((x_n, y_n)) = H$.
 - Let $F(x, y, u, v,)$ be a continuous function defined on $[0, 1]^4$ and assume that $\lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{m,n=1}^N F(x_m, y_m, x_n, y_n) = 0$. Then every d.f. $g(x, y) \in G((x_n, y_n))$ satisfies the following equation:

$$\begin{aligned}
 & \int_0^1 \int_0^1 g(u, v) \, d_u \, d_v \, F(1, 1, u, v) + \int_0^1 \int_0^1 g(x, y) \, d_x \, d_y \, F(x, y, 1, 1) \\
 & - \int_0^1 \int_0^1 \int_0^1 g(u, v) y \, d_y \, d_u \, d_v \, F(1, y, u, v) - \int_0^1 \int_0^1 \int_0^1 g(u, v) x \, d_x \, d_u \, d_v \, F(x, 1, u, v) \\
 & - \int_0^1 \int_0^1 \int_0^1 g(x, y) v \, d_v \, d_x \, d_y \, F(x, y, 1, v) - \int_0^1 \int_0^1 \int_0^1 g(x, y) u \, d_u \, d_x \, d_y \, F(x, y, u, 1) \\
 & + \int_0^1 \int_0^1 \int_0^1 \int_0^1 g(x, y) g(u, v) \, d_u \, d_v \, d_x \, d_y \, F(x, y, u, v) = \\
 & = -F(1, 1, 1, 1) + \int_0^1 v \, d_v \, F(1, 1, 1, v) + \int_0^1 u \, d_u \, F(1, 1, u, 1) \\
 & + \int_0^1 x \, d_x \, F(x, 1, 1, 1) + \int_0^1 y \, d_y \, F(1, y, 1, 1) \\
 & - \int_0^1 \int_0^1 y v \, d_y \, d_v \, F(1, y, 1, v) - \int_0^1 \int_0^1 y u \, d_y \, d_u \, F(1, y, u, 1) \\
 & - \int_0^1 \int_0^1 x v \, d_x \, d_v \, F(x, 1, 1, v) - \int_0^1 \int_0^1 x u \, d_x \, d_u \, F(x, 1, u, 1).
 \end{aligned}$$


- By definition of $G_{s,k}$, the $G_{3,2}$ is the set of all three-dimensional d.f.s $g(x, y, z)$ defined on $[0, 1]^3$ such that their two-dimensional marginals (or faces) d.f.'s satisfy $g(x, y, 1) = xy$, $g(1, y, z) = yz$ and $g(x, 1, z) = xz$. The

$G_{3,2}$ contains

- $g_1(x, y, z) = xyz$,
- $g_2(x, y, z) = \min(xy, xz, yz)$,
- $g_3(x, y, z) = \frac{1}{u_0} \min(xyz, xyu_0, xzu_0, yzu_0)$, for fixed u_0 , $0 < u_0 \leq 1$,
- $g_4(x, y, z)$ is a.d.f. of a three-dimensional sequence $(u_n, v_n, \{u_n - v_n\})$, where two-dimensional (u_n, v_n) is u.d. in $[0, 1]^2$. Applying Weyl's criterion we see that also $(u_n, \{u_n - v_n\})$ and $(v_n, \{u_n - v_n\})$ are u.d. and the d.f. $g_4(x, y, z)$ has the following explicit form (cf. O. Strauch (2003).)

$$g_4(x_1, x_2, x_3) = \begin{cases} x_1x_2, & \text{if } (x_1, x_2) \in A, \\ -\frac{1}{2}(x_1^2 + x_2^2 + x_3^2) + x_1x_2 + x_2x_3, & \text{if } (x_1, x_2) \in B, \\ -\frac{1}{2}x_1^2 + x_1x_2, & \text{if } (x_1, x_2) \in C, \\ \frac{1}{2}x_2^2, & \text{if } (x_1, x_2) \in D, \\ -\frac{1}{2}x_3^2 + x_2x_3, & \text{if } (x_1, x_2) \in E, \\ -\frac{1}{2}x_2^2 + x_1x_2 + x_1x_3 + x_2x_3 - x_1 - x_3 + \frac{1}{2}, & \text{if } (x_1, x_2) \in F, \\ \frac{1}{2}x_1^2 + x_1x_3 + x_2x_3 - x_1 - x_3 + \frac{1}{2}, & \text{if } (x_1, x_2) \in G, \\ \frac{1}{2}(x_1^2 + x_2^2 + x_3^2) + x_1x_3 - x_1 - x_3 + \frac{1}{2}, & \text{if } (x_1, x_2) \in H, \\ x_1x_2 + x_2x_3 - x_2 & \text{if } (x_1, x_2) \in I. \end{cases}$$

where the regions $A, B, C, D, E, F, G, H, I$ are shown on the following Figure



- For every $g(x, y, z) \in G_{3,2}$ and fixed z_0 , $0 < z_0 \leq 1$ we have $\frac{1}{z_0}g(x, y, z_0) \in G_{2,1}$. Vice versa, if $g_z(x, y)$, $z \in [0, 1]$ is a system of d.f.s in $G_{2,1}$ such that

$g_1(x, y) = xy$ and for every $z' \leq z$, we have $z' d_x d_y g_{z'}(x, y) \leq z d_x d_y g_z(x, y)$ on $[0, 1]^2$, then $g(x, y, z) = z g_z(x, y) \in G_{3,2}$.

- Multi-dimensional case: Let $g(x_1, x_2, \dots, x_s)$ be an s -dimensional d.f. and let $g_1(x_1) = g(x_1, 1, \dots, 1)$, $g_2(x_2) = g(1, x_2, 1, \dots, 1), \dots$ be margins of $g(x_1, x_2, \dots, x_s)$. By Sklar's theorem there exists s -dimensional copula $c(x_1, x_2, \dots, x_s)$ such that

$$g(x_1, x_2, \dots, x_s) = c(g_1(x_1), g_2(x_2), \dots, g_s(x_s)).$$

Furthermore, for arbitrary continuous $F(x_1, x_2, \dots, x_s)$ we have

$$\begin{aligned} & \int_{[0,1]^s} F(x_1, x_2, \dots, x_s) dg(x_1, x_2, \dots, x_s) \\ &= \int_{[0,1]^s} F(g_1^{-1}(x_1), g_2^{-1}(x_2), \dots, g_s^{-1}(x_s)) dc(x_1, x_2, \dots, x_s). \end{aligned}$$

• In the direction (II) of Open problem for testing of u.d. of \mathbf{x}_n it can be used statistical independence of marginal sequences, but formulas for L^2 discrepancy of statistical independence and classical L^2 discrepancy have the following similar structures: For $\mathbf{x} = (x_1, \dots, x_s)$ and $\mathbf{y} = (y_1, \dots, y_s)$ denote $(\mathbf{1} - \max(\mathbf{x}, \mathbf{y})) = (1 - \max(x_1, y_1)) \dots (1 - \max(x_s, y_s))$, $\mathbf{0} = (0, \dots, 0)$ and $\mathbf{1} = (1, \dots, 1)$. For every two d.f.'s $g_1(\mathbf{x})$ and $g_2(\mathbf{x})$ defined in $[0, 1]^s$ we have (see Strauch (2003))

$$\int_{\mathbf{0}}^{\mathbf{1}} (g_1(\mathbf{x}) - g_2(\mathbf{x}))^2 d\mathbf{x} = \int_{\mathbf{0}}^{\mathbf{1}} \int_{\mathbf{0}}^{\mathbf{1}} (\mathbf{1} - \max(\mathbf{x}, \mathbf{y})) d(g_1(\mathbf{x}) - g_2(\mathbf{x})) d(g_1(\mathbf{y}) - g_2(\mathbf{y})) \quad (1)$$

Now, divide the vector $\mathbf{x} = (x_1, \dots, x_s)$ into two face vectors $\mathbf{x}^{(1)} = (x_{i_1}, \dots, x_{i_l})$ and $\mathbf{x}^{(2)} = (x_{j_1}, \dots, x_{j_k})$, $l + k = s$. Similarly, divide the s -dimensional sequence \mathbf{x}_n , $n = 1, 2, \dots$ in $[0, 1]^s$ with step d.f. $F_N(\mathbf{x}) = F_N(\mathbf{x}^{(1)}, \mathbf{x}^{(2)})$ into two face sequences

l -dimensional $\mathbf{x}_n^{(1)}$, $n = 1, 2, \dots$, with step d.f. $F_N(\mathbf{x}^{(1)}, \mathbf{1})$, and k -dimensional $\mathbf{x}_n^{(2)}$, $n = 1, 2, \dots$, with step d.f. $F_N(\mathbf{1}, \mathbf{x}^{(2)})$.

Using (1) we see that the L^2 discrepancy (with respect to $g(\mathbf{x})$) and statistical L^2 discrepancy have the following similar structures

$$\int_{\mathbf{0}}^{\mathbf{1}} (F_N(\mathbf{x}) - g(\mathbf{x}))^2 d\mathbf{x} = \int_{\mathbf{0}}^{\mathbf{1}} \int_{\mathbf{0}}^{\mathbf{1}} (\mathbf{1} - \max(\mathbf{x}, \mathbf{y})) \cdot d(F_N(\mathbf{x}) - g(\mathbf{x})) d(F_N(\mathbf{y}) - g(\mathbf{y})),$$

$$\begin{aligned} & \int_{\mathbf{0}}^{\mathbf{1}} (F_N(\mathbf{x}) - F_N(\mathbf{x}^{(1)}, \mathbf{1}) F_N(\mathbf{1}, \mathbf{x}^{(2)}))^2 d\mathbf{x} = \\ &= \int_{\mathbf{0}}^{\mathbf{1}} \int_{\mathbf{0}}^{\mathbf{1}} \int_{\mathbf{0}}^{\mathbf{1}} \int_{\mathbf{0}}^{\mathbf{1}} (\mathbf{1} - \max(\mathbf{x}^{(1)}, \mathbf{y}^{(1)})) (\mathbf{1} - \max(\mathbf{x}^{(2)}, \mathbf{y}^{(2)})) \cdot \\ & \cdot d(F_N(\mathbf{x}^{(1)}, \mathbf{x}^{(2)}) - F_N(\mathbf{x}^{(1)}, \mathbf{1}) F_N(\mathbf{1}, \mathbf{x}^{(2)})) \cdot \\ & \cdot d(F_N(\mathbf{y}^{(1)}, \mathbf{y}^{(2)}) - F_N(\mathbf{y}^{(1)}, \mathbf{1}) F_N(\mathbf{1}, \mathbf{y}^{(2)})). \end{aligned}$$

Expressing L^2 discrepancy as

$$\begin{aligned} & \int_{\mathbf{0}}^{\mathbf{1}} (F_N(\mathbf{x}) - g(\mathbf{x}))^2 d\mathbf{x} = \\ & = \int_{\mathbf{0}}^{\mathbf{1}} \int_{\mathbf{0}}^{\mathbf{1}} \left[(F_N(\mathbf{x}^{(1)}, \mathbf{x}^{(2)}) - F_N(\mathbf{x}^{(1)}, \mathbf{1})F_N(\mathbf{1}, \mathbf{x}^{(2)})) + \right. \\ & + (F_N(\mathbf{x}^{(1)}, \mathbf{1}) - g(\mathbf{x}^{(1)}, \mathbf{1}))F_N(\mathbf{1}, \mathbf{x}^{(2)}) + \\ & + g(\mathbf{x}^{(1)}, \mathbf{1})(F_N(\mathbf{1}, \mathbf{x}^{(2)}) - g(\mathbf{1}, \mathbf{x}^{(2)})) + \\ & \left. + (g(\mathbf{x}^{(1)}, \mathbf{1})g(\mathbf{1}, \mathbf{x}^{(2)}) - g(\mathbf{x}^{(1)}, \mathbf{x}^{(2)})) \right]^2 d\mathbf{x}^{(1)} d\mathbf{x}^{(2)} \end{aligned}$$

the Cauchy inequality implies

$$\begin{aligned} & \sqrt{\int_{\mathbf{0}}^{\mathbf{1}} (F_N(\mathbf{x}) - g(\mathbf{x}))^2 d\mathbf{x}} \leq \\ & \leq \sqrt{\int_{\mathbf{0}}^{\mathbf{1}} \int_{\mathbf{0}}^{\mathbf{1}} (F_N(\mathbf{x}^{(1)}, \mathbf{x}^{(2)}) - F_N(\mathbf{x}^{(1)}, \mathbf{1})F_N(\mathbf{1}, \mathbf{x}^{(2)}))^2 d\mathbf{x}^{(1)} d\mathbf{x}^{(2)} +} \\ & + \sqrt{\int_{\mathbf{0}}^{\mathbf{1}} \int_{\mathbf{0}}^{\mathbf{1}} (g(\mathbf{x}^{(1)}, \mathbf{x}^{(2)}) - g(\mathbf{x}^{(1)}, \mathbf{1})g(\mathbf{1}, \mathbf{x}^{(2)}))^2 d\mathbf{x}^{(1)} d\mathbf{x}^{(2)} +} \\ & + \sqrt{\int_{\mathbf{0}}^{\mathbf{1}} (F_N(\mathbf{x}^{(1)}, \mathbf{1}) - g(\mathbf{x}^{(1)}, \mathbf{1}))^2 d\mathbf{x}^{(1)} \int_{\mathbf{0}}^{\mathbf{1}} F_N^2(\mathbf{1}, \mathbf{x}^{(2)}) d\mathbf{x}^{(2)} +} \\ & + \sqrt{\int_{\mathbf{0}}^{\mathbf{1}} (F_N(\mathbf{1}, \mathbf{x}^{(2)}) - g(\mathbf{1}, \mathbf{x}^{(2)}))^2 d\mathbf{x}^{(2)} \int_{\mathbf{0}}^{\mathbf{1}} g^2(\mathbf{x}^{(1)}, \mathbf{1}) d\mathbf{x}^{(1)}}. \end{aligned}$$

Thus we have an upper bound of the classical L^2 discrepancy of $\mathbf{x}_1, \dots, \mathbf{x}_N$ which contains the L^2 discrepancy of statistical independence of partial sequences $\mathbf{x}_1^{(1)}, \dots, \mathbf{x}_N^{(1)}$ and $\mathbf{x}_1^{(2)}, \dots, \mathbf{x}_N^{(2)}$. Note that the infinite partial sequences $\mathbf{x}_n^{(1)}$, $n = 1, 2, \dots$, and $\mathbf{x}_n^{(2)}$, $n = 1, 2, \dots$ of the sequence \mathbf{x}_n , $n = 1, 2, \dots$ are statistically independent if and only if for every d.f. $g(\mathbf{x}) \in G(\mathbf{x}_n)$ we have $g(\mathbf{x}) = g(\mathbf{x}^{(1)}, \mathbf{1}) \cdot g(\mathbf{1}, \mathbf{x}^{(2)})$ in common points of continuity of d.f.s. It can be used as a definition of independence.

Proposed by O. Strauch.

R.B. NELSEN: *An Introduction to Copulas. Properties and Applications*, Lecture Notes in Statistics, Vol. 139, Springer-Verlag, New York, 1999.

M. SKLAR: *Fonctions de répartition à n dimensions et leurs marges*, Publ. Inst. Statist. Univ. Paris **8** (1959), 229–231 (MR **23**# A2899).

O. STRAUCH: *Reconstruction of distribution function by its marginals*, Math. Institute, Slovak Acad. Sci., Bratislava, Slovak Republic, 2003, 10 pp.

2.4 Exponential sequences.

The theory of the sequences $\lambda\theta^n \bmod 1$, $n = 1, 2, \dots$, $\theta > 1$ is not satisfactory. Characterization of distribution of such sequences is a well-known

and largely **unsolved problem**, see [SP, p. 2–149]. In the following we listed some conjectures and some positive results.

NOTES:

(1) J.F. Koksma (1935) proved that the sequence $\lambda\theta^n \bmod 1$ with $\lambda \neq 0$ fixed is u.d. for almost all real $\theta > 1$. If we take $\lambda = 1$ then we get that the sequence $\theta^n \bmod 1$ is u.d. for almost all real numbers $\theta > 1$. However, no explicit example of a real number θ is known for which this sequence is u.d.

(2) If $\theta > 1$ is fixed then H. Weyl (1916) proved that the sequence $\lambda\theta^n \bmod 1$ is u.d. for almost all real λ .

(3) A.D. Pollington (1983) proved that the Hausdorff dimension of the set of all $\lambda \in \mathbb{R}$ for which the sequence $\lambda\theta^n \bmod 1$ is nowhere dense is $\geq \frac{1}{2}$.

(4) $(3/2)^n \bmod 1$ is u.d. in $[0, 1]$ (**conjecture**).

(5) $(3/2)^n \bmod 1$ is dense in $[0, 1]$ (**conjecture**).

(6) $\limsup_{n \rightarrow \infty} \{(3/2)^n\} - \liminf_{n \rightarrow \infty} \{(3/2)^n\} > 1/2$ (T. Vijayaraghavan's (1940) **conjecture**).

(7) K. Mahler's (1968) **conjecture**: There is no $0 \neq \xi \in \mathbb{R}$ such that $0 \leq \{\xi(3/2)^n\} < 1/2$ for all $n = 0, 1, 2, \dots$. Such ξ does not exist if for each $\xi > 0$ the sequence of integer parts $[\xi(3/2)^n]$, $n = 1, 2, \dots$, contains infinitely many odd numbers.

(8) There is no $0 \neq \xi \in \mathbb{R}$ such that the closure of $\{\{\xi(3/2)^n\}; n = 0, 1, 2, \dots\}$ is nowhere dense in $[0, 1]$ (**conjecture**).

(9) L. Flatto, J.C. Lagarias and A.D. Pollington (1995) showed that for every $\xi > 0$ we have $\limsup_{n \rightarrow \infty} \{\xi(3/2)^n\} - \liminf_{n \rightarrow \infty} \{\xi(3/2)^n\} \geq 1/3$.

(10) G. Choquet (1980) proved the existence of infinitely many $\xi \in \mathbb{R}$ for which $1/19 \leq \{\xi(3/2)^n\} \leq 18/19$ for $n = 0, 1, 2, \dots$. Him is ascribed the conjecture (v).

(11) A. Dubickas (2006[a]) proved that for any $\xi \neq 0$ the sequence of fractional part $\{\xi(3/2)^n\}$, $n = 1, 2, \dots$, contains at least one limit point in the interval $[0.238117\dots, 0.761882\dots]$ of length $0.523764\dots$. This immediately follows from:

(12) A. Dubickas (2006[a]): Set $T(x) = \prod_{n=0}^{\infty} (1 - x^{2^n})$. If $\xi \neq 0$ then the sequence $\|\xi(3/2)^n\|$, $n = 1, 2, \dots$, has a limit point $\geq (3 - T(2/3))/12 = 0.238117\dots$ and a limit point $\leq (1 + T(2/3))/4 = 0.285647\dots$.

(12') A. Dubickas (2007) from (22') derived: $\{\xi(-3/2)^n\}$ has a limit point ≤ 0.533547 and a limit point ≥ 0.466452 .

(13) S. Akiyama, C. Frougny and J. Sakarovitch (2006): There is $\xi \neq 0$ such that $\|\xi(3/2)^n\| < 1/3$ for $n = 1, 2, \dots$.

(14) A. Pollington: There is $\xi \neq 0$ such that $\|\xi(3/2)^n\| > 4/65$ for $n = 1, 2, \dots$.

(15) R. Tijdeman (1972) showed that for every pair of integers k, m with $k \geq 2$ and $m \geq 1$ there exists $\xi \in [m, m+1)$ such that $0 \leq \{\xi((2k+1)/2)^n\} \leq 1/(2k-1)$ for $n = 0, 1, 2, \dots$.

(16) O. Strauch (1997) proved that every d.f. $g(x)$ of $\xi(3/2)^n \bmod 1$ satisfies the functional equation

$$g(x/2) + g((x+1)/2) - g(1/2) = g(x/3) + g((x+1)/3) + g((x+2)/3) - g(1/3) - g(2/3).$$

A non-trivial solution (cf. O. Strauch (1999, p. 126)) is

$$g(x) = \begin{cases} 0, & \text{if } x \in [0, 1/6], \\ 2x - 1/3, & \text{if } x \in [1/6, 3/12], \\ 4x - 5/6, & \text{if } x \in [3/12, 5/18], \\ 2x - 5/18, & \text{if } x \in [5/18, 2/6], \\ 7/18, & \text{if } x \in [2/6, 8/18], \\ x - 1/18, & \text{if } x \in [8/18, 3/6], \\ 8/18, & \text{if } x \in [3/6, 7/9], \\ 2x - 20/18, & \text{if } x \in [7/9, 5/6], \\ 4x - 50/18, & \text{if } x \in [5/6, 11/12], \\ 2x - 17/18, & \text{if } x \in [11/12, 17/18], \\ x, & \text{if } x \in [17/18, 1]. \end{cases}$$

(17) Strauch (1997) introduced : The set $X \subset [0, 1]$ is said to be a set of uniqueness of d.f.s of $\xi(3/2)^n \bmod 1$, if for every two d.f.s $g_1(x), g_2(x)$ of $\xi(3/2)^n \bmod 1$ with $g_1(x) = g_2(x)$ for $x \in X$ then $g_1(x) = g_2(x)$ for every $x \in [0, 1]$. He gives the following sets of uniqueness: $X = [0, 2/3]$, $X = [1/3, 1]$, $X = [0, 1/3] \cup [2/3, 1]$, $X = [2/9, 1/3] \cup [1/2, 1]$ or $X = [0, 1/2] \cup [2/3, 7/9]$.

(18) The elements of the sequence $(3/2)^n$ appear in the Waring problem. Let

$$g(k) = \min \{s; a = n_1^k + \dots + n_s^k \text{ for all } a \in \mathbb{N} \text{ and suitable } n_i \in \mathbb{N}_0\}.$$

S. Pillai (1936) proved that if $k \geq 5$ and if we write $3^k = q2^k + r$ with $0 < r < 2^k$, then $g(k) = 2^k + [(3/2)^k] - 2$, provided that $r + q < 2^k$, i.e. $3^k - 2^k[(3/2)^k] < 2^k - [(3/2)^k]$.

(19) **Open problem** is to characterize distribution of $e^n \bmod 1$ and $\pi^n \bmod 1$.

(20) If $p > q > 1$ are integers and $\gcd(p, q) = 1$ then the sequence $(p/q)^n \bmod 1$, $n = 1, 2, \dots$, has an infinite number of points of accumulation. This was firstly proved by Ch. Pisot (1938), then by T. Vijayaraghavan (1940) and L. Rédei (1942). The density of $(p/q)^n \bmod 1$ in $[0, 1]$ is an **open problem** posed by Pisot and Vijayaraghavan.

(21) L. Flatto, J.C. Lagarias and A.D. Pollington (1995) proved that if $\xi > 0$, then $\limsup_{n \rightarrow \infty} \{\xi(p/q)^n\} - \liminf_{n \rightarrow \infty} \{\xi(p/q)^n\} \geq 1/p$.

(22) A. Dubickas (2006[a]): Denote $T(x) = \prod_{n=0}^{\infty} (1 - x^{2^n})$, $E(x) = \frac{1 - (1-x)T(x)}{2x}$. If $\xi \neq 0$ and $p > q > 1$, $\gcd(p, q) = 1$, then the sequence $\|\xi(p/q)^n\|$, $n = 1, 2, \dots$ has a limit point $\geq E(q/p)/p$ and a limit point $\leq 1/2 - (1 - e(q/p))T(q/p)/2q$, where $e(q/p) = 1 - (q/p)$ if $p + q$ is even and $e(q/p) = 1$ if $p + q$ is odd.

(22') A. Dubickas (2007): Set $F(x) = \prod_{k=1}^{\infty} (1 - x^{(2^k + (-1)^{k-1})/3})$. For two coprime positive integers $p > q > 1$ and any real number $\xi \neq 0$, the sequence of fractional part $\{\xi(-p/q)^n\}$, $n = 0, 1, 2, \dots$, has a limit point $\leq 1 - (1 - F(q/p))/q$ and a limit point $\geq (1 - F(q/p))/q$.

(22'') A. Dubickas (2006[a]): Set $T(x) = \prod_{n=0}^{\infty} (1 - x^{2^n})$ and $E(x) = \frac{1 - (1-x)T(x)}{2x}$. Let ξ be an irrational number and let $p > 1$ be an integer. Then the sequence

$\|\xi p^n\|$, $n = 1, 2, \dots$ has a limit point $\geq \xi_p = E(1/p)/p$, and a limit point $\leq \hat{\xi}_p = e(1/p)T(1/p)/2$, where $e(1/p) = 1 - (1/p)$ if p is odd and $e(1/p) = 1$ if p is even. Furthermore, both bounds are best possible: in particular, $\xi_p, \hat{\xi}_p$ are irrational and $\|\xi_p p^n\| < \xi_p, \|\hat{\xi}_p p^n\| > \hat{\xi}_p$ for every $n = 1, 2, \dots$

(23) S.D. Adhikari, P. Rath and N. Saradha (2005) prove that every d.f. $g(x)$ of $\{\xi(p/q)^n\}$ satisfies the functional equation

$$\sum_{i=0}^{q-1} g\left(\frac{x+i}{q}\right) - \sum_{i=0}^{q-1} g\left(\frac{i}{q}\right) = \sum_{i=0}^{p-1} g\left(\frac{x+i}{p}\right) - \sum_{i=0}^{p-1} g\left(\frac{i}{p}\right).$$

(24) S.D. Adhikari, P. Rath and N. Saradha (2005) prove that every interval $I \subset [0, 1]$ of the length $|I| = (p-1)/q$ and every complement $[0, 1] - [(i-1)/p, i/p]$, $i = 1, 2, \dots, p$, are sets of uniqueness of d.f.s of $\{\xi(p/q)^n\}$, for definition see (17). In the second case, if $j/q \in [(i-1)/p, i/p]$ for some $1 \leq j < q$ they assume $p \geq q^2 - q$.

(25) T. Vijayaraghavan (1940a): Let $\theta = q^{\frac{1}{k}}$ be irrational, where k and $q \geq 2$ are integers. Then the set of limit points of the sequence $\theta^n \bmod 1$ is infinite.

(26) H. Helson and J.-P. Kahane (1965): Let $\theta > 1$ be a real number. There exists uncountably many ξ such that the sequence $\xi\theta^n \bmod 1$ does not have the a.d.f.

(27) A. Zame (1967): For an arbitrary d.f. $g(x)$ and for any sequence u_n of real numbers which satisfies $\lim_{n \rightarrow \infty} (u_{n+1} - u_n) = \infty$, there exists a real number θ such that the sequence $\theta^{u_n} \bmod 1$ has $g(x)$ as its a.d.f.

• A real algebraic integer $\theta > 1$ is called a P.V. number (Pisot – Vijayaraghavan number) if all its conjugates $\neq \theta$ lie strictly inside the unit circle.

(28) Let θ be a P.V. number. Then $\theta^n \bmod 1 \rightarrow 0$ as $n \rightarrow \infty$.

(29) A. Thue (1912) proved that θ is a P.V. number if and only if $\{\theta^n\} = \mathcal{O}(c^n)$ for some $0 < c < 1$.

(30) G.H. Hardy (1919) proved that if $\theta > 1$ is any algebraic number and $\lambda > 0$ is a real number so that $\{\lambda\theta^n\} = \mathcal{O}(c^n)$, ($0 < c < 1$), then θ is a P.V. number. Hardy posed an interesting and still **unanswered question** whether there is a transcendental numbers $\theta > 1$ for which a $\lambda > 0$ exists such that $\{\lambda\theta^n\} \rightarrow 0$.

(31) T. Vijayaraghavan (1941) proved that if $\theta > 1$ is an algebraic and if θ^n , $n = 1, 3, \dots$, has only a finite set of limit points, then θ is a P.V. number.

(32) Ch. Pisot (1937, [a]1937) proved that if $\theta > 1$ and $\lambda > 0$ are real numbers such that $\sum_{n=1}^{\infty} \{\lambda\theta^n\} < +\infty$, then θ is a P.V. number.

(33) The set S of all P.V. numbers is closed (R. Salem (1944)). Two smallest elements of S are $1.324717\dots$, and $1.380277\dots$, the real roots of $x^3 - x - 1$, and $x^4 - x^3 - 1$, respectively. Both are isolated points of S and S contains no other point in the interval $(1, \sqrt{2}]$ (C.L. Siegel (1944)). The next one is $1.443269\dots$, the real root of $x^5 - x^4 - x^3 + x^2 - 1$ and $1.465571\dots$, the real root of $x^3 - x^2 - 1$. The smallest limit point of S is the root $\frac{(1+\sqrt{5})}{2} = 1.618033\dots$ of $x^2 - x - 1$, an isolated point of the derived set S' of S (J. Dufresnoy and Ch. Pisot (1952), (1953)). The smallest number S'' is 2.

• The real algebraic integer $\theta > 1$ is called a Salem number if all its conjugates lie inside or on the circumference of the unit circle and at least one of conjugates of θ lies on the circumference of the unit circle. It is well known that if θ is a Salem number of degree d , then d is even, $d \geq 4$ and $1/\theta$ is the only conjugate of θ with modulus less than 1, all the other conjugates are of modulus 1.

(34) Let θ be a Salem number. the sequence $\theta^n \bmod 1$ is dense in $[0, 1]$, but not u.d. (Ch. Pisot and R. Salem (1964)) Salem numbers are the only known concrete numbers whose powers are dense mod 1 in $[0, 1]$, see the monograph of M.J. Bertin, A. Decomps-Guilloux, M. Grandet-Hugot, M. Pathiaux-Delefosse, and J.P. Schreiber (1992, pp. 87–89). The survey paper of E. Ghate and E. Hironaka (2001) deals with the following **open problem**: Is the set of Salem numbers bounded away from 1? D.H. Lehmer (1933) found the monic polynomial $L(x) = x^{10} + x^9 - x^7 - x^6 - x^5 - x^4 - x^3 + x + 1$, where its real root $\theta = 1.17628\dots$ is both the smallest known Salem number.

(35) Let θ be the Salem numbers of degree greater than or equal to 8. Then the sequence $x_n = \theta^n \bmod 1$, $n = 1, 2, \dots$, has a.d.f. $g(x) \neq x$ which satisfies $|(g(y) - g(x)) - (y - x)| \leq 2\zeta\left(\frac{\deg(\theta)-2}{4}\right)(2\pi)^{1-\frac{\deg(\theta)}{2}}(y - x)$, where $\zeta(z)$ is the Riemann zeta function, $\deg(\theta)$ is the degree of θ over \mathbb{Q} and $0 \leq x < y \leq 1$. This was proved by S. Akiyama and Y. Tanigawa (2004).

(35') Toufik Zaimi (2006): Let θ be a Salem number and let λ be a nonzero element of the field $\mathbb{Q}(\theta)$ and denote $\Delta = \limsup_{n \rightarrow \infty} \{\lambda\theta^n\} - \liminf_{n \rightarrow \infty} \{\lambda\theta^n\}$. Then (i) $\Delta > 0$. (ii) If λ is an algebraic integer, then $\Delta = 1$. Furthermore, for any $0 < t < 1$ there is an algebraic integer λ and a subinterval $I \subset [0, 1]$ with the length t such that the sequence $\{\lambda\theta^n\}$, $n = 1, 2, \dots$ has no limit point in I . (iii) If $\theta - 1$ is a unit, then $\Delta \geq 1/L$, where L is the sum of the absolute values of the coefficients of the minimal polynomial of θ . (iv) If $\theta - 1$ is not a unit, then $\inf_{\lambda} \Delta = 0$.

(35'') A. Dubickas (2006[b]): If θ is either a P.V. or Salem number and $\lambda \neq 0$ and $\lambda \notin \mathbb{Q}(\theta)$, then $\Delta \geq 1/L$, where Δ and λ are defined as in (35').

(35''') A. Dubickas (2006[b], see MR2201605=2006i:11080): Let $d \geq 2$ be a positive integer. Suppose that $\alpha > 1$ is a root of the polynomial $x^d - x - 1$. Let ξ be an arbitrary positive number that lies outside the field $\mathbb{Q}(\alpha)$ if $d = 2$ or $d = 3$. Then the sequence $[\xi\alpha^n]$, $n = 1, 2, \dots$, contains infinitely many even numbers and infinitely many odd numbers. Thus α satisfies Mahler's conjecture (7), i.e. $0 \leq \{\xi\alpha^n\} < 1/2$, does not holds for all $n = 1, 2, \dots$.

(35''') D. Berend and G. Kolesnik (2011): If λ is a Salem number of degree 4, then the sequence $n\lambda^n \bmod 1$, $n = 1, 2, \dots$ is u.d. Precisely they proved: Let λ be a Salem number of degree 4 and $P(x)$ a nonconstant polynomial with integer coefficients. Then the sequence

$(P(n)\lambda^n, P(n+1)\lambda^{n+1}, P(n+2)\lambda^{n+2}, P(n+3)\lambda^{n+3}) \bmod 1, n = 1, 2, \dots$ is u.d.

(36) I.I. Pjateckiĭ – Šapiro (1951) proved that every distribution function $g(x)$ of the sequence $\alpha q^n \bmod 1$ with integer $q > 1$ satisfies the functional equation

$$g(x) = \sum_{i=0}^{q-1} (g((x+i)/q) - g(i/q)).$$

(37) If α is a non-zero real number and $q \geq 2$ an integer then the sequence $\alpha q^n \bmod 1$ has a.d.f. $g(x)$ if and only if $\int_0^1 f(x) dg(x) = \int_0^1 f(qx) dg(x)$ for every continuous $f(x)$ which is defined on $[0, 1]$. (I.I. Pjateckiĭ – Šapiro (1951)).

(38) Let α be a non-zero real and $q \geq 2$ be an integer. If the sequence $x_n = \alpha q^n \bmod 1$ has absolutely continuous a.d.f. $g(x)$, then $g(x) = x$ and thus the sequence x_n is u.d.

(39) If α is irrational, then for any integer $q \geq 2$ the set of all limit points of the

sequence $\alpha q^n \bmod 1$ is infinite (T. Vijayaraghavan (1940a)).

(39') A. Dubickas (2007): For an integer $b \leq -2$ and any irrational ξ we have $\liminf_{n \rightarrow \infty} \{\xi b^n\} \leq F(-1/b)/q$ and $\limsup_{n \rightarrow \infty} \{\xi b^n\} \geq (1 - F(q/p))/q$, where $F(x) = \prod_{k=1}^{\infty} (1 - x^{2^k + (-1)^{k-1}/3})$. From it he derives: (i) $\liminf_{n \rightarrow \infty} \{\xi(-2)^n\} < 0.211811$ and $\limsup_{n \rightarrow \infty} \{\xi(-2)^n\} > 0.788189$; (ii) The sequence of integer parts $[\xi(-2)^n]$, $n = 0, 1, 2, \dots$, contains infinitely many numbers divisible by 3 and infinitely many numbers divisible by 4.

• The number α is normal in the base q if and only if $\alpha q^n \bmod 1$ is u.d. The number α is called absolutely normal if it is normal in the base q for all integers $q \geq 2$. The number α is called simply normal to base q if each digit from 0 to $q - 1$ appears with the asymptotic frequency $\frac{1}{q}$.

(40) **It is not known** whether the following constants of general interest e , π , $\sqrt{2}$, $\log 2$, $\zeta(3)$, $\zeta(5)$, .. are normal in the base 10. All are, conjecturally, absolutely normal.

(41) The first classical example $\alpha_0 = 0.123456789101112\dots$ of a simple normal number in base $q = 10$ is given by Champernowne (1933). It is also normal in $q = 10$.

(42) Let $f(x) = \alpha_0 x^{\beta_0} + \alpha_1 x^{\beta_1} + \dots + \alpha_k x^{\beta_k}$ be a generalized polynomial where α 's and β 's are real numbers such that $\beta_0 > \beta_1 > \dots > \beta_k \geq 0$. Assume that $f(x) \geq 1$ for $x \geq 1$ and that $q \geq 2$ is a fixed integer. Put $\alpha = 0.[f(1)][f(2)]\dots$, where the integer part $[f(n)]$ is represented in the q -adic digit expansion. Then α is normal in the base q . This was proved by Y.-N. Nakai and I. Shiokawa in the series of papers (1990, [a]1990, 1992). They give the following examples $\alpha = 0.1247912151822\dots$ with $f(x) = x^{\sqrt{2}}$, and $\alpha = 0.151222355069\dots$ with $f(x) = \sqrt{2}x^2$.

(43) If $f(x)$ is a non-constant polynomial with rational coefficients all of whose values at $x = 1, 2, \dots$, are positive integers, then the normality of α in base 10 was proved by H. Davenport and P. Erdős (1952).

(44) K. Mahler (1937) proved that α defined by an integer polynomial $f(x)$ is a transcendental number of the non-Liouville type.

(45) Y.-N. Nakai and I. Shiokawa (1997): Let $f(x)$ be a non-constant polynomial which takes positive integral values at all positive integers. The number $\alpha = 0.f(2)f(3)f(5)f(7)f(11)\dots$, where $f(p)$ is represented in the q -adic digit expansion and p runs through the primes, is normal in the integral base q . The normality of $\alpha = 0.235711\dots$ with respect to base $q = 10$ was conjectured by D.G. Champernowne (1933) and proved by A.H. Copeland and P. Erdős (1946).

(45') M.G. Madritsch, J.M. Thuswaldner and R.F. Tichy (2008) extended the results of Nakai and Shiokawa by showing that, if f is an entire function of logarithmic order, then the numbers $0.[f(1)][f(2)][f(3)]\dots$ and $0.[f(2)][f(3)][f(5)][f(7)]\dots$, where $[f(n)]$ stands for the base q expansion of the integer part of $f(n)$, are normal.

(46) H. Furstenberg (1967) proved that if $p, q > 1$ are integers not both integer powers of the same integer (i.e. p and q are *multiplicatively independent*), then for every irrational α the sequence $p^m q^n \alpha \bmod 1$, $m, n = 1, 2, \dots$ is everywhere dense

in $[0, 1]$.

(47) B. Kra (1999) extended (46) to the following: Let p_i and q_i be integers and α_i real, $i = 1, 2, \dots, k$. If $p_1, q_1 > 1$ are multiplicatively independent, α_1 is irrational, and $(p_i, q_i) \neq (p_1, q_1)$ for $i > 1$ then the sequence $\sum_{i=1}^k p_i^m q_i^n \alpha_i \pmod{1}$, $m, n = 1, 2, \dots$ is dense in $[0, 1]$. He also gave the following:

(48) Let $p, q > 1$ be multiplicatively independent integers and let x_n , $n = 1, 2, \dots$, be any sequence of real numbers. Then for any irrational α the sequence $p^m q^n \alpha + x_n \pmod{1}$, $m, n = 1, 2, \dots$ is dense in $[0, 1]$.

(49) **Conjecture:** Let λ_i, μ_i , for $i = 1, 2, \dots, k$ be real algebraic numbers, $|\lambda_i|, |\mu_i| > 1$, λ_i, μ_i are multiplicatively independent, and $(\lambda_i, \mu_i) \neq (\lambda_j, \mu_j)$ for $i \neq j$. Then for any real numbers $\alpha_1, \dots, \alpha_k$ with at least one $\alpha_i \notin \mathbb{Q}(\cup_{i=1}^k \{\lambda_i, \mu_i\})$ the sequence $\sum_{i=1}^k \lambda_i^m \mu_i^n \alpha_i \pmod{1}$, $m, n = 1, 2, \dots$ is dense in $[0, 1]$.

(50) Conjecture (49) was stated by R. Urban (2007). He proved it for special algebraic integers of degree 2, see 1.28. As illustrating examples he gave:

For any α_1, α_2 with at least one non-zero, the sequence $\{(\sqrt{23}+1)^m (\sqrt{23}+2)^n \alpha_1 + (\sqrt{61}+1)^m (\sqrt{61}-6)^n \alpha_2\}$, $m, n = 1, 2, \dots$ is everywhere dense and also for irrational α_2 the sequence $\{(3+\sqrt{3})^m 2^n + 5^m 7^n \alpha_2 \sqrt{2}\}$, $m, n = 1, 2, \dots$, is everywhere dense in $[0, 1]$. For more information see Problem 1.28.

Entered by O. Strauch

S.D. ADHIKARI – P. RATH – N. SARADHA: *On the set of uniqueness of a distribution function of $\{\zeta(p/q)^n\}$* , Acta Arith. **119** (2005), no. 4, 307–316.

S. AKIYAMA – C. FROUGNY – J. SAKAROVITCH: *On number representation in a rational base*, (submitted).

A. AKIYAMA – Y. TANIGAWA: *Salem numbers and uniform distribution modulo 1*, Publ. Math. Debrecen **64** (2004), no. 3–4, 329–341 (MR 2058906).

D. BEREND – G. KOLESNIK: *Complete uniform distribution of some oscillating sequences*, J. Ramanujan Math. Soc. **26** (2011), no. 2, 127–144.

M.–J. BERTIN – A. DECOMPS–GUILLOUX – M. GRANDET–HUGOT – M. PATHIAUX–DELEFOSSE – J.–P. SCHREIBER: *Pisot and Salem numbers*, Birkhäuser Verlag, Basel, 1992 (MR 93k:11095).

D.G. CHAMPERNOWNE: *The construction of decimals normal in the scale ten*, J. London Math. Soc., **8** (1933), 254–260 (Zbl 7, 337).

G. CHOQUET: *Construction effective de suites $(k(3/2)^n)$. Étude des mesures $(3/2)$ -stables*, C.R. Acad. Sci. Paris, Ser. A–B **291** (1980), A69–A74 (MR 82h:10062a–d).

A.H. COPELAND – P. ERDŐS: *Note of normal numbers*, Bull. Amer. Math. Soc. **52** (1946), 857–860 (MR **8**, 194b).

H. DAVENPORT – P. ERDŐS: *Note on normal numbers*, Canad. J. Math. **4** (1952), 58–63 (MR **13**, 825).

[a] A. DUBICKAS: *On the distance from a rational power to the nearest integer*, J. Number Theory **117** (2006), 222–239.

[b] A. DUBICKAS: *Arithmetical properties of powers of algebraic numbers*, Bull. London Math. Soc. **38** (2006), no. 1, 70–80 (MR2201605=2006i:11080).

A. DUBICKAS: *On a sequence related to that of Thue–Morse and its applications*, Discrete Mathematics **307** (2007), no. 1, 1082–1093.

J. DUFRESNOY – CH. PISOT: *Sur un problème de M. Siegel relatif à un ensemble fermé d'entiers algébriques*, C. R. Acad. Sci. Paris **235** (1952), 1592–1593 (MR **14**, 538c).

J. DUFRESNOY – CH. PISOT: *Sur un point particulier de la solution d'un problème de M. Siegel*, C. R. Acad. Sci. Paris **236** (1953), 30–31 (MR **14**, 538d).

L. FLATTO – J.C. LAGARIAS – A.D. POLLINGTON: *On the range of fractional parts $\{\zeta(p/q)^n\}$* , Acta Arith. **70** (1995), no. 2, 125–147 (MR 96a:11073).

H. FURSTENBERG: *Disjointness in ergodic theory, minimal sets, and a problem in Diophantine*

- approximation*, Math. System Theory **1** (1967), 1–49.
- E. GHATE – E. HIRONAKA: *The arithmetic and geometry of Salem numbers*, Bull. Amer. Math. Soc. (N.S.) **38** (2001), no. 3, 293–314 (MR 2002c:11137).
- G.H. HARDY: *A problem of diophantine approximation*, Jour. Indian. Math. Soc. **11** (1919), 162–166.
- H. HELSON – J.-P. KAHANE: *A Fourier method in diophantine problems*, J. Analyse Math. **15** (1965), 245–262 (MR **31**#5856).
- J.F. KOKSMA: *Ein mengentheoretischer Satz ueber die Gleichverteilung modulo Eins*, Compositio Math. **2** (1935), 250–258 (Zbl 12, 14; JFM 61.0205.01).
- B. KRA: *A generalization of Furstenberg’s Diophantine theorem*, Proc. Amer. Math. Soc. **127** (1999), 1951–1956.
- D.H. LEHMER: *Factorization of certain cyclotomic functions*, Ann. Math. **34** (1933), 461–469 (Zbl 0007.19904).
- M.G. MADRITSCH – J.M. THUSWALDNER – R.F. TICHY: *Normality of numbers generated by the values of entire functions*, J. Number Theory **128** (2008), 1127–1145.
- K. MAHLER: *Arithmetische Eigenschaften einer Klasse von Dezimalbrüchen*, Nederl. Akad. Wetensch. Proc. Ser. A **40** (1937), 421–428 (Zbl 17, 56).
- K. MAHLER: *An unsolved problem on the powers of $3/2$* , J. Austral. Math. Soc. **8** (1968), 313–321 (MR **37**#2694).
- Y.-N. NAKAI – I. SHIOKAWA: *A class of normal numbers*, Japan. J. Math. (N.S.) **16** (1990), no. 1, 17–29 (MR 91g:11081).
- [a] Y.-N. NAKAI – I. SHIOKAWA: *A class of normal numbers. II*, in: *Number Theory and Cryptography* (Sydney, 1989), (J.H. Loxton Ed.), Cambridge University Press, Cambridge, London Math. Soc. Lecture Note Ser., Vol. 154, 1990, pp. 204–210 (MR 91h:11074).
- Y.-N. NAKAI – I. SHIOKAWA: *Discrepancy estimates for a class of normal numbers*, Acta Arith. **62** (1992), no. 3, 271–284 (MR 94a:11113).
- Y.-N. NAKAI – I. SHIOKAWA: *Normality of numbers generated by the values of polynomials at primes*, Acta Arith. **81** (1997), no. 4, 345–356 (MR 98h:11098).
- I.I. PJATECKIJ – ŠAPIRO (I.I. ŠAPIRO – PJATECKIJ): *On the laws of distribution of the fractional parts of an exponential function* (Russian), Izv. Akad. Nauk SSSR, Ser. Mat., Ser. Mat. **15** (1951), 47–52 (MR **13**, 213d).
- S.S. PILLAI: *On Waring’s problem*, Journal of Indian Math. Soc. (2) **2** (1936), 16–44; Errata *ibid.* p. 131 (JFM 62.1132.02).
- CH. PISOT: *Sur la répartition modulo 1 des puissances successives d’un même nombre*, C.R. Acad. Sci. Paris **204** (1937), 312–314 (Zbl 16, 53).
- [a] CH. PISOT: *Sur la répartition modulo 1*, C. R. Acad. Sci. Paris **204** (1937), 1853–1855 (Zbl 16, 392).
- CH. PISOT: *La répartition modulo 1 et les nombres algébriques*, (French), (Diss.), Paris, 1938 (Zbl 19, 7).
- CH. PISOT: *La répartition modulo 1 et les nombres algébriques*, Ann. Scuola norm. sup. Pisa, Sci. fis. mat. (2) **7** (1938), 205–248 (Identical with the previous item) (Zbl 19, 155).
- A.D. POLLINGTON: *Progressions arithmétiques généralisées et le problème des $(3/2)^n$* , C. R. Acad. Sci. Paris Sér. I Math. **292** (1981), 383–384.
- A.D. POLLINGTON: *Sur les suites $\{k\theta^n\}$* , C. R. Acad. Sci. Paris Sér. I Math. **296** (1983), no. 23, 941–943 (MR 86i:11034).
- L. RÉDEI: *Zu einem Approximationssatz von Koksma*, Math. Z. **48** (1942), 500–502 (MR **4**, 266c).
- R. SALEM: *A remarkable class of algebraic integers. Proof of a conjecture of Vijayaraghavan*, Duke Math. J. **11** (1944), 103–108 (MR **5**, 254a).
- C.L. SIEGEL: *Algebraic integers whose conjugates lie in the unit circle*, Duke Math. J. **11** (1944), 597–611 (MR **6**, 39b).
- O. STRAUCH: *On distribution functions of $\zeta(3/2)^n \bmod 1$* , Acta Arith. **81** (1997), no. 1, 25–35 (MR 98c:11075).
- O. STRAUCH: *Distribution of Sequences* (in Slovak), Mathematical Institute of the Slovak Academy of Sciences, (DSc Thesis), Bratislava, Slovakia, 1999.
- A. THUE: *Über eine Eigenschaft, die keine transcendente Größe haben kann*, Norske Vid. Skrift.

20 (1912), 1–15.

R. TIJDEMAN: *Note on Mahler's 3/2-problem*, Norske Vid. Selske. Skr. **16** (1972), 1–4 (Zbl 227.10025).

TOUFIK ZAÏMI: *An arithmetical property of powers of Salem numbers*, J. Number Theory **120** (2006), 179–191 (MR 2256803).

R. URBAN: *On density modulo 1 of some expressions containing algebraic integers*, Acta Arith. **127** (2007), no. 3 217–229.

[a] T. VIJAYARAGHAVAN: *On decimals of irrational numbers*, Proc. Indian Acad. Sci., Sect. A **12** (1940), 20 (MR **2**, 33d).

T. VIJAYARAGHAVAN: *On the fractional parts of the powers of a number. I*, J. London Math. Soc. **15** (1940), 159–160 (MR **2**, 33e).

[a] T. VIJAYARAGHAVAN: *On decimals of irrational numbers*, Proc. Indian Acad. Sci., Sect. A **12** (1940), 20 (MR **2**, 33d).

T. VIJAYARAGHAVAN: *On the fractional parts of the powers of a number (II)*, Proc. Cambridge Philos. Soc. **37** (1941), 349–357 (MR **3**, 274c).

H. WEYL: *Über die Gleichverteilung von Zahlen mod. Eins*, Math. Ann. **77** (1916), 313–352 (JFM 46.0278.06).

A. ZAME: *The distribution of sequences modulo 1*, Canad. J. Math. **19** (1967), 697–709 (MR **36**#115).

2.5 Duffin-Schaeffer conjecture and related sequences.

D.S.C.: Let $f(q)$ be a function defined on the positive integers and let $\varphi(q)$ be the Euler totient function. The Duffin and Schaeffer conjecture (D.S.C.) says that for an arbitrary function $f \geq 0$ defined on positive integers (zero values are also allowed for f) the diophantine inequality

$$\left| x - \frac{p}{q} \right| < f(q), \quad \gcd(p, q) = 1, \quad q > 0 \quad (1)$$

has infinitely many integer solutions p and q for almost all $x \in [0, 1]$ (in the sense of Lebesgue measure) if and only if the following series diverges $\sum_{q=1}^{\infty} \varphi(q)f(q) = \infty$.

NOTES:

(I) The D.S.C. is one of the most important unsolved problems in metric number theory, cf. Encyclopaedia of Mathematics 2000 (M. Hazewinkel ed.).

(II) It was inspired by A. Khintchine (1924) result: If $q^2 f(q)$ is nonincreasing and $\sum_{q=1}^{\infty} qf(q)$ diverges, then (1) has infinitely many integer solutions for almost all x . Originally, he did not assume $\gcd(p, q) = 1$.

(III) By the Borel-Cantelli lemma, (1) has only finitely many solutions for almost all x if $\sum_{q=1}^{\infty} \varphi(q)f(q)$ converges.

(IV) By the Gallagher ergodic theorem (P. Gallagher (1965)) the set of all $x \in [0, 1]$ for which (1) has infinitely many integer solutions has measure either 0 or 1.

(V) R.J. Duffin and A.C. Schaeffer (1941) improved Khintchine's theorem in (II) for $f(q)$ satisfying $\sum_{q=1}^Q qf(q) \leq c \sum_{q=1}^Q \varphi(q)f(q)$ for infinitely many Q and some positive constant c . They also have given an example of $f(q)$ such that $\sum_{q=1}^{\infty} qf(q)$ diverges, $\sum_{q=1}^{\infty} \varphi(q)f(q)$ converges and (1) has for almost all $x \in [0, 1]$ only finitely many solutions p and q , where the $\gcd(p, q) = 1$ is omitted (cf. (VI')). This naturally leads to D.S.C. with $\sum_{q=1}^{\infty} \varphi(q)f(q)$ replaced of $\sum_{q=1}^{\infty} qf(q)$.

(VI) In the following a class of sequences q_n $n = 1, 2, \dots$, distinct positive integers and a class of functions f is said to satisfy D.S.C. if the divergence $\sum_{n=1}^{\infty} \varphi(q_n) f(q_n)$ implies that for almost all $x \in [0, 1]$ there exist infinitely many n such that the diophantine inequality

$$\left| x - \frac{p}{q_n} \right| < f(q_n), \quad \gcd(p, q_n) = 1 \quad (2)$$

has an integer solution p . There are three types of results of q_n, f satisfying D.S.C.:

- (a) any one-to-one sequence q_n and special f ;
- (b) any $f \geq 0$ and a special q_n (e.g., $q_n = n^k$);
- (c) special q_n, f .

For example:

(VIa) Following f 's satisfy D.S.C. with every one-to-one sequence q_n :

- (i) $f(n) = \frac{c}{n^2}$, where $c > 0$ is a constant.
- (ii) $f(n) = O(n^{-2})$.
- (iii) $f(n) = O\left(\frac{\exp(g(n))^\gamma}{n^2}\right)$, where $\gamma = e^{\frac{1}{2}} - \varepsilon$, $\varepsilon > 0$ and $g(n)$ is the first positive integer for which $\sum_{\substack{p|n, p > g(n) \\ p\text{-prime}}} \frac{1}{p} < 1$.

Note that (i) was proved by P. Erdős (1970), (ii) by J.D. Vaaler (1978) and (iii) by V.T. Vil'chinskii (1979).

(VIb) Following one-to-one sequences q_n satisfy D.S.C. with every $f \geq 0$ (zero values are also allowed):

- (i) $\frac{\varphi(q_n)}{q_n} \geq c > 0$ for every n .
- (ii) $\frac{\varphi(q_n)}{\varphi(q_{n+1})} \leq c < 1$ for all sufficiently large n .
- (iii) $\sum_{i \neq j=1}^{\infty} \frac{4^{\omega(q_{ij})}}{\varphi(q_{ij})} < +\infty$, where $q_{ij} = \frac{q_i q_j}{\gcd(q_i, q_j)^2}$ and $\omega(n) = \#\{p\text{-prime}, p|n\}$.
- (iv) $\sum_{n=1}^{\infty} \frac{\varphi(q_n)}{q_n} < +\infty$.
- (v) $(q_m, q_n) = 1$ for every $m \neq n$.
- (vi) $\sum_{i, j=1}^{\infty} \frac{(\log q_{ij})^2}{q_{ij}} \frac{\varphi(q_i)}{q_i} \frac{\varphi(q_j)}{q_j} < +\infty$.
- (vii) $\sum_{n=1}^{\infty} \frac{(\log q_n)^2}{q_n^{2\varepsilon}} < +\infty$ and $d_{ij} \leq (q_i q_j)^{\frac{1}{2} - \varepsilon}$ for some $\varepsilon > 0$ and every $i \neq j$, where $d_{ij} = \gcd(q_i, q_j)$.
- (viii) The sequence $d_{ij} = \gcd(q_i, q_j)$, $i, j = 1, 2, \dots$, has only finitely many different terms.
- (ix) $\frac{q_n}{q_{n+1}} \leq c < 1$ for every n .
- (x) $\frac{\varphi(q_n)}{q_n} < K n^{-\delta}$ for some $K, \delta > 0$ and $n = 1, 2, \dots$.
- (xi) $q_n = n^k$, for $k \geq 2$.
- (xii) $q_n = q^n$, $q_n = n!$, $q_n = 2^{2^n} + 1$ -Fermat numbers, $q_n = F_n$ -Fibonacci numbers, $q_n = q^n - 1$, $q_n = q^n + 1$ (for every positive integer $q \geq 2$).
- (xiii) q_n is a one-to-one sequence of primes.

Note that (i) and (ii) can be found in Duffin and Schaeffer (1941).

(iii) and (iv) was proved by O. Strauch (1982, Th. 14 and 15).

(v) and (vi) by Strauch (1983, Th. 7 and 2).

(vii) by Strauch (1984, Th. 6).

(viii) by Strauch (1986, Th. 8).

(ix), (x), (xi) by G. Harman (1990, Th. 1).

(xii) by Strauch (1986).

(xiii) is an example in Duffin and Schaeffer (1941, p. 245). Also G. Herman (1998, p. 27, Cor. 2) notes that Duffin-Schaeffer criterion in (VIc(i)) (also in (VIb(i))) holds for one-to-one sequence q_n of primes, since it satisfies D.S.C. with every $f \geq 0$.

(VIc) The following special sequences q_n and functions f satisfy D.S.C.:

(i) $\sum_{n=1}^N q_n f(q_n) \leq c \sum_{n=1}^N \varphi(q_n) f(q_n)$ for $N = 1, 2, \dots$

(i') $q_n = n$ and $q_n^c f(q_n)$ is non-increasing. Here c may be any real constant.

(ii) $\sum_{n=1}^N \varphi(q_n) f(q_n) > cN^\delta$ for infinitely many N , where $c, \delta > 0$ are constants.

(iii) $f(q_n) > cq_n(\varphi(q_n)/q_n)^R$ for $n = 1, 2, \dots$, where $c, R > 0$ are constants.

(iv) q_n strictly increase, $q_n f(q_n)$ does not increase and the lower asymptotic density $\underline{d}(q_n) > 0$.

(v) $f(q_n)\varphi(q_n) = O((n \log n \log \log n)^{-1})$ and $c_1 n^A \leq q_n \leq c_2 n^B$, where c_1, c_2, A, B are positive constants and $B \geq 1$.

(vi) $\frac{\max_{i \leq n} \varphi(q_i)}{\sum_{i=1}^n \varphi(q_i)} \geq c > 0$ for $n = 1, 2, \dots$ and $f(q_n)$ is nonincreasing.

(vii) q_n is strictly increasing with $\underline{d}(q_n) > 0$ and $f(q) \geq cf(s)$ for all $q = 1, 2, \dots$, every $s \in \{q+1, q+2, \dots, 2q\}$ and some constant $c > 0$.

Note that (i) and (i') was proved by R.J. Duffin and A.C. Schaeffer (1942); (ii) and (iii) G. Harman (1990, 1998, p. 66, Th. 3.7); (iv) G. Harman (1998, p. 41, Cor. 3); (v) G. Harman (1998, p. 57, Th. 2.10); (vi) O. Strauch (1982); and (vii) E. Zoli (2008).

(VI') O. Strauch (1982): Let $p_n, n = 1, 2, \dots$ be the increasing sequence of all primes and put $q_n = p_1 p_2 \dots p_n$ and $f(q_n) = (q_n n \log n)^{-1}$. Then $\sum_{n=1}^{\infty} \varphi(q_n) f(q_n)$ converges, $\sum_{n=1}^{\infty} q_n f(q_n)$ diverges and for almost all $x \in [0, 1]$ the inequality (2) has only finitely many solutions p, q_n , but infinitely many if the assumption $\gcd(p, q_n) = 1$ is omitted.

(VI'') P.A. Caltin (1976) **conjectured** that the divergence $\sum_{q=1}^{\infty} \varphi(q) \max_{m \geq 1} f(m \cdot q)$

is the necessary and sufficient condition for (1) to have infinitely many solutions.

The D.S.C. implies this conjecture, cf. Harman (1998, pp. 28–29).

(VI*) It is interesting that one-dimensional D.S.C. is open, but multidimensional D.S.C. was proved by A.D. Pollington and R.C. Vaughan (1999) in the following form: For every $k = 2, 3, \dots$, for every one-to-one sequence $q_n, n = 1, 2, \dots$, of positive integers and for any nonnegative function f , if the series $\sum_{n=1}^{\infty} (\varphi(q_n) f(q_n))^k$ diverges, then for almost all $\mathbf{x} = (x_1, \dots, x_k)$ and for infinitely many n there exists an integer vector $\mathbf{p} = (p_1, \dots, p_k)$ such that the inequalities

$$\left| x - \frac{p_1}{q_n} \right| < f(q_n), \dots, \left| x - \frac{p_k}{q_n} \right| < f(q_n),$$

hold, where $\gcd(p_1 p_2 \dots p_k, q_n) = 1$.

(VI*b) G. Harman (1998, p. 65, Th. 3.6) proved another multidimensional D.S.C.:

Let $f_1(n), \dots, f_k(n)$ be functions of n taking values in $[0, c)$ for some $c > 0$. Write $\theta(n) = \prod_{j=1}^k (nf_j(n))$ and suppose, for some positive reals ε and K , that for each n for which $\theta(n) \neq 0$ we have $\max_{1 \leq j \leq k} \frac{\theta(n)}{nf_j(n)} \leq K(\theta(n))^\varepsilon$. Let q_n be a sequence of distinct positive integers for which

$$\sum_{n=1}^{\infty} (\varphi(q_n)f_1(q_n)) \dots (\varphi(q_n)f_k(q_n)) = \infty.$$

Then for almost all $\mathbf{x} = (x_1, \dots, x_k) \in [0, 1]^k$ there are infinitely many solutions

$$\left| x - \frac{p_1}{q_n} \right| < f_1(q_n), \dots, \left| x - \frac{p_k}{q_n} \right| < f_k(q_n), \quad \gcd(p_1 p_2 \dots p_k, q_n) = 1.$$

There are two following types of sequences inspired by D.S.C., namely eutaxic and quick.

Eutaxic sequences. Let $x_n \in [0, 1)$, $z_n \in \mathbb{R}^+$, $n = 1, 2, \dots$, be two sequences and $x \in [0, 1]$. Strauch (1994) introduced a new counting function

$$A(x; N; (x_n, z_n)) = \#\{n \leq N; |x - x_n| < z_n\}.$$

• The sequence x_n is said to be **eutaxic** if for every non-increasing sequence z_n the divergence of $\sum_{n=1}^{\infty} z_n$ implies that

$$\lim_{N \rightarrow \infty} A(x; N; (x_n, z_n)) = \infty$$

for almost all $x \in [0, 1]$. If furthermore

$$\lim_{N \rightarrow \infty} \frac{A(x; N; (x_n, z_n))}{2 \sum_{n=1}^N z_n} = 1$$

then x_n is called **strongly eutaxic**.

NOTES:

(VII) Eutaxic sequences were introduced by J. Lesca (1968). He proved that if θ is irrational then the sequence $n\theta \bmod 1$ is eutaxic if and only if θ has bounded partial quotients.

(VIII) M. Reversat proved the same for the strong eutaxicity of $n\theta \bmod 1$, i.e., for sequence $n\theta \bmod 1$ both notions coincide.

(IX) B. de Mathan (1971) defined the counting function

$$A^*(N, x_n) = \#\{0 \leq k < N; \exists n \leq N (x_n \in [k/N, (k+1)/N])\}$$

and proved that $\liminf_{N \rightarrow \infty} A^*(N, x_n)/N = 0$ implies that x_n is not eutaxic. Since for the sequence $x_n = n\theta \bmod 1$ and for θ with unbounded partial quotients we have $\liminf_{N \rightarrow \infty} A^*(N, x_n)/N = 0$, de Mathan (1971) recovered half of Lesca's

result.

(X) A characterization of strong eutaxicity in terms of L^2 discrepancy is an **open problem**, cf. O. Strauch (1994).

Quick sequences. Let $X = \cup_{m=1}^{\infty} I_m$ be a decomposition of an open set $X \subset [0, 1]$ into a sequence I_m , $m = 1, 2, \dots$, of pairwise disjoint open subintervals of $[0, 1]$ (empty intervals are allowed). Let x_n be an infinite sequence in $[0, 1)$. Define a new counting function

$$\begin{aligned} \tilde{A}(X; N; x_n) = \#\{m \in \mathbb{N}; \exists n \leq N \text{ such that } x_n \in I_m\} + \\ + \#\{n \leq N; x_n \notin X\}, \end{aligned}$$

i.e., if $x_n \in X$ for $n = 1, 2, \dots$, then $\tilde{A}(X; N; x_n)$ is the number of intervals I_m containing at least one element of x_1, x_2, \dots, x_N .

• The sequence x_n is said to be **quick** if for any open set $X \subset [0, 1]$ with the Lebesgue measure $|X| < 1$, there exists a constant $c = c(X)$ such that

$$\frac{\tilde{A}(X; N; x_n)}{N} \geq c > 0.$$

NOTES:

(X') Examples of quick sequences:

- (i) The sequence x_n of all dyadic rational numbers from $[0, 1]$ ordered by $\frac{0}{2}, \frac{1}{2}, \frac{1}{4}, \frac{3}{4}, \frac{1}{8}, \frac{3}{8}, \frac{5}{8}, \frac{7}{8}, \frac{1}{16}, \dots$
- (ii) The sequence x_n of all rational numbers (reduced fractions) from $(0, 1]$ ordered by $\frac{1}{1}, \frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{1}{4}, \frac{3}{4}, \frac{1}{5}, \frac{2}{5}, \frac{3}{5}, \frac{4}{5}, \frac{1}{6}, \dots$
- (iii) The sequence x_n of all non-reduced fractions from $(0, 1]$ ordered by $\frac{1}{1}, \frac{1}{2}, \frac{2}{2}, \frac{1}{3}, \frac{2}{3}, \frac{3}{3}, \frac{1}{4}, \frac{2}{4}, \frac{3}{4}, \frac{4}{4}, \frac{1}{5}, \dots$
- (iv) The denominators $1, 2, 3, \dots$ in (iii) can be replaced by any arithmetic subsequence of positive integers.

• The sequence x_n is said to be **uniformly quick** (abbreviated u.q.) if for any open set $X \subset [0, 1]$ we have

$$\lim_{N \rightarrow \infty} \frac{\tilde{A}(X; N; x_n)}{N} = 1 - |X|.$$

• If this limit holds for a special sequence of indices $N_1 < N_2 < \dots$, then x_n is said **almost u.q.**

NOTES:

(XI) Quick and u.q. sequences were introduced and studied by O. Strauch (1982, 1983, 1984, [a]1984, 1986) in connection with D.S.C.

(XII) Any quick sequence x_n is eutaxic, i.e., for every non-increasing sequence z_n , $\sum_{n=1}^{\infty} z_n = \infty$, for almost all $x \in [0, 1]$, the inequality $|x - y_n| < z_n$ holds for

infinitely many n .

(XII') Any u.q. sequence x_n is u.d. in $(0, 1]$ and it is also strongly eutaxic.

(XIII) The sequence $x_n = n\theta \bmod 1$ is u.q. if and only if the simple continued fraction expansion of the irrational θ has bounded partial quotients (cf. O. Strauch ([a]1984)).

(XIV) Strauch (1982, Th. 3): The u.d. sequence x_n is u.q. if for infinitely many M there exist c_M, c'_M , and $N_0(M)$ such that $c'_M \rightarrow 0$ as $M \rightarrow \infty$ and

$$\sum_{\substack{|x_i - x_j| \leq t \\ M < i \neq j \leq N}} 1 \leq c_M t (N - M)^2 + c'_M (N - M)$$

for every $N \geq N_0(M)$ and every $t \geq 0$.

(XV) The u.q. sequences x_n can be used in the numerical evaluation of integrals $\int_X f(x) dx$ over open subsets X of $[0, 1]$. Thus also for Jordan non-measurable sets X whose boundaries ∂X are of positive measure $|\partial X| > 0$, cf. Strauch (1997).

(XVI) Let $q_n, n = 1, 2, \dots$, be a one-to-one sequence of positive integers and let $A_n, n = 1, 2, \dots$ be a sequence composed from blocks

$$A_n = \left(\frac{1}{q_n}, \frac{a_2}{q_n}, \dots, \frac{a_{\varphi(q_n)}}{q_n} \right),$$

where $1 = a_1 < a_2 < a_3 < \dots < a_{\varphi(q_n)}$ are the integers $< q_n$ coprime to q_n . If $A_n, n = 1, 2, \dots$ is almost u.q. (with respect to the set of indices $N_n = \sum_{i=1}^n \varphi(q_i)$), then the D.S.C. holds for q_n and for every non-increasing $f(q_n)$. Thus the D.S.C. follows from the following conjecture immediately:

Conjecture: For every one-to-one sequence of positive integers q_n , the block sequence A_n is almost u.q.

(XVII) For every one-to-one sequence q_n in (VIb) the block sequence A_n is almost u.q., e.g., (v) relatively prime q_n ; (xii) $q_n =$ Fibonacci numbers; (ix) lacunary q_n ;

(XVIII) From (XIV) it follows immediately: Assume that, for infinitely many m there exist c_m and c'_m such that $c'_m \rightarrow 0$ as $m \rightarrow \infty$ and for every $0 < x < 1$ and for all sufficiently large n we have the estimation

$$\sum_{\substack{0 < \frac{a}{q_i} - \frac{b}{q_j} < x \\ (a, q_i) = (b, q_j) = 1 \\ m < i, j \leq n}} 1 \leq c_m x \left(\sum_{m < i \leq n} \phi(q_i) \right)^2 + c'_m \left(\sum_{m < i \leq n} \phi(q_i) \right). \quad (3)$$

Then the sequence $q_n, n = 1, 2, \dots$, satisfies D.S.C. with every non-increasing function f . If the estimation (3) holds for every permuted $q_{\pi(n)}$ and for any subsequences $q_{\pi(n_k)}, k = 1, 2, \dots$, then the sequence $q_n, n = 1, 2, \dots$, satisfies D.S.C. with every $f \geq 0$, zero values are also allowed. We **conjectured** that (3) holds for any one-to-one integers q_n , where the constants c_m and c'_m depend on q_n . All results in (VIb) can be found by proving (3), where we used the following

partial estimations: For given two integers q_i and q_j denote

$$a(x) = \prod_{\substack{p > x, p|a \\ p\text{-prime}}} p, \quad d_{ij} = \gcd(q_i, q_j), \quad q_{ij} = \frac{q_i q_j}{d_{ij}^2}, \quad x_{ij} = x d_{ij} q_{ij}.$$

Then the sum $\sum_{\substack{0 < \frac{a}{q_i} - \frac{b}{q_j} < x \\ (a, q_i) = (b, q_j) = 1}} 1$

has the following upper bounds:

- (i) $c_0 x \varphi(q_i) \varphi(q_j) \frac{q_{ij}(x_{ij})}{\varphi(q_{ij}(x_{ij}))}$;
- (ii) $c_0 x \varphi(q_i) \varphi(q_j)$;
- (iii) $c_0 x q_i q_j$;
- (iv) $c_0 x \varphi(q_i) \varphi(q_j) \frac{q_{ij}}{\varphi(q_{ij})}$;
- (v) $c_0 x \varphi(q_i) \varphi(q_j) + 2^{\omega(q_{ij})} \varphi(d_{ij})$;

where $\omega(n)$ is the number of distinct prime divisors of n and c_0 is an absolute constant.

(XIX) Let $g(x)$ be an integer polynomial. In generally, the D.S.C. for $q_n = g(n)$ is an **open problem** (we know only $g(x) = x^k$, see (VIb(xi))). In connection with (VIb(vii)) there is a **question**: When for polynomial $g(x)$ we have

$$\gcd(g(m), g(n)) \leq c(g(m)g(n))^{\frac{1}{2}-\varepsilon} \tag{4}$$

for every sufficiently large integers $m \neq n$, where the constants c and ε depend on $g(x)$.

A. Schinzel (personal communication) has shown that the Masser *abc*-hypothesis implies (4) for $g(x) = x^3 + k$, $k = 1, 2, \dots$, with exponent $\frac{1}{2} - \frac{1}{18} + \varepsilon'$. He conjectured that (4) holds also for $g(x) = x^k + 1$, $k = 3, 4, \dots$, but he proved that (4) does not hold for $g(x) = x^2 + 1$.

The problem of restricting both numerators p and denominators q in (1) to sets of number-theoretic interest was investigated by G. Harman (1988).

NOTES:

(XX) Firstly, he considers (1), where p, q are both primes. In this case D.S.C. has the following form.

Conjecture: For any function $f \geq 0$ if the sum

$$\sum_{\substack{q=2 \\ q\text{-prime}}}^{\infty} f(q) \frac{q}{\log q} \tag{5}$$

diverges, then for almost all x there are infinitely many primes p, q which satisfy (1).

(XXI) G. Harman (1988) established this D.S.C. for $f(q) \geq 0$ satisfying $0 < \sigma_1 \leq \frac{mf(m)}{nf(n)} \leq \sigma_2$ for all m with $n_0 \leq n < m < 2n$, where σ_1, σ_2, n_0 are positive

constants.

(XXII) V.T. Vil'chinskii (1990) replaced (5) by

$$\sum_{\substack{q=2 \\ q\text{-prime}}}^{\infty} f(q) \frac{q^{-m+1+(m/n)}}{\log q} \quad (6)$$

where integers m, n satisfy one from $m = n$, $m > 2n$, or $n > 2m$. He proved that for special $f > 0$, if the series (6) diverges, then for almost all x , there exists infinitely many primes p, q such that $\left| x - \frac{p}{q^m} \right| < f(q)$.

(XXIII) Using the theory of u.q. sequences, the D.S.C. for prime numbers and non-increasing f holds if the block sequence $A_n, n = 1, 2, \dots$,

$$A_n = \left(\frac{q_1}{q_n}, \frac{q_2}{q_n}, \dots, \frac{q_n}{q_n} \right),$$

is u.q., where $q_n, n = 1, 2, \dots$ is the increasing sequence of all primes. Note that the sequence of blocks $A_n, n = 1, 2, \dots$, is u.d., see 1.9 Block sequence, Example (I).

(XXIV) G. Harman (1998, p. 168, Th. 6.2): Let A and B be sets of positive integers and denote

- (i) $\underline{d}(B) > 0$;
- (ii) $A(kn)/A(n) > c + 1$ for all n and for some constants $k > 1, c > 0$. Here $A(n) = \#\{i \leq n; i \in A\}$;
- (iii) $A(2n) - A(n) < C$ for all n , where C depends only on A ,
- (iv) $\lim_{q \rightarrow \infty} \lim_{p \rightarrow \infty} \gcd(p, q) = \infty$, where $q \in A$ and $p \in B$;
- (v) $f(n)$ is non-increasing;
- (vi) $0 < \sigma_1 < \frac{mf(m)}{nf(n)} < \sigma_2$ for all m with $n_0 \leq n < m < 2n$, where σ_1, σ_2, n_0 are positive constants;
- (vii) $\liminf_{n \rightarrow \infty} \frac{1}{A(n)} \sum_{\substack{q=1 \\ q \in A}}^n \frac{1}{q} \sum_{\substack{p \in B \\ p/q \in [x, y] \\ \gcd(p, q) = 1}} 1 > c(y - x)$ for all intervals $[x, y]$, where

$c > 0$ is a constant depending on A and B .

Suppose that (i), (vii) and at least one of (ii), (iii), (iv) holds for A and at least one of (v), (vi) holds for f . If the series $\sum_{q=1, q \in A}^{\infty} f(q)$ diverges, then there are infinitely many solutions to

$$\left| x - \frac{p}{q} \right| < f(q), \quad q \in A, \quad p \in B, \quad \gcd(p, q) = 1 \quad (7)$$

for almost all x .

- $\mathcal{K}(f) = \left\{ x \in \mathbb{R}; \left| x - \frac{p}{q} \right| < f(q) \text{ for infinitely many rationals } \frac{p}{q} \right\}$;
- $\text{Exact}(f) = \mathcal{K}(f) - \cup_{m \geq 2} \mathcal{K}((1 - 1/m)f)$;

(XXV) Y. Bugeaud (2008) collected known results:

- (i) If $f(q)$ is non-increasing and $f(q) = o(q^{-2})$ then $\mathcal{K}(f) \neq \emptyset$ (V. Jarník (1931));
(ii) If $q^2 f(q)$ is non-increasing and $\sum_{q=1}^{\infty} f(q)$ converges then for Hausdorff dimension $\dim \text{Exact}(f) = \dim \mathcal{K}(f) = \frac{2}{\lambda}$, where $\lambda = \liminf_{x \rightarrow \infty} -\frac{\log f(x)}{\log x}$ (M.M. Dodson (1992)).

Since for convergent $\sum_{q=1}^{\infty} f(q)$ the result (i) is very satisfactory, Y. Bugeaud (2008) proposed two following problems:

Open problem 1. Let $f(q)$ be a non-increasing, $f(q) = o(q^{-2})$ and $\sum_{q=1}^{\infty} f(q)$ diverges then to find Hausdorff dimension of $\text{Exact}(f)$.

- (iii) Y. Bugeaud (2008): Let $q^2 f(q)$ be a non-increasing, $\sum_{q=1}^{\infty} f(q)$ diverges and $1/(q^{2+\varepsilon}) \leq f(q) \leq 1/(100q^2 \log q)$ for any $\varepsilon > 0$ and sufficiently large q . Then $\dim \text{Exact}(f) = \dim {}_H\mathcal{K}(f) = 1$.

Open problem 2. Study the set $\text{Exact}(c/n^2)$.

- (iv) Y. Bugeaud (2008): For any $0 < c < 1/6$ the set $\text{Exact}(c/n^2)$ is non-empty. (XXVI) D. Berend and A. Dubickas (2009) studied diophantine approximation in the form (VII) $|x - x_n| < z_n$. Putting

$$\bullet \mathcal{G}(x_n, z_n) = \{x \in \mathbb{R}; |x - x_n| < z_n \text{ for infinitely many } n\}$$

they proved:

- (i) Let x_n be an arbitrary dense sequence in $[0, 1]$, and z_n be an arbitrary sequence of positive numbers. Then the set $\mathcal{G}(x_n, z_n)$ is an uncountable dense subset of the interval $[0, 1]$.

- (ii) Let x_n be an arbitrary sequence in $[0, 1]$, and z_n be an arbitrary sequence of positive numbers. If $\sum_{n=1}^{\infty} z_n^s < \infty$ for some $0 < s < 1$, then $\dim_H \mathcal{G}(x_n, z_n) \leq s$ (the Hausdorff dimension).

- (iii) For any sequence $z_n > 0$, $\lim_{n \rightarrow \infty} z_n = 0$, there exists well distributed $x_n \in [0, 1)$ such that $\dim_H \mathcal{G}(x_n, z_n) = 0$.

- (iv) For every z_n , $\sum_{n=1}^{\infty} z_n = \infty$, there exists $x_n \in [0, 1]$ such that $\mathcal{G}(x_n, z_n) = [0, 1]$.

- (v) Note that, by definition of eutaxic sequence and by (VII), for $x_n = n\theta \bmod 1$, if θ has bounded partial quotients, then the measure $|\mathcal{G}(x_n, z_n)| = 1$, for an arbitrary sequence $z_n > 0$, $\sum_{n=1}^{\infty} z_n = \infty$.

(XXVII) L. Mišík and O. Strauch (2012) are linked to (XXVI) results:

- (i) Let x_n be a sequence in $[0, 1)$ such that the set $G(x_n)$ of all d.f.s of x_n contains only continuous d.f.s. Then for every sequence $z_n > 0$, $z_n \rightarrow 0$, and every $x \in [0, 1]$ we have: If $|x - x_{n_k}| < z_{n_k}$, $k = 1, 2, \dots$, then the asymptotic density $d(n_k) = 0$.

- (ii) Ex.: The set $G(\log n)$ (was found by A. Wintner (1935)) has only continuous functions. Thus if $|x - \{\log n_k\}| < z_{n_k}$, $k = 1, 2, \dots$ then $\frac{k}{n_k} \rightarrow 0$ for every sequence $z_n > 0$, $z_n \rightarrow 0$. Note that, recently Y. Ohkubo (2011) proved: Let p_n , $n = 1, 2, \dots$, be the increasing sequence of all primes. The sequence $\{\log p_n\}$, $n = 1, 2, \dots$, has the same d.f.s as $\log n \bmod 1$. Thus n_k with $|x - \{\log p_{n_k}\}| < z_{n_k}$, $k = 1, 2, \dots$ satisfies $\frac{k}{n_k} \rightarrow 0$, again.

- G. Myerson (1993)(see [SP, 1.8.10]) introduced the sequence x_n in $[0, 1)$ to be *uniformly maldistributed* if for every subinterval $I \subset [0, 1)$ with positive length $|I| > 0$ we have both $\liminf_{n \rightarrow \infty} \frac{\#\{i \leq n; x_i \in I\}}{n} = 0$, $\limsup_{n \rightarrow \infty} \frac{\#\{i \leq n; x_i \in I\}}{n} = 1$.

(iii) Let x_n be a uniformly maldistributed sequence in $[0, 1]$. Then there exists a decreasing sequence $z_n > 0$, $n = 1, 2, \dots$, $z_n \rightarrow 0$, such that for every $x \in [0, 1]$, the sequence of all indices n_k , $|x - x_{n_k}| < z_{n_k}$, $k = 1, 2, \dots$, has the upper asymptotic density $\bar{d}(n_k) = 1$.

(iv) Ex.: The sequence $\{\log \log n\}$, $n = 2, 3, \dots$, is uniformly maldistributed, thus there exists a decreasing sequence $z_n > 0$, $n = 1, 2, \dots$, $z_n \rightarrow 0$, such that for every $x \in [0, 1]$, $\bar{d}(n_k) = 1$ for all possible n_k , $|x - \{\log \log n_k\}| < z_{n_k}$.

(XXVIII) For Hausdorff dimension the analogue of the D.S.C. is true. G. Harman (1998, Th. 10,7) proved that if $\sum_{q=1}^{\infty} qf(q) = \infty$, then the set

$X = \{x \in [0, 1]; \text{ there exists infinitely many integer solutions } |x - \frac{p}{q}|, \gcd(p, q) = 1, q > 0\}$ has Hausdorff dimension 1.

(Note that $\sum_{q=1}^{\infty} \varphi(q)f(q) = \infty \Rightarrow \sum_{q=1}^{\infty} qf(q) = \infty$). The dimensional analogue of D.S.C. also proved A. Haynes, A. Pollington and S. Velani (2012).

(XXIX) O. Strauch (1983) also studied the diophantine inequality

$$|x - x_n| < z_n/z. \quad (8)$$

He proved

(i) If x_n is dense in $[0, 1]$, $z_n \rightarrow 0$, the following sets

$X_1 = \{x \in [0, 1]; \text{ for every } z > 0, (8) \text{ and } x_n > x \text{ holds for infinitely many } n\}$,

$X_2 = \{x \in [0, 1]; \text{ for every } z > 0, (8) \text{ and } x_n < x \text{ holds for infinitely many } n\}$,

are also dense and they have a power c , and

$X_0 = \{x \in [0, 1]; \text{ there exists } z > 0, (8) \text{ holds only for finitely many } n\}$

is of the first category.

(ii) For every $x_n \in [0, 1]$, $z_n \rightarrow 0$, with the possible exception of a nullset, the unit interval $[0, 1]$ can be decomposed into two sets:

$X_3 = \{x \in [0, 1]; \text{ for every } z > 0, (8) \text{ holds only for finitely many } n\}$,

$X_4 = \{x \in [0, 1]; \text{ for every } z > 0, (8) \text{ and } x_n > x \text{ hold for infinitely many } n$

and also (8) and $x_n < x$ hold for infinitely many $n\}$.

(XXX) A. Haynes, A. Pollington and S. Velani (2012) replace Duffin-Schaeffer series $\sum_{q=1}^{\infty} \varphi(q)f(q)$ by series $\sum_{q=1}^{\infty} \varphi(q)(f(q))^{1+\varepsilon}$ and prove that the divergence $\sum_{q=1}^{\infty} \varphi(q)(f(q))^{1+\varepsilon} = \infty$ implies that the diophantine inequality (1) has infinitely many integer solutions p and q for almost all $x \in [0, 1]$ and for arbitrary $f(q) \geq 0$. Thus D.S.C. holds in this form. Liangpan Li (2013) replace $\sum_{q=1}^{\infty} \varphi(q)(f(q))^{1+\varepsilon}$ by $\sum_{q=1}^{\infty} \varphi(q)q^\varepsilon(f(q))^{1+\varepsilon}$ in this result.

Entered by O. Strauch

D. BEREND – A. DUBICKAS: *Good points for diophantine approximation*, Proc. Indian Acad. Sci. (Math. Sci.) **119** (2009), no. 4, 423–429.

P.A. CALTIN: *Two problems in metric Diophantine approximation I*, J. Number Theory **8** (1976), 289–297.

R.J. DUFFIN AND A.C. SCHAEFFER: *Khintchine's problem in metric diophantine approximation*, Duke Math. J. **8** (1941), 243–255 (MR **3**, 71).

P. GALLAGHER: *Metric simultaneous diophantine approximation. II*, Mathematika **12** (1965), 123–127.

- G. HARMAN: *Some cases of the Duffin and Schaeffer conjecture*, Quart. J. Math. Oxford Ser.(2) **41** (1990), no. 2, 395–404 (MR 92c:11073).
- G. HARMAN: *Metric diophantine approximation with two restricted variables. III. Two prime numbers*, J. Number Theory **29** (1988), 364–375.
- G. HARMAN: *Metric Number Theory*, London Math. Soc. Monographs, New Series, Vol. 18, Clarendon Press, Oxford, 1998 (MR 99k:11112).
- A.K. HAYNES – A.D. POLLINGTON – S.L. VELANI: *The Duffin-Schaeffer Conjecture with extra divergence*, Math. Ann. **353** (2012), 259–273.
- A. KHINTCHINE (A.J. CHINČIN): *Ein Satz über Kettenbrüche, mit arithmetischen Anwendungen*, Math. Z. **18** (1923), 289–306 (JFM 49.0159.03).
- B. DE MATHAN: *Un critère de non-eutaxie*, C. R. Acad. Sci. Paris Sér. A–B **273** (1971), A433–A436 (MR 44#6610; Zbl 0219.10061).
- J. LESCA: *Sur les approximationes a'une dimension*, Univ. Grenoble, (Thèse Sc. Math.), Grenoble, 1968.
- LIANGPAN LI: *A note on the Duffin-Schaeffer conjecture*, Uniform Distribution Theory **8** (2013), no. 2, 151–156.
- L. MIŠÍK – O. STRAUCH: *Diophantine approximation generalized*, Proceedings of the Steklov Institute of Mathematics **276** (2012), 193–207.
- MYERSON, G.: *A sampler of recent developments in the distribution of sequences*, in: Number theory with an emphasis on the Markoff spectrum (Provo, UT, 1991), Lecture Notes in Pure and App. Math. **147** Marcel Dekker, New York, 1993, 163–190.
- OHKUBO, Y.: *On sequences involving primes*, Uniform Distribution Theory **6** (2011), no. 2, 221–238.
- A.D. POLLINGTON – R.C. VAUGHAN: *The k -dimensional Duffin and Schaeffer conjecture*, Matematika **37** no. 2, (1990), 190–200 (MR 92d:11084).
- M. REVERSAT: *Un résultat de forte eutaxie*, C. R. Acad. Sci. Paris Sér. A–B **280** (1975), A153–A155 (MR 51#3075).
- V.G. SPRINDZUK: *Metric Theory of Diophantine Approximations*, (Russian), Izd. Nauka, Moscow, 1977 (English translation by R. A. Silverma, Winston/Wiley, Washington, DC, 1979).
- STRAUCH, O.: *A coherence between the diophantine approximations and the Dini derivatives of some real functions*, Acta Math. Univ. Comenian. **42–43** (1983), 97–109 (MR 86c:11021).
- O. STRAUCH: *L^2 discrepancy*, Math. Slovaca **44** (1994), 601–632 (MR 96c:11085).
- O. STRAUCH: *Duffin – Schaeffer conjecture and some new types of real sequences*, Acta Math. Univ. Comenian. **40–41** (1982), 233–265 (MR 84f:10065).
- O. STRAUCH: *Some new criteria for sequences which satisfy Duffin – Schaeffer conjecture, I*, Acta Math. Univ. Comenian. **42–43** (1983), 87–95 (MR 86a:11031).
- O. STRAUCH: *Some new criteria for sequences which satisfy Duffin – Schaeffer conjecture, II*, Acta Math. Univ. Comenian. **44–45** (1984), 55–65 (MR 86d:11059).
- [a] O. STRAUCH: *Two properties of the sequence $n\alpha \pmod{1}$* , Acta Math. Univ. Comenian. **44–45** (1984), 67–73 (MR 86d:11057).
- O. STRAUCH: *Some new criteria for sequences which satisfy Duffin – Schaeffer conjecture, III*, Acta Math. Univ. Comenian. **48–49** (1986), 37–50 (MR 88h:11053).
- O. STRAUCH: *A numerical integration method employing the Fibonacci numbers*, Grazer Math. Ber. **333** (1997), 19–33 (MR 99h:65038).
- O. STRAUCH: *Distribution of Sequences* (in Slovak), Mathematical Institute of the Slovak Academy of Sciences, (DSc Thesis), Bratislava, Slovakia, 1999.
- O. STRAUCH: *Duffin-Schaeffer conjecture; Gallagher ergodic theorem*, in: *Encyclopaedia of Mathematics, Supplement II*, (M. Hazewinkel Eds.), Kluwer Academic Publishers, Dordrecht, 2000, pp. 172–174, 242–243.
- V.T. VIL'CHINSKIĪ: *On the metric theory of nonlinear diophantine approximation*, (Russian), Dokl. Akad. Nauk. BSSR **34** (1990), no. 8, 677–680.
- WINTNER, A.: *On the cyclical distribution of the logarithms of the prime numbers*, Quart. J. Math. Oxford (1) **6** (1935), 65–68 (Zbl 11, 149).
- E. ZOLI: *A theorem of Khintchine type*, Uniform Distribution Theory **3** (2008), no. 1, 73–83.
- E. ZOLI: *Addendum to: A theorem of Khintchine type*, Uniform Distribution Theory **3** (2008), no. 1, 153–155